

to MTL group for all genes. Upregulation for BCRP, MRP1 and UGT1A4 were statistically significant for Control versus FCD groups ( $p < 0.05$ ). Further studies on bigger cohort of patients are required to conclude these findings.

<http://dx.doi.org/10.1016/j.ijep.2015.12.017>

### Endoscopic assisted inter hemispheric trans-callosal hemispherotomy: Preliminary description of a novel technique

P. Sarat Chandra\*, Ajay Garg, Chitra Sarkar, Nilesh Kurwale, Sarabjit Singh Chibber, Madhavi Tripathi, C.S. Bal, Manjari Tripathi

Dept of Neurosurgery and Center of Excellence  
AIIMS, New Delhi, India

E-mail address: [saratpchandra3@gmail.com](mailto:saratpchandra3@gmail.com)  
(P.S. Chandra).

**Background:** Various hemispherotomy techniques have been developed to reduce complication rates and achieve the best possible seizure control.

**Objective:** We present a novel and minimally invasive endoscopic assisted approach to perform this procedure.

**Method:** Endoscopic assisted inter hemispheric trans-callosal hemispherotomy was performed in 20 children (April 2013–June 2014). The procedure consisted of the use of a small craniotomy (4 × 3 cm), just lateral to midline using a transverse skin incision. Following dural opening, the surgery was performed with the assistance of a rigid high-definition endoscope, bayoneted self-irrigating bipolar, and other standard endoscopic instruments. Steps included a complete corpus callosotomy followed by the disconnection of the hemisphere at the level of the basal nuclei and thalamus. The surgeries were performed in a dedicated operating room with intra-operative MRI and neuro navigation. Intra-operative MRI confirmed a total disconnection.

**Results:** The pathologies for which surgeries were performed included sequelae of middle cerebral artery infarct (8), Rasmussen's (4), and hemimegalencephaly (8). Four patients had a class I Engel and one patient had a class II outcome at a mean follow up of 10.2 months (range: 3–14 months). The mean blood loss was 80 cc and mean operating time was 220 min. There were no complications in this study.

**Conclusion:** The present study describes a pilot novel technique and the feasibility of performing a minimally invasive endoscopic assisted hemispherotomy.

<http://dx.doi.org/10.1016/j.ijep.2015.12.018>



### Endoscopic assisted (through a mini craniotomy) corpus callostomy combined with anterior, hippocampal, and posterior commissurotomy in Lennox Gastaut syndrome: A pilot study to establish its safety and efficacy

P. Sarat Chandra\*, Ajay Garg, Chitra Sarkar, Nilesh Kurwale, Sarabjit Singh Chibber, Madhavi Tripathi, C.S. Bal, Manjari Tripathi

Dept of Neurosurgery and Center of Excellence  
AIIMS, New Delhi, India

E-mail address: [saratpchandra3@gmail.com](mailto:saratpchandra3@gmail.com)  
(P.S. Chandra).

**Background:** Corpus callosotomy (CC) is a palliative procedure especially for Lennox Gastaut semiology without localization with drop attacks.

**Objective:** To describe endoscopic assisted complete CC combined with anterior, hippocampal and posterior commissurotomy.

**Methods:** Patients with drug refractory epilepsy (DRE) having drop attacks as predominant seizure type, bilateral abnormalities on imaging, moderate to severe mental retardation. All underwent a complete work up (including MRI).

**Results:** Patients ( $n = 16$ , mean age  $11.4 \pm 6.4$  years, range 6–19 years) mean seizure frequency:  $24.5 \pm 19.8$ /days (range 1–60); mean intelligence quotient:  $25.23 \pm 10.71$ . All had syndromic diagnosis of Lennox Gastaut syndrome (LGS), with etiologies: hypoxic insult (10), lissencephaly (2), bilateral band heterotropia (2) microgyria and pachygyria (2). Surgery: complete callosotomy and section of anterior and posterior commissure by microscopic approach through a mini craniotomy (11) and endoscopic assisted approach (5). Complications: meningitis (1) hyperammonemic encephalopathy (2) and acute transient disconnection (5). No mortality or long-term morbidity. Mean follow-up:  $18 \pm 4.7$  months (range 16–27 months). Drop attacks stopped in all. Seizure frequency/duration decreased >90% in 10 and >50% in 5, increased in one patient. All patients attained pre-surgical functional levels in 3–6 months. Child behavior checklist scores: no deterioration. Parental questionnaire reported 90% satisfaction, attributing to control of drop attacks. The series was compared retrospectively with age/sex matched cohort (where a callosotomy only was performed), showed better outcome for drop attacks ( $p < 0.003$ ).

**Conclusion:** This preliminary study demonstrated efficacy, safety of complete callosotomy with anterior, hippocampal and posterior commissurotomy in LGS (drop attacks) with moderate-severe mental retardation.

<http://dx.doi.org/10.1016/j.ijep.2015.12.019>

