

Lesson learned from early experience in pediatric epilepsy surgery



Wihasto Suryaningtyas, Heri Subiyanto, Muhammad Arifin

Section on Pediatric, Department of Neurosurgery, Faculty of Medicine, Airlangga University, Soetomo General Hospital, Surabaya, Indonesia

About 1% of children population may have seizure that is recurrent and occurred without provocation, known as epilepsy. Those children with epilepsy fall into two main categories: well-controlled with drug(s) and uncontrollable with medication. The latest is now called intractable epilepsy or medically refractory epilepsy or drug resistant epilepsy.

For intractable epilepsy, pediatric epilepsy surgery has grown as a treatment of choice in carefully selected candidates. Decades ago, surgical avenue for epilepsy was considered as a last resort in medically refractory focal epilepsy after failure of numerous antiepileptic drug trials spanning many years. There have been certain key developments that have catalyzed this paradigm to change. Now, epilepsy surgery for children is considered earlier.

Sixteen cases of pediatric epilepsy surgery will be presented here along with type of pathology, features, and syndromic cases, and outcome during follow up. Cortical dysplasia was the most frequent pathology encountered including gyral, lobar and hemispheric type. Family of syndromic epilepsy such as West syndrome was also found. It is obvious that epilepsy in childhood age has many different features compared to their adult counterpart. The challenge is increasing when there is no sophisticated facilities available, and the physicians have to rely on their capacity to diagnose, analyze, and make the decision whether he/she is the best candidate for surgery or not.

<http://dx.doi.org/10.1016/j.ijep.2015.12.049>

Tailored temporal lobectomy for mesial temporal lobe epilepsy: Can we minimize visual field defect? Result of a prospective study



Xian-Lun Zhu, on behalf of the Epilepsy Surgery Group

Prince of Wales Hospital, Chinese University of Hong Kong, Hong Kong

Background: Traditional anterior temporal lobectomy for mesial temporal epilepsy is associated with high chance of visual field (VF) defect ranged from 50% to 100%.¹⁻⁶ A review in 2008 of our own series showed 68% upper quadrant VF defect contralateral to the side of operation (unpublished). Our further study on the distance between the anterior tip of Myer's lobe and the temporal lobe pole using Diffusion Tensor Tractography (DTT) among the Southern Chinese population showed significant individual variations ranged from 15 to 17 mm.⁷ We hypothesized that tailored anterior temporal lobectomy could minimize the VF defect. We have carried out a prospective study comparing with the historical series.

Materials and methods: A prospective study for all epilepsy surgery cases requiring resection of mesial temporal structure from 2009 till 2015. The Myer's loop was delineated using DTT. Navigation was used to guide the anterior temporal lobectomy. All cases had pre-operative and post-operative VF perimetry done.

Results: The prospective cohort (study group) composed of 25 cases. 3 cases were excluded from VF analysis because pre-existing VF defect. Among the 23 cases, new VF defect was found in 8 cases (32%). Comparing with the historical group, the reduction of VF defect was very close to significance ($P = 0.052$). Seizure outcome of Engel classification I were 73% and 69% for the study and historical group respectively.

Conclusion: Using DTT to delineate the Myer's loop and to operate under navigation guidance, tailored temporal lobectomy for mesial temporal resection may reduce the chance of VF defect significantly post-operatively without compromise seizure free outcome.

REFERENCES

1. Marino P, Rasmussen T. Visual field changes after temporal lobectomy in man. *Neurology*. 1968;18:825-835.
2. Katz A, Awad I, Kong AK, et al. Extent of resection in temporal lobectomy for epilepsy. II. Memory changes and neurologic complications. *Epilepsia*. 1989;30:763-771.
3. Tecoma E, Laxer K, Barbaro N, et al. Frequency and characteristics of visual field deficits after surgery for mesial temporal sclerosis. *Neurology*. 1993;43:1235-1238.
4. Bjork A, Kugelberg E. Visual field defects after temporal lobectomy. *Acta Ophthalmol*. 1957;35:210-216.
5. Falconer M, Wilson J. Visual field changes following anterior temporal lobectomy: their significance in relation to Meyer's loop. *Brain*. 1958;81:1-14.
6. Hughes T, Abou-Khalil B, Lavin P, et al. Visual field defects after temporal lobe resection. A prospective quantitative analysis. *Neurology*. 1999;53:167-172.
7. Wang YX, Zhu XL, Siu D, et al. The use of diffusion tensor tractography to measure the distance between the anterior tip of the Meyer loop and the temporal pole in a cohort from Southern China. *J Neurosurg*. 2010;113:1144-1151.

<http://dx.doi.org/10.1016/j.ijep.2015.12.050>

Accurate surgical treatment for rolandic epilepsy



Zhang Xinwei

Neurosurgical Department, The 3rd Affiliated Hospital of Southern Medical University, Southern Research Institute Against Epilepsy, Guangzhou 516030, China

Objective: Study the method of operative for rolandic epilepsy.

Method: We collected subjects who underwent epilepsy surgeries consecutively in southern research institute against epilepsy with the epileptogenic zone located in rolandic areas from January 2008 to March 2015. We analysed the method of location, technology of resection and follow-up result. The epileptogenic zone located by clinical symptoms, MR,