

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

International Journal of Epilepsy

journal homepage: <http://www.journals.elsevier.com/international-journal-of-epilepsy>




Image corner

Focal gigantism in tuberous sclerosis

Sanat R. Bhatkar*, Aastha Takkar

Department of Neurology, Postgraduate Institute of Medical Research (PGIMER), Chandigarh, 160012, India



ARTICLE INFO

Article history:

Received 19 October 2016

Accepted 21 February 2017

Available online 17 March 2017

Keywords:

Tuberous sclerosis

Focal gigantism

ABSTRACT

A 38 year lady had recurrent generalized and myoclonic seizures since 6 years of age. On examination she had facial adenoma sebaceum and periungual fibromas, cutaneous angiofibromas on the back. Contrast enhanced magnetic Resonance Imaging of brain was suggestive of multiple cortical tubers with subependymal nodules consistent with the diagnosis of Tuberous sclerosis. Focal hypertrophy of the skin of the dorsal aspect of the left hand with focal gigantism was noted.

© 2017 Indian Epilepsy Society. Published by Elsevier, a division of RELX India, Pvt. Ltd. All rights reserved.

A 38 year lady had recurrent generalized and myoclonic seizures since 6 years of age. On examination she had facial adenoma sebaceum (Fig. 1a) and periungual fibromas, cutaneous angiofibromas on the back. Contrast enhanced magnetic Resonance Imaging of brain was suggestive of multiple cortical tubers with subependymal nodules consistent with the diagnosis of Tuberous sclerosis. Focal hypertrophy of the skin of the dorsal aspect of the left hand with **focal gigantism** was noted (Fig. 1b).

Tuberous sclerosis complex in an autosomal dominant neurocutaneous disorder frequently recognized by presence of facial angiofibromas, periungual fibromas, shagreen patches; and presence of cortical tubers and subependymal nodules in MRI Brain.¹ Focal gigantism in the form of digital enlargement is an uncommon musculoskeletal association in tuberous sclerosis. The mechanism for its occurrence is unknown and it could be implicated to be due to localized tissue hyperplasia.²



Fig. 1. Showing patient of tuberous sclerosis with adenoma sebaceum (Fig. 1a) and focal gigantism and localized digital enlargement (Fig. 1b).

* Corresponding author.

E-mail addresses: bhatkar_sr@yahoo.com (S.R. Bhatkar), draastha49@yahoo.com (A. Takkar).

Conflicts of interest

None.

Financial disclosures/funding source

None.

Author contribution

Sanat R Bhatkar Drafting the article or revising it intellectual content, analysis or interpretation of the data, manuscript preparation.

Aastha Takkar Drafting the article and revising it's intellectual content. Manuscript preparation.

References

1. Northrup H, Krueger DA, on behalf of the International Tuberous Sclerosis Complex Consensus Group. Tuberous sclerosis complex diagnostic criteria update: recommendations of the 2012 international tuberous sclerosis complex consensus conference. *Pediatr Neurol.* 2013;49(4):243–254.
2. Soeiro E Sá M, Moldovan O, Sousa AB. Macrodactyly in tuberous sclerosis complex: case report and review of the literature. *Am J Med Genet A.* 2016;170 (July (7)):1903–1907.