


## Reintervention for stent occlusion after endoscopic ultrasound-guided hepaticogastrostomy with novel use of a precut needle-knife

Endoscopic ultrasound-guided hepaticogastrostomy (EUS-HGS) has gained popularity as an alternative biliary drainage method [1, 2]; however, reintervention after EUS-HGS remains to be elucidated. In EUS-HGS, use of a biliary stent that is longer than 100 mm is recommended in order to prevent stent migration [2, 3]. However, such stent placement occasionally makes reintervention challenging owing to the long length of the stent in the gastric lumen. A few reports have described technical efforts involved in reintervention after EUS-HGS [4, 5]. We describe a patient who underwent successful reintervention via a novel use of a precut needle-knife.

A 74-year-old woman with recurrent pancreatic cancer after pancreaticoduodenectomy presented with recurrent cholangitis. An 8 × 100 mm covered metal stent (Niti-S biliary covered stent; Taewoong Medical, Seoul, South Korea) had been previously deployed during EUS-HGS for biliary obstruction at the hepatic hilum. Stent occlusion occurred 4 months after EUS-HGS. Abdominal computed tomography showed a dilated intrahepatic bile duct, and stent occlusion was confirmed on endoscopy (► Fig. 1). Revisionary stent placement was attempted.

First, the advancement of an endoscopic retrograde cholangiopancreatography (ERCP) catheter was attempted via the proximal end of the HGS stent; however, the long stent length in the gastric lumen rendered catheter insertion impossible. Therefore, reintervention through the stent mesh was attempted. A 0.035-inch guidewire (Jagwire; Boston Scientific, Marlborough, Massachusetts, USA) was successfully passed through the stent mesh (► Fig. 2); however, an ERCP catheter could not be passed. Subsequently, a diathermic dilator was utilized, but it failed to break the stent mesh. Next, the use of a precut needle-knife (NeedleCut3V; Olympus, Tokyo, Japan) was



► Fig. 1 Stent occlusion after endoscopic ultrasound-guided hepaticogastrostomy. **a** Abdominal computed tomography showed a dilated intrahepatic bile duct. **b** Gastroscopy showed an occluded hepaticogastrostomy stent.



► Fig. 2 A 0.035-inch guidewire (Jagwire; Boston Scientific, Marlborough, Massachusetts, USA) was passed successfully through the mesh of the previously deployed hepaticogastrostomy stent (Niti-S biliary covered stent, 8 × 100 mm; Taewoong Medical, Seoul, South Korea).

considered. Using this knife, the stent mesh was broken easily (► Fig. 3), and a 7-Fr plastic stent (Flexima; Boston Scientific) was successfully deployed via the stent mesh into the left intrahepatic bile duct (► Fig. 4, ► Video 1). Cholangitis resolved in a few days.

The use of a precut needle-knife is simple and may be considered as a useful treatment option for reintervention after EUS-HGS.

Endoscopy\_UCTN\_Code\_CPL\_1AL\_2AD


### Competing interests

None


### The authors

Kosuke Minaga, Mamoru Takenaka, Ayana Okamoto, Shunsuke Omoto, Takeshi Miyata, Hajime Imai, Masatoshi Kudo


Department of Gastroenterology and Hepatology, Kindai University Faculty of Medicine, Osaka-Sayama, Japan



► **Fig. 3** A precut needle-knife (NeedleCut3V; Olympus, Tokyo, Japan) was inserted over the guidewire and could break the stent mesh easily.



► **Fig. 4** A 7-Fr biliary plastic stent (70 mm long, Flexima; Boston Scientific, Marlborough, Massachusetts, USA) was deployed successfully via the stent mesh into the left intrahepatic bile duct.



► **Video 1** Using a precut needle-knife, the mesh of the previously deployed hepaticogastrostomy stent was broken easily. Thereafter, a 7-Fr biliary plastic stent was deployed successfully via the stent mesh into the left intrahepatic bile duct.



## Corresponding author

### Mamoru Takenaka, MD

Department of Gastroenterology and Hepatology, Kindai University Faculty of Medicine, 377-2 Ohno-Higashi, Osaka-Sayama 589-8511, Japan  
Fax: +81-72-3672880  
mamoxyo45@gmail.com

## References

- [1] Wang K, Zhu J, Xing L et al. Assessment of efficacy and safety of EUS-guided biliary drainage: a systematic review. *Gastrointest Endosc* 2016; 83: 1218–1227
- [2] Nakai Y, Isayama H, Yamamoto N et al. Safety and effectiveness of a long, partially covered metal stent for endoscopic ultrasound-guided hepaticogastrostomy in patients with malignant biliary obstruction. *Endoscopy* 2016; 48: 1125–1128
- [3] Okuno N, Hara K, Mizuno N et al. Stent migration into the peritoneal cavity following endoscopic ultrasound-guided hepaticogastrostomy. *Endoscopy* 2015; 47 (Suppl. 01): E311
- [4] Ogura T, Masuda D, Takeuchi T et al. Simplified reintervention method of EUS-guided hepaticogastrostomy stent obstruction. *Gastrointest Endosc* 2016; 83: 831
- [5] Minaga K, Takenaka M, Miyata T et al. Through-the-mesh technique after endoscopic ultrasonography-guided hepaticogastrostomy: a novel re-intervention method. *Endoscopy* 2016; 48: E369–E370

## Bibliography

DOI <https://doi.org/10.1055/a-0596-7171>  
Published online: 13.4.2018  
*Endoscopy* 2018; 50: E153–E154  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

## ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>