



Reverse rendezvous with endoscopic retrograde cholangiography and percutaneous transhepatic cholangio drainage: who meets whom?



► **Fig. 1** Cholangiographic images. **a** High grade common bile duct (CBD) stenoses, common hepatic duct (CHD) dilation to 15 mm, and left hilar obstruction not amenable to endoscopic retrograde cholangiography-guided therapy. Right-sided loss of intrahepatic ducts was suspected previously. **b** Rendezvous of wires in the dilated CHD, but access across the significant stenoses was not possible. **c, d** The transpapillary wire was grabbed with a forceps over an 8-Fr bougie and exteriorized percutaneously. **e** Resolution of stenoses in the CBD and CHD 14 months after initial rendezvous.

A 47-year-old severely ill Caucasian man presented with cholestasis (bilirubin 17.8 mg/dL) due to primary sclerosing cholangitis. Endoscopic retrograde cholangiography (ERC) showed high grade strictures of the common bile duct (CBD), dilation of the common hepatic duct (CHD), and left hilar obstruction (► **Fig. 1 a**). Attempts to maneuver 5–7-Fr bougies across the distal CBD stenosis were not successful.

Via left-sided percutaneous transhepatic cholangio drainage (PTCD), retrograde access to the CBD was not possible even after simultaneous transpapillary wire guidance (► **Fig. 1 b**). Therefore, a 1.2-mm biopsy forceps (SpyBite; Boston Scientific, Ratingen, Germany) was introduced percutaneously through an 8-Fr bougie into the dilated CHD to grab the transpapillary 0.025-inch wire. The wire was carefully exteriorized in a reverse rendezvous maneuver (► **Fig. 1 c, d**, ► **Video 1**). Given the lack of bougienage options, a 5.2-Fr angiography catheter (Super Torque Plus; Cordis,



► **Video 1** Reverse rendezvous with endoscopic retrograde cholangiography and percutaneous transhepatic biliary drainage.

Baar, Switzerland) was inserted as a temporary spacer across the papilla under duodenoscopic view. Upon PTCD exchange, spurting bleeding from the access site was stopped by upgrade to

an 8.5-Fr Yamakawa drain (Peter Pflugbeil GmbH, Zorneding, Germany). Parenchymal damage from initial wire manipulation was suspected, so the percutaneous tract was subsequently

occluded with hemostyptic gelatine (Gelita; B. Braun, Melsungen, Germany), and a transpapillary 8.5-Fr pigtail stent was inserted.

The patient gained 10 kg in weight and the bilirubin level persistently dropped to 0.8 mg/dL. After repeated stent upgrades and dilations (► **Fig. 1 e**), dysplasia was ruled out by cholangioscopic biopsies. After 20 months, the patient was well and continued to have regular follow-up with no evidence of recurrence of cholestasis.

To our knowledge, reverse rendezvous, with percutaneous uptake of a transpapillary wire, has not been reported previously. The “lucky punch” of being able to grab the transpapillary wire with a port-guided forceps can be facilitated by C-arm rotation. Unsheathed transparenchymal wire extraction is not recommended as the wire may cut the liver parenchyma, necessitating hemostyptic occlusion of the percutaneous tract, as in our patient. Reverse ERC-PTCD rendezvous is a nonstandard rescue maneuver that can offer significant benefit in technically demanding situations.

Endoscopy_UCTN_Code_CPL_1AK_2AH

Competing interests

Martin Goetz is in the advisory board for Boston Scientific.

The authors

Martin Goetz¹, Jakob Fisch¹, Jürgen Hetzel², Gerd Grözinger³

- 1 Innere Medizin I, Universitätsklinikum Tübingen, Tübingen, Germany
- 2 Innere Medizin II, Universitätsklinikum Tübingen, Tübingen, Germany
- 3 Radiologische Universitätsklinik, Universitätsklinikum Tübingen, Tübingen, Germany

Corresponding author

Martin Goetz, MD

Innere Medizin I, Universitätsklinikum Tübingen, 72076 Tübingen, Germany
 Fax: +49-7071-2925034
 m_goetz@web.de

Bibliography

DOI <https://doi.org/10.1055/a-0800-8342>

Published online: 14.12.2018

Endoscopy 2019; 51: E47–E48

© Georg Thieme Verlag KG

Stuttgart · New York

ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>