

The Terroir of *Cannabis*: Terpene Metabolomics as a Tool to Understand *Cannabis sativa* Selections

Authors

Elizabeth M. Mudge^{1,2}, Paula N. Brown^{2,3}, Susan J. Murch¹

Affiliations

- 1 Chemistry, University of British Columbia, Kelowna, British Columbia, Canada
- 2 Natural Health & Food Products Research, British Columbia Institute of Technology, Burnaby, British Columbia, Canada
- 3 Biology, University of British Columbia, Kelowna, British Columbia, Canada

Key words

Cannabaceae, *Cannabis sativa*, metabolomics, terpenes, headspace GC-MS, entourage

received February 4, 2019

revised April 17, 2019

accepted April 30, 2019


Bibliography

DOI <https://doi.org/10.1055/a-0915-2550>

Published online May 16, 2019 | *Planta Med* 2019; 85: 781–796 © Georg Thieme Verlag KG Stuttgart · New York | ISSN 0032-0943

Correspondence

Susan J. Murch, PhD
University of British Columbia
3247 University Way, Kelowna, BC, Canada
Phone: +1 250 807 9566
susan.murch@ubc.ca

 Supporting information available online at <http://www.thieme-connect.de/products>

ABSTRACT

The phytochemical diversity of *Cannabis* chemovars is not well understood, and many chemovars were created in informal breeding programs without records of parentage or the criteria for selection. Key criteria for selection sometimes included aroma notes and visual cues, which some breeders associated with pharmacological activity. We hypothesized that the process of selection for scents believed to be related to specific tetrahydrocannabinol levels has resulted in modified terpene biosynthesis in these chemovars. Thirty-two cannabinoids, 29 monoterpenes and 38 sesquiterpenes were measured in 33 chemovars from 5 licensed producers. A classification system based on cannabinoid content was used with targeted metabolomic tools to determine relationships in the phytochemistry. Three monoterpenes, limonene, β -myrcene, and α -pinene, and two sesquiterpenes, caryophyllene and humulene, were abundant in the majority of chemovars. Nine terpenes were present in tetrahydrocannabinol-dominant chemovars. Three monoterpenes and four sesquiterpenes were predominantly found in cannabidiol-containing chemovars. Low abundance terpenes may have been the aromatic cues identified by breeders. The medicinal activity of some of the terpenes is likely to contribute to the pharmacological effect of specific chemovars. Together, these data demonstrate the synergy of compounds in *Cannabis* chemovars and point to the need for additional research to understand the phytochemical complexity.

Introduction

Until recently, informal breeding has been the only source of *Cannabis* varieties (chemovars), where normal crop breeding protocols have not been strictly followed [1–3]. Key selection criteria used to select *Cannabis* chemovars included aroma, morphology, and other visual cues that some growers associated with tetrahydrocannabinol (THC) potency [1, 2]. The aroma of chemovars has been shown to play a significant role in the selection, preference, and quality indication of chemovars [4, 5]. The resultant marijuana chemovars may not be genetically distinct from one another but can have different chemistry, and some chemovars with the same name are not genetically similar [6, 7]. Genetic variation

within chemovars is highlighted by the expression of phytochemicals present, and terpenes are one of the major classes of compounds responsible for aroma, and, therefore, are impacted by these breeding practices.

As with many high value products such as wine and hops, the variation in aroma notes is the result of variation in the volatile constituents, including monoterpenes and sesquiterpenes [5, 8, 9]. Terpenes are particularly interesting in *Cannabis* because they are sequestered in glandular trichomes and co-accumulate with the cannabinoids. Both terpenes and cannabinoids are derived from the same precursor molecule, geranyl pyrophosphate, and more than 240 different cannabinoids and terpenes have been described in *Cannabis* [10–13]. Recent data has signified the pres-

► **Table 1** Chemovars of *Cannabis* were clustered into five distinct groups that could be separated by their CBD and THC contents.

Group	Color code	CBD range (% w/w)	THC range (% w/w)	# Chemovars
A	Blue	< MDL – 0.08	11.3–19.1	20
B	Purple	< MDL – 0.02	8.0–9.9	3
C	Orange	7.1–9.7	5.0–6.7	6
D	Green	5.3–8.8	1.7–3.1	3
E	Red	16.1	0.7	1

ence of several terpene synthases in *Cannabis*, mainly producing the major monoterpenes and sesquiterpenes identified in *Cannabis* [14].

Plant metabolomics provides the ability to study small molecules within samples to understand the underlying impacts of genetics, environment, or stressors [15, 16]. Approaches can be used to combine information from targeted and untargeted metabolites to discover relationships, clusters, families, biochemical pathways, genetic expression, and post-translational modifications that would be missed when performing univariate analysis of single metabolites [16–19]. Several data reduction strategies and unsupervised classification techniques have been developed that reduce complex phytochemical diversity issues like those found in *Cannabis* chemovars and can be used to identify relationships between metabolites [17, 20, 21]. Multivariate statistics provide avenues to explore these relationships, which have recently been used to describe the impacts of domestication and breeding on cannabinoid biosynthesis but has not been used to evaluate terpene biosynthesis [22].

It has been suggested that *Cannabis* breeders selected for scent notes that they believe are indicative of high potency chemovars. We hypothesize that this process of selection for scents believed to be related to specific THC levels has resulted in modified terpene biosynthesis in these chemovars. To investigate this hypothesis, we assembled a collection of 33 *Cannabis* chemovars from 5 different producers and profiled the terpenes. Previous analysis had classified the chemovars as THC-dominant or cannabidiol (CBD)-THC hybrid chemovars [22]. Our data indicate that there are groups of terpenes with characteristic aromas that are associated with major cannabinoid content, which was the major focus of many clandestine breeding programs.

Results

A total of 67 terpenes were detected and comprised 29 monoterpenes and 38 sesquiterpenes. Monoterpenes accounted for 87.1 to 99.5% of the terpene profiles, while sesquiterpenes accounted for the remaining 0.5 to 12.9%. Four chemovars had less than 1% sesquiterpenes, while the average content was 5.4%.


The classification system based on cannabinoid content, which has been used to highlight breeding based on cannabinoid potency described by Mudge et al. was used to identify relationships of different terpenes across the classes [22]. These classes are summarized in ► **Table 1**. To assess the relationships between THC, CBD, and the terpenes, each terpene was graphed according

to THC content from lowest to highest and color coded to chemovar class [22]. Five monoterpenes and seven sesquiterpenes were ubiquitous across all chemovars (► **Figs. 1** and **2**). Three monoterpenes, limonene, β -myrcene, and α -pinene, were abundant in the majority of chemovars, while the two most abundant sesquiterpenes, caryophyllene and humulene, ranged from 0.2 to 5.5% and 0.3 to 1.5% respectively.

Seventeen terpenes were found in chemovars from a range of cannabinoid groupings, but not in all chemovars (**Fig. 1S**, Supporting Information). β -Cubebene was found in all chemovars except the very low THC, high CBD chemovar (**Fig. 1Sd**, Supporting Information). There were considerable correlations among the lower abundance sesquiterpenes with correlation coefficients above 0.8, as visually represented in ► **Fig. 3**. Correlations were observed between γ -muurolene, copaene, β -cubebene, elemol, germacrene A, guaia-3,9-diene, β -maaliene, γ -maaliene, selina-3,7(11)-diene, α -selinene, and δ -selinene (► **Fig. 3**), for which many of these metabolites were observed in either all or almost all cannabinoid chemovars and clusters (**Fig. 1S**, Supporting Information).

Eight sesquiterpenes and one monoterpene were present in THC-dominant chemovars (► **Fig. 4** and **Fig. 2S**, Supporting Information) and four that were found to be in chemovars identified as mid-range THC (**Fig. 3S**, Supporting Information). (*Z,Z*)- α -Farnesene was found only in the chemovar CAN36, and β -sesquiphellandrene was found only in the chemovar CAN27 (► **Fig. 4F, H**). δ -Cadiene and an unidentified sesquiterpene were found only in one chemovar, CAN23 (**Fig. 3S**, Supporting Information). Santolina triene (tentative identification) was one of two monoterpenes observed to have correlations with sesquiterpenes sesquiterp-1 (unidentified) and δ -cadinene, all present in this grouping (► **Fig. 3** and **Fig. 3S**, Supporting Information).

There were 18 terpenes present in high abundance in chemovars identified as high THC and mid-level THC/CBD. Ten monoterpenes were identified as strongly correlated to one another, and are shown in ► **Fig. 5**. The remaining eight monoterpene and sesquiterpene profiles observed in this group are summarized in **Fig. 4S**, Supporting Information. Terpinolene was the most dominant monoterpene, which was less than 0.3% in 27 of the 33 chemovars, but ranged from 13.4 to 41.2% in the six chemovars that have this distinctive monoterpene profile: CAN16, CAN17, CAN19, CAN21, CAN32, and CAN33. Terpinolene was correlated with other monoterpenes, such as α -thujene, α -phellandrene, 3-carene, α -terpinene, *p*-cymene, β -phellandrene, α -terpinene, and terpinen-4-ol, with correlation coefficients ranging from 0.95 to




0.99 (► **Fig. 3**). Two sesquiterpene alcohols were also classed in this group and were highly correlated to one another (**Fig. 4S**, Supporting Information).


The final three monoterpenes and four sesquiterpenes were predominantly found in CBD-containing chemovars (**Fig. 5S**, Sup-

porting Information). Two sesquiterpene alcohols, guaiol and 10-epi- γ -eudesmol, were highly correlated to one another (► **Fig. 3**).

The aromas that describe each of the terpenes detected and identified in this collection were compiled from published sources [23] and are grouped according to their presence within the ter-



► **Fig. 2** Sesquiterpene profiles identified as those present across the across the entire dataset. **a** β -Caryophyllene, **b** guaia-3,9-diene, **c** α -guaiene, **d** humulene, **e** β -maaliene, **f** selina-3,7(11)-diene, and **g** valencene.




► Fig. 3 Pearson correlations between monoterpenes and sesquiterpenes within the Cannabis dataset.

pene groupings (► Table 2). The aromas range from pine and woody to spicy, floral, and citrus. While Group 1 terpenes are present in all chemovars and invoke most of the major aromas, Groups 2 to 5 are considered undertones, contributing to unique aromas within each terpene cluster. Group 2 is a combination of woody, floral, and herbal undertones. Group 3, which is found only in THC-dominant chemovars, contained herbal, floral, woody, sweet, and spicy undertones. Group 4 appears to have a considerable amount of citrus, woody, musty, floral, and sweet undertones and Group 5, which has the CBD-dominant chemovars, has primarily citrus, tropical, and sweet undertones.


A principal component analysis (PCA) was performed in the autoscaled terpene profiles to evaluate the clustering and multivariate correlations between the metabolites. The PCA is shown in ► Fig. 6A. The first two principal components (PCs) describe 47.53% of the variance within the data. There is no clear clustering of the chemovars according to their THC/CBD classifications as all five cluster groups overlap significantly. Based on the loading plots of the first two PCs (► Fig. 6B), the majority of the sesquiterpenes cluster together in the top right quadrant of the plot, while the terpinolene-correlated monoterpenes appear to cluster sepa-

rately from the Cannabis groups in the top left quadrant. PC2 appears to have some influence by different monoterpenes; α -pinene and β -myrcene are negatively correlated from the terpinolene-correlated terpenes on this PC.

It was previously noted that many of the monoterpenes and sesquiterpenes were identified across every cannabinoid class. Therefore, a data reduction strategy was undertaken to remove these metabolites and identify any unique clustering of the chemovars when removing these terpenes. In this case, the number of metabolites was reduced from 67 to 38 and then subjected to PCA (Fig. 6S, Supporting Information). The first two PCs of this reduced dataset describe 40.02% of the data. The loading plots indicate that the first PC is clearly influenced by the terpinolene-correlated monoterpenes, for which the chemovars all cluster together on the right side of the scores plot. PC2 appears to cluster a few chemovars on the top left and bottom left quadrant from the majority of the remaining chemovars. These are influenced by the contents of several sesquiterpenes. The chemovars in the top left quadrant are impacted by δ -selinene, germacrene B, α -cubebene, and γ -elementene, all metabolites identified to be present only in THC-dominant chemovars.



► **Fig. 4** Sesquiterpene profiles present primarily in THC-dominant chemovars. **a** α -Amorphene, **b** caryophyllene oxide, **c** α -cubenene, **d** β -elemene, **e** γ -elemene, **f** (Z,Z)- α -farnesene, **g** germacrene B, and **h** β -sesquiphellandrene.




► **Fig. 5** Monoterpene profiles representing a unique group of terpenes that dominate both THC-dominant and CBD-THC hybrid chemovars found to be strongly correlated with terpinolene. a 3-Carene, b *p*-cymene, c *p*-cymenene, d α -phellandrene, e β -phellandrene, f α -terpinene, g γ -terpinene, h terpinen-4-ol, i terpinolene, and j α -thujene. *continued next page*

Discussion

We hypothesized that the practice of selecting *Cannabis* chemovars by aromas thought to be indicative of THC content would result in a set of common scent tones characteristic of high-THC

chemovars, and that the comprehensive and sensitive analysis of terpene profiles in *Cannabis* chemovars could then provide new insights into the impact of the domestication on *Cannabis*. Anecdotal evidence suggests that the informal breeding history of the crop predicted the potency of THC chemovars based on slight



► Fig. 5 continued

aromatic undertones and breeders selected for CBD-containing chemovars by choosing to clone individuals with specific aromas believed to predict these metabolites [1, 3]. *Cannabis* aromas play many roles in chemovar selection, euphoria, and product quality, and are strongly associated with clandestine breeding [1, 4]. Many of the terpenes have similar characteristic aromas, which can be impacted by concentration, synergy with other aromatic compounds, and subjective interpretation of the aroma [24]. Subjective interpretation of “desirable” *Cannabis* aromas during breeding could impact terpene profiles, and many chemovar names are indicative of their aroma. Several dominant aromas described in *Cannabis* names include lemon, sour, skunk, berry/fruit, diesel, or cheese. There has been considerable variation observed between the chemovar name and composition, suggesting that some chemovar names may not accurately describe aroma due to phytochemical variance [25].

Headspace GC-MS analysis was employed for the profiling of monoterpenes and sesquiterpenes in *Cannabis* because of its sensitivity in comparison to solvent extraction methods and the ability to highlight the aromatic expression (headspace) of the che-

movars. This method detected 67 metabolites identified with reference standards and the NIST spectral database with considerable matching capabilities. In many previous characterizations of *Cannabis*, the number of terpenes ranged from 14 to 37, focusing only on high abundance terpenes and leaving a considerable number not evaluated [25–29]. Over 120 different terpenes have been previously detected in *Cannabis*, but many of those not detected in this study are typically present in trace levels [29]. The implementation of this more sensitive technique provides a deeper insight into the phytochemical variation within chemovars and the underlying variances that would otherwise be overlooked in traditional solvent extraction-based methods. Evaluating terpene profiles and potential aromatic characteristics provides a deeper insight into aroma selection by breeders and patients [4].

Positive correlations were observed of many low abundance terpenes with the cannabinoid classes. High-THC chemovars had a higher prevalence of herbal and floral undertones and a higher prevalence of several sesquiterpenes. Interestingly, caryophyllene oxide was correlated with high-THC chemovars and is a sesquiterpene identified by canine enforcement officers to detect drugs

► **Table 2** Aroma descriptors for each of the terpenes identified within the *Cannabis* chemovars grouped based on their presence within the cannabinoid classes.

Group	Terpene	Scent descriptors	Aroma
Group 1	α -pinene	woody/pine	woody, pine, citrus, spicy, floral
	β -pinene	woody/pine	
	<i>trans</i> -2-pinanol	pine	
	camphene	woody/camphor	
	α -gurjunene derivative	woody/balsamic	
	β -maaliene	woody	
	selina-3,7(11)-diene	possible woody	
	camphene hydrate	woody/camphor	
	α -bergamotene	woody	
	4,11-selinadiene	woody	
	endo-borneol	camphor	
	fenchone	camphor	
	Z-sabinene hydrate	balsam	
	γ -gurjunene	musty	
	β -myrcene	spicy/balsamic/peppery	
	caryophyllene	spicy/cloves/roses	
	copaene	spicy/honey	
	γ -muurolene	spicy	
	D-limonene	citrus	
	α -terpineol	citrus	
	β -cubebene	citrus	
	valencene	citrus	
	guaia-3,9-diene	floral, rose, geranium	
	germacrene A	floral	
	ylangene	ylang ylang	
humulene	hoppy		
α -selinene	celery		
exo-fenchol	basil		
β -selinene	celery		
Group 2	α -gurjunene	woody/balsamic	floral, woody, herbal
	santolina triene	floral	
	sesquiterp-1	n/a	
	δ -cadinene	herbal/thyme	
Group 3	α -amorphene	woody	herbal, woody, floral, citrus
	caryophyllene oxide	spicy, woody/carrot	
	germacrene B	floral/roses	
	γ -elemene	floral	
	2-carene	sweet	
	(Z,Z)- α -farnesene	citrus	
	α -cubenene	herbal	
	β -elemene	herbal	
	β -sesquiphellandrene	herbal/oregano	

continued next page

► Table 2 Continued

Group	Terpene	Scent descriptors	Aroma
Group 4	α -thujene	woody/frankincense	citrus, woody, sweet, spicy
	α -terpinene	woody	
	<i>cis</i> - β -terpineol	woody	
	α -santalene	woody	
	α -bulnesene	Patchouli	
	α -fenchene	camphor	
	<i>cis</i> - β -farnesene	citrus/sweet	
	<i>p</i> -cymene	citrus/sweet	
	α -phellandrene	citrus/pepper	
	γ -terpinene	citrus	
	terpinolene	sweet/pine/citrus	
	linalool	citrus/floral/sweet	
	δ -selinene	floral	
	3-carene	sweet	
	α -eudesmol	sweet	
	terpinen-4-ol	peppery/musty/sweet	
	<i>p</i> -cymenene	spicy/cloves	
β -phellandrene	minty		
bulnesol	spicy		
Group 5	alloaromadendrene	woody	citrus, woody, sweet, tropical
	guaiol	rose wood	
	10- <i>epi</i> - γ -eudesmol	sweet	
	<i>cis</i> - α -bisabolene	citrus/myrrh/balsamic	
	<i>cis</i> - β -ocimene	citrus/tropical	
	<i>trans</i> - β -ocimene	citrus/tropical	
	sabinene	citrus/pine/spicy	

[30]. The CBD-containing chemovars were higher in citrus and tropical undertones, which were attributed to several monoterpenes and sesquiterpene alcohols. Aromas are determined based on volatility, threshold, concentration, and interactions with other aromatic compounds, therefore, the data described are only a preliminary estimation of aromatic characteristics from each compound [24].

Cannabis chemovars with similar THC/CBD contents exhibit varying pharmacological effects [3,22,29,31], and previous authors have proposed an “entourage effect” theory, suggesting that cannabinoids and terpenes act synergistically to invoke varying pharmacological effects [5,31,32]. There are over 30 000 known terpenes in plants [23]. A summary of the pharmacological activities of terpenes identified in this collection that have been described in the literature through *in vitro*, *in vivo*, and clinical studies are presented in ► Table 3. Major monoterpenes such as α -pinene, β -myrcene, and limonene have been shown to have anti-inflammatory, analgesic, and sedative properties evaluated in animal models, respectively [33–36] (► Table 3). Terpinolene, present in high abundance in only a select few chemovars, also showed anti-inflammatory and sedative properties in animal models [37,38] (► Table 3). Minor terpenes may also play a signif-

icant role. Linalool has been shown to have anti-inflammatory, sedative, anxiolytic, anticonvulsant, and antidepressant activities [31,39]. Cymene has antinociceptive activity [40]. Terpinen-4-ol has been studied extensively for its anticonvulsant and anticancer activities [41,42].

Data from other medicinal plants can aid in understanding the pharmacological effects of many of the terpenes in *Cannabis*. For example, *Salvia* sp. and *Ocimum sanctum* (holy basil) are used for their analgesic, antidepressant, anxiolytic, and anti-inflammatory activities [43,44]. These plants have many similar terpenes including borneol, β -pinene, α -pinene, camphene α -thujene, β -caryophyllene, sabinene, limonene, *p*-cymene, terpinolene, ocimene, α -cubebene, linalool, β -elemene, β -caryophyllene, α -guaiene, α -amorphene, α -humulene, isoborneol, borneol, α -selinene, β -selinene, and α -muurolene. *Myrcia* spp. have many similar terpenes and exhibit anti-inflammatory, antiproliferative, and antinociceptive activities [45]. Similarly, *Ocimum basicicum* has reported antidepressant and anticonvulsant activities and similar terpene chemistry [46]. Further research is needed to understand the synergy of these bioactive compounds and the pharmacological significance for humans.

► **Table 3** Reported activities of monoterpenes and sesquiterpenes identified in *Cannabis* through *in vitro* cell-based models, *in vivo* animal models, and clinical data.

Group	Terpene	Pharmacological activity	References
Group 1	camphene	expectorant	[64]
	caryophyllene	anti-inflammatory, antinociceptive, anxiolytic, antispasmodic, antidepressant, gastroprotective	[47, 65–69]
	fenchone	antinociceptive activity	[70]
	humulene	anti-inflammatory, antitumor	[67, 71]
	limonene	antioxidant, tumor reduction, sedative	[33, 34, 72, 73]
	β -maaliene	sedative	[49]
	β -myrcene	sedative, analgesic, antioxidant,	[36, 74, 75]
	α -pinene	antinociceptive activity, anti-inflammatory, anxiolytic	[35, 70, 76]
	β -pinene	antidepressant	[77]
	α -terpineol	anti-inflammatory, antinociceptive, gastroprotective	[78, 79]
	valencene	anti-melanogenesis activity, UV protectant, anti-inflammatory	[80, 81]
Group 3	caryophyllene oxide	analgesic, anti-inflammatory, tumor inhibition	[82, 83]
	β -elemene	antitumor	[55]
	β -sesquiphellandrene	tumor inhibition	[84]
Group 4	α -bulnesene	antiplatelet	[85]
	<i>p</i> -cymene	antinociceptive, anti-inflammatory, antioxidant	[86, 87]
	β -eudesmol	anti-inflammatory, muscle relaxant, anti-cholangiocarcinoma activity, appetite stimulation, antiangiogenic, gastroprotective, anticonvulsant	[88–92]
	linalool	antidepressant, antinociceptive, sedative	[77, 93, 94]
	α -phellandrene	antinociceptive	[65]
	γ -terpinene	antioxidant, antinociceptive	[95, 96]
	terpinen-4-ol	antimicrobial, antihypertensive, anticonvulsant, tumor inhibition	[41, 42, 97, 98]
Group 5	terpinolene	antinociceptive, anti-inflammatory, sedative	[37, 38]
	alloaromadendrene	antioxidant	[99]
	<i>cis</i> - α -bisabolene	anticonvulsant	[57]
	guaiol	anti-inflammatory	[100]
	sabinene	antimicrobial	[97]

The sensitive analytical method employed in this work allowed a significantly higher number of terpenes to be detected and identified in the chemovars. This expansion of chemical composition allowed for increased chemical characterization and identified several low abundance terpenes associated with cannabinoid potency. The data suggest that domestication syndrome, resulting from informal breeding and selection, has impacted phytochemical diversity, which may be associated with the pharmacological variance observed across chemovars. Future research is needed to understand the activities of low abundance terpenes and synergistic effects in *Cannabis* chemovars and to determine the importance for medical efficacy and their roles in plant biosynthesis.

Materials and Methods

Reagents

HPLC grade methanol and acetonitrile were purchased from VWR International. Water was deionized and purified to 18.2 M Ω using a Barnstead Smart2Pure nanopure system (Thermo Scientific). Ammonium formate, HPLC grade (>99.0%), was purchased from Sigma-Aldrich and formic acid (98%) was HPLC grade and purchased from Fisher Scientific. Cannabinoid certified reference material standards purchased as 1 mL solutions in ampules were purchased from Cerilliant Corp. They included tetrahydrocannabinolic acid (THCA, 1.000 mg/mL), Δ 9-tetrahydrocannabinol (THC, 1.001 mg/mL), cannabidiolic acid (CBDA, 1.000 mg/mL), cannabidiol (CBD, 1.000 mg/mL), cannabigerol (CBG, 1.000 mg/mL), cannabichromene (CBC, 1.000 mg/mL), tetrahydrocannabivarin (THCV, 1.00 mg/mL), and cannabinol (CBN, 1.000 mg/mL). Additional standards were purchased for peak identification from Cerilliant Corp., which included Δ 8-THC (1.000 mg/mL), cannabi-

divarinic acid (CBDVA, 1.000 mg/mL), cannabidivarin (CBDV, 1.000 mg/mL), cannabigerolic acid (CBGA, 1.000 mg/mL), and cannabicyclol (CBL, 1.000 mg/mL). All cannabinoid standards were provided in either methanol or acetonitrile. Cannabis Terpene Mix A and Mix B containing 20 and 15 terpenes, respectively, at 2000 µg/mL in methanol were purchased from Sigma-Aldrich. Cannabis Terpene Mix A contained α -pinene, camphene, β -pinene, 3-carene, α -terpinene, limonene, γ -terpinene, fenchone, fenchol, camphor, isoborneol, menthol, citronellol, pulegone, geranyl acetate, α -cedrene, α -humulene, nerolidol, cedrol, and α -bisabolol. Cannabis Terpene Mix B contained β -pinene, 3-carene, *p*-cymene, limonene, terpinolene, linalool, camphor, borneol, α -terpineol, geraniol, β -caryophyllene, *cis*-nerolidol, β -eudesmol, and phytol.

Plant materials

Thirty-three chemovars of *Cannabis sativa* L. were purchased from five licensed producers in Canada under the Access to *Cannabis* for Medical Purposes Regulation (ACMPR), and laboratory analysis was performed under a Health Canada Research License. The test samples were provided as whole or milled flowers in 5-, 10-, and 15-gram packages and stored at room temperature until use. Due to the legal restrictions pertaining to the storage of *Cannabis* chemovars, submission of voucher specimens to an herbarium was not possible, but given the regulatory framework associated with these plants, their identify has been confirmed as *C. sativa* L.

Cannabinoid analysis

The content of 32 cannabinoids was determined previously [22, 61]. In brief, ground *Cannabis* flowers (0.200 g) were extracted with 25 mL of 80% methanol for 15 min, followed by centrifugation at 4500 *g* for 5 min and filtration with a 0.22-µm PTFE filter. Extracts were diluted to within the calibration range using the extraction solvent and placed in the 4 °C sample holder for same-day analysis. Chromatographic separation was performed on an Agilent 1200 UHPLC with a Kinetex C18 100 mm × 3.0 mm, 1.8 µm column (Phenomenex) using a gradient elution with 10 mM ammonium formate (pH 3.6) and acetonitrile with detection at 220 nm. Chemovars were classified into five clusters based on the range of CBD/THC values determined [22].

Evaluation of volatile constituents

Terpene profiles were determined using an in-house developed method, adapted from a previous terpene method [13]. Immediately after grinding, *Cannabis* flowers (0.100 g) were added to a 20-mL gas tight headspace vial. Samples were prepared in triplicate. Using a CTC Analytics Combi-PAL headspace autosampler, the vials were transferred to a heated incubator at 80 °C for 15 min and agitated at 500 rpm. Next, 1000 µL of the vial headspace was injected using a syringe at 120 °C. The injector temperature was 230 °C with a split ratio of 10:1. GC analysis was undertaken on an Agilent 7890A GC coupled to a 5975B mass spectrometer (MS). Separation was achieved on a 20 m × 180 µm ID, 0.18 µm film thickness J&W DB-5MS column. Helium was used as the carrier gas at a flow rate of 1.3 mL/min. The column was held at 50 °C for 3 min followed by a ramp to 170 °C at 5 °C/min for a total run time of 27 min. MS detection with electron impact

ionization at 70 eV was used to collect mass spectra from *m/z* 50 to 500. The MS quad and source temperatures were 230 and 150 °C, respectively.

Chemometrics

Metabolite profiling

Terpenes were identified based on comparison of mass spectra with the NIST spectral database (NIST 11). Additionally, retention indices were compared to published literature to confirm elution order and identity [62]. Several monoterpene standards were also analyzed to confirm identity. Multivariate curve resolution using SOLO+MIA software (version 8.5; Eigenvector Research) was employed to separate coeluting terpenes, and peak areas were determined using the software program R, version 3.5.2 [63]. Peaks were manually aligned based on compound identity and retention time using Excel. Missing values (zeros) were replaced with half of the lowest value in the dataset.

Identification of metabolite relationships

Individual terpenes were plotted according to their cannabinoid profiles, previously described by Mudge et al. to identify trends within the datasets and classify them into unique groups [22]. Trends evaluated included those present across all chemovars, those found primarily in THC-dominant chemovars, those present primarily in CBD-THC hybrid chemovars, and other terpene correlations independent of cannabinoid content. Correlations between terpenes and cannabinoids were confirmed by evaluating Pearson correlation coefficients using the R program *cor*.

Multivariate classification

The data were autoscaled by mean centering and scaling to unit variance in order to give each metabolite equal weight prior to multivariate analyses. PCA was subsequently performed using Solo+MIA.

Supporting information

Terpene profiles identified across different cannabinoid classes, but not present in all chemovars (endo-borneol, camphene hydrate, copaene, β -cubebene, exo-fenchol, fenchone, germacrene A, α -gurjunene derivative, γ -gurjunene, γ -muurolene, *trans*-2-pinanol, *z*-sabinine hydrate, 4,11-selinadiene, α -selinene, β -selinene, α -terpineol, ylangene) are described in Fig. 1S. The profile of the monoterpene 2-carene, detected primarily in THC-dominant chemovars is described in Fig. 2S. The terpene profiles for those found primarily in mid-level THC-dominant chemovars (δ -cadiene, α -gurjunene, santolina triene, sesquiterp-1) are summarized in Fig. 3S. Several additional terpenes representing a unique group of terpenes that dominate both THC-dominant and THC-CBD hybrid chemovars (α -bulnesene, bulnesol, α -eudesmol, *cis*- β -farnesene, α -fenchene, linalool, α -santolene, δ -selinene) are summarized in Fig. 4S. The profiles of terpenes found predominantly in higher CBD chemovars (alloaromadendrene, *cis*- α -bisabolene, 10-epi- γ -eudesmol, guaial, *cis*- β -ocimene, *trans*- β -ocimene, sabinene) are summarized in Fig. 5S. A PCA of the monoterpene and sesquiterpene profiles after undertaking a data reduction strategy, and the associated loading plots are summarized in Fig. 6S.

Acknowledgements

This research was undertaken, in part, thanks to funding from the Canada Research Chairs program.

Conflict of Interest

The authors declare no conflicts of interest.

References

- Clarke RC, Merlin MD. Cannabis domestication, breeding history, present-day genetic diversity, and future prospects. *CRC Crit Rev Plant Sci* 2016; 35: 293–327
- Small E. Evolution and classification of *Cannabis sativa* (Marijuana, Hemp) in relation to human utilization. *Bot Rev* 2015; 81: 189–294
- Lewis MA, Russo EB, Smith KM. Pharmacological foundations of *Cannabis* chemovars. *Planta Med* 2018; 84: 225–233
- Gilbert AN, DiVerdi JA. Consumer perceptions of strain differences in *Cannabis* aroma. *PLoS One* 2018; 13: 1–14
- Russo EB. The case for the entourage effect and conventional breeding of clinical cannabis: No “strain,” no gain. *Front Plant Sci* 2019; 9: 1969
- Sawler J, Stout JM, Gardner KM, Hudson D, Vidmar J, Butler L, Page JE, Myles S. The genetic structure of marijuana and hemp. *PLoS One* 2015; 10: 1–9
- Soler S, Gramazio P, Figàs MR, Vilanova S, Rosa E, Lloa ER, Borràs D, Plazas M, Prohens J. Genetic structure of *Cannabis sativa* var. *indica* cultivars based on genomic SSR (gSSR) markers: Implications for breeding and germplasm management. *Ind Crops Prod* 2017; 104: 171–178
- Arrhenius SP, McCloskey LP, Sylvan M. Chemical markers for aroma of *Vitis vinifera* Var. Chardonnay regional wines. *J Agric Food Chem* 1996; 44: 1085–1090
- Inui T, Tsuchiya F, Ishimaru M, Oka K, Komura H. Different beers with different hops. Relevant compounds for their aroma characteristics. *J Agric Food Chem* 2013; 61: 4758–4764
- Flores-Sanchez IJ, Verpoorte R. Secondary metabolism in *Cannabis*. *Phytochem Rev* 2008; 7: 615–639
- ElSohly MA, Gul W. Constituents of *Cannabis sativa*. In: Pertwee R, editor. *Handbook of Cannabis*. Oxford UK: Oxford University Press; 2014
- Turner CE, Elsohly MA, Boeren EG. Constituents of *Cannabis sativa* L. XVII. A review of the natural constituents. *J Nat Prod* 1980; 43: 169–234
- Giese MW, Lewis MA, Giese L, Smith KM. Development and validation of a reliable and robust method for the analysis of cannabinoids and terpenes in *Cannabis*. *J AOAC Int* 2015; 98: 1503–1522
- Booth JK, Page JE, Bohlmann J. Terpene synthases from *Cannabis sativa*. *PLoS One* 2017; 12: 1–21
- Fiehn O, Kopka J, Dörmann P, Altmann T, Trethewey RN, Willmitzer L. Metabolite profiling for plant functional genomics. *Nat Biotechnol* 2000; 18: 1157–1161
- Turi CE, Finley J, Shipley PR, Murch SJ, Brown PN. Metabolomics for phytochemical discovery: development of statistical approaches using a cranberry model system. *J Nat Prod* 2015; 78: 953–966
- Brown PN, Murch SJ, Shipley P. Phytochemical diversity of cranberry (*Vaccinium macrocarpon* Aiton) cultivars by anthocyanin determination and metabolomic profiling with chemometric analysis. *J Agric Food Chem* 2012; 60: 261–271
- Turi CE, Murch SJ. Targeted and untargeted phytochemistry of *Ligusticum canbyi*: indoleamines, phthalides, antioxidant potential, and use of metabolomics as a hypothesis-generating technique for compound discovery. *Planta Med* 2013; 79: 1370–1379
- Turi CE, Axwik KE, Smith A, Jones AMP, Saxena PK, Murch SJ. Galanthamine, an anticholinesterase drug, effects plant growth and development in *Artemisia tridentata* Nutt. via modulation of auxin and neurotransmitter signaling. *Plant Signal Behav* 2014; 9: e28645
- Brown PN, Turi CE, Shipley PR, Murch SJ. Comparisons of large (*Vaccinium macrocarpon* Ait.) and small (*Vaccinium oxycoccos* L., *Vaccinium vitis-idaea* L.) cranberry in British Columbia by phytochemical determination, antioxidant potential, and metabolomic profiling with chemometric analysis. *Planta Med* 2012; 78: 630–640
- Murch SJ, Rupasinghe HPV, Goodenow D, Saxena PK. A metabolomic analysis of medicinal diversity in Huang-qin (*Scutellaria baicalensis* Georgi) genotypes: discovery of novel compounds. *Plant Cell Rep* 2004; 23: 419–425
- Mudge EM, Murch SJ, Brown PN. Chemometric Analysis of Cannabinoids: Chemotaxonomy and Domestication Syndrome. *Sci Rep* 2018; 8: 13090
- Breitmaier E. Terpenes: flavors, fragrances, pharmaca, pheromones. Morlenbach, Germany: Wiley-VCH; 2006
- Wu Y, Duan S, Zhao L, Gao Z, Luo M, Song S, Xu W, Zhang C, Ma C, Wang S. Aroma characterization based on aromatic series analysis in table grapes. *Sci Rep* 2016; 6: 1–16
- Elzinga S, Fishedick JT. Cannabinoids and terpenes as chemotaxonomic markers in *Cannabis*. *Nat Prod Chem Res* 2015; 3: 4
- Hazekamp A, Fishedick J. *Cannabis* – from cultivar to chemovar. *Drug Test Anal* 2012; 4: 660–667
- Fishedick JT, Hazekamp A, Erkelens T, Choi YH, Verpoorte R. Metabolic fingerprinting of *Cannabis sativa* L., cannabinoids and terpenoids for chemotaxonomic and drug standardization purposes. *Phytochemistry* 2010; 71: 2058–2073
- Jin D, Jin S, Yu Y, Lee C, Chen J. Classification of *Cannabis* cultivars marketed in Canada for medical purposes by quantification of cannabinoids and terpenes using HPLC-DAD and GC-MS. *J Anal Bioanal Tech* 2017; 8: 1–9
- Hazekamp A, Tejkalová K, Papadimitriou S. *Cannabis*: from cultivar to chemovar II—a metabolomics approach to *Cannabis* classification. *Cannabis Cannabinoid Res* 2016; 1: 202–215
- Mediavilla V, Steinemann S. Essential oil of *Cannabis sativa* L. strains. *J Intl Hemp Assoc* 1997; 4: 82–84
- Russo EB. Taming THC: potential cannabis synergy and phytocannabinoid-terpenoid entourage effects. *Br J Pharmacol* 2011; 163: 1344–1364
- McPartland JM. *Cannabis* systematics at the levels of family, genus, and species. *Cannabis Cannabinoid Res* 2018; 3: 203–212
- Yun J. Limonene inhibits methamphetamine-induced locomotor activity via regulation of 5-HT neuronal function and dopamine release. *Phyto-medicine* 2014; 21: 883–887
- Do Vale TG, Furtado EC, Santos JG, Viana GSB. Central effects of citral, myrcene and limonene, constituents of essential oil chemotypes from *Lippia alba* (mill.) N.E. Brown. *Phytomedicine* 2002; 9: 709–714
- Kim D, Lee H, Jeon Y, Han Y, Kee J, Kim H, Shin H, Kang J, Lee B, Kim S, Kim S, Park S, Choi B, Park S, Um J, Hong S. Alpha-pinene exhibits anti-inflammatory activity through the suppression of MAPKs and NF- κ B pathway in mouse peritoneal macrophages. *Am J Chin Med* 2015; 43: 731–742
- Freitas J, Presgrave O, Fingola F, Menezes M, Paumgarten F. Effect of beta-myrcene on pentobarbital sleeping time. *Brazilian J Med Biol Res* 1993; 26: 519–523
- Macedo E, Santos W, Sousa BP Neto, Lopes E, Piauilino C, Cunha F, Sousa D, Oliveira F, Almeida F. Association of terpinolene and diclofenac presents antinociceptive and anti-inflammatory synergistic effects in a model of chronic inflammation. *Brazilian J Med Biol Res* 2016; 49: e5103
- Ito K, Ito M. The sedative effect of inhaled terpinolene in mice and its structure-activity relationships. *J Nat Med* 2013; 67: 833–837

- [39] Linck VM, da Silva AL, Figueiró M, Caramão EB, Moreno PRH, Elisabetsky E. Effects of inhaled Linalool in anxiety, social interaction and aggressive behavior in mice. *Phytomedicine* 2010; 17: 679–683
- [40] Bonjardim L, Cunha E, Guimaraes A, Santana M, Oliveira M, Serafini M, Araujo A, Antonioli A, Cavalcanti S, Santos M, Quintans-Junior L. Evaluations of the anti-inflammatory and antinociceptive properties of *p*-cymene in mice. *Zeitschrift für Naturforsch* 2012; 67: 15–21
- [41] de Sousa DP, Nóbrega FFF, de Morais LCSL, de Almeida RN. Evaluation of the anticonvulsant activity of terpinen-4-ol. *Z Naturforsch C* 2009; 64: 1–5
- [42] Calcabrini A, Stringaro A, Toccaceli L, Meschini S, Marra M, Colone M, Salvatore G, Mondello F, Arancia G, Molinari A. Terpinen-4-ol, the main component of *Melaleuca alternifolia* (tea tree) oil inhibits the *in vitro* growth of human melanoma cells. *J Invest Dermatol* 2004; 122: 349–360
- [43] Garg P, Sardana S. Pharmacological and therapeutic effects of *Ocimum sanctum*. *Eur J Pharm Med Res* 2016; 3: 637–640
- [44] Fu Z, Wang H, Hu X, Sun Z, Han C. The pharmacological properties of salvia essential oils. *J Appl Pharm Sci* 2013; 3: 122–127
- [45] Cascaes MM, Guilhon GMSP, de Aguiar Andrade EH, das Graças Bichara Zoghbi M, da Silva Santos L. Constituents and pharmacological activities of Myrcia (Myrtaceae): A review of an aromatic and medicinal group of plants. *Int J Mol Sci* 2015; 16: 23881–23904
- [46] Khair-ul-Bariyah S, Ahmed D, Ikram M. *Ocimum basilicum*: a review on phytochemical and pharmacological studies. *Pakistan J Chem* 2012; 2: 78–85
- [47] Bahi A, Al Mansouri S, Al Memari E, Al Ameri M, Nurulain SM, Ojha S. β -Caryophyllene, a CB2 receptor agonist produces multiple behavioral changes relevant to anxiety and depression in mice. *Physiol Behav* 2014; 135: 119–124
- [48] Rogerio AP, Andrade EL, Leite DFP, Figueiredo CP, Calixto JB. Preventive and therapeutic anti-inflammatory properties of the sesquiterpene α -humulene in experimental airways allergic inflammation. *Br J Pharmacol* 2009; 158: 1074–1087
- [49] Takemoto H, Yagura T, Ito M. Evaluation of volatile components from spikenard: valerenen-4,7(11)-diene is a highly active sedative compound. *J Nat Med* 2009; 63: 380–385
- [50] Peng W, Han T, Xin WB, Zhang XG, Zhang QY, Jia M, Qin LP. Comparative research of chemical constituents and bioactivities between petroleum ether extracts of the aerial part and the rhizome of *Atractylodes macrocephala*. *Med Chem Res* 2011; 20: 146–151
- [51] Pang Y, Wang D, Fan Z, Chen X, Yu F, Hu X, Wang K, Yuan L. *Blumea balsamifera*—a phytochemical and pharmacological review. *Molecules* 2014; 19: 9453–9477
- [52] Al-Snafi PDAE. The pharmacological and therapeutic importance of *Eucalyptus* species grown in Iraq. *IOSR J Pharm* 2017; 7: 72–91
- [53] Hieu LD, Hoi TM, Thang TD, Eresanya OI. Fonenol, the main constituent of the essential oil of the leaf of *Piper longum* L. *Am J Essent Oil Nat Prod* 2018; 6: 16–19
- [54] de Albuquerque IL, Alves LA, Lemos TLG, Dorneles CA, de Morais MO. Constituents of the essential oil of Brazilian green propolis from Brazil. *J Essent Oil Res* 2008; 20: 414–415
- [55] Chen W, Lu Y, Wu J, Gao M, Wang A, Xu B. Beta-elemene inhibits melanoma growth and metastasis via suppressing vascular endothelial growth factor-mediated angiogenesis. *Cancer Chemother Pharmacol* 2011; 66: 799–808
- [56] Tsuneki H, Ma EL, Kobayashi S, Sekizaki N, Maekawa K, Sasaoka T, Wang MW, Kimura I. Antiangiogenic activity of β -eudesmol *in vitro* and *in vivo*. *Eur J Pharmacol* 2005; 512: 105–115
- [57] Orellana-Paucar AM, Serruys ASK, Afrikanova T, Maes J, De Borggraeve W, Alen J, León-Tamariz F, Wilches-Arizábalá IM, Crawford AD, de Witte PAM, Esguerra CV. Anticonvulsant activity of bisabolene sesquiterpenoids of *Curcuma longa* in zebrafish and mouse seizure models. *Epilepsy Behav* 2012; 24: 14–22
- [58] Devinsky O, Cilio MR, Cross H, Fernandez-Ruiz J, French J, Hill C, Katz R, Di Marzo V, Jutras-Aswad D, Notcutt WG, Martinez-Orgado J, Robson PJ, Rohrback BG, Thiele E, Whalley B, Friedman D. Cannabidiol: pharmacology and potential therapeutic role in epilepsy and other neuropsychiatric disorders. *Epilepsia* 2014; 55: 791–802
- [59] Meyer RS, Duval AE, Jensen HR. Patterns and processes in crop domestication: an historical review and quantitative analysis of 203 global food crops. *New Phytol* 2012; 196: 29–48
- [60] Tholl D. Terpene synthases and the regulation, diversity and biological roles of terpene metabolism. *Curr Opin Plant Biol* 2006; 9: 297–304
- [61] Mudge EM, Liu Y, Lund JA, Brown PN. Single-laboratory validation for the determination of flavonoids in Hawthorn leaves and finished products by LC-UV. *Planta Med* 2016; 82: 1487–1492
- [62] Babushok VI, Linstrom PJ, Zenkevich IG. Retention indices for frequently reported compounds of plant essential oils. *J Phys Chem Ref Data* 2011; 40: 043101
- [63] Ruckebusch C, Blanchet L. Multivariate curve resolution: a review of advanced and tailored applications and challenges. *Anal Chim Acta* 2013; 765: 28–36
- [64] Boyd E, Sheppard P. Nutmeg oil and camphene as inhaled expectorants. *Arch Otolaryngol* 1970; 92: 372–378
- [65] Galdino PM, Nascimento MVM, Florentino IF, Lino RC, Fajemiroye JO, Chaibub BA, de Paula JR, de Lima TCM, Costa EA. The anxiolytic-like effect of an essential oil derived from *Spiranthera odoratissima* A. St. Hil. leaves and its major component, β -caryophyllene, in male mice. *Prog Neuropsychopharmacol Biol Psychiatry* 2012; 38: 276–284
- [66] Katsuyama S, Mizoguchi H, Kuwahata H, Komatsu T, Nagaoka K, Nakamura H, Bagetta G, Sakurada T, Sakurada S. Involvement of peripheral cannabinoid and opioid receptors in β -caryophyllene-induced antinociception. *Eur J Pain* 2013; 17: 664–675
- [67] Fernandes ES, Passos GF, Medeiros R, da Cunha FM, Ferreira J, Campos MM, Pianowski LF, Calixto JB. Anti-inflammatory effects of compounds alpha-humulene and (-)-*trans*-caryophyllene isolated from the essential oil of *Cordia verbenacea*. *Eur J Pharmacol* 2007; 569: 228–236
- [68] Leonhardt V, Leal-Cardoso JH, Lahlou S, Albuquerque AAC, Porto RS, Celedônio NR, Oliveira AC, Pereira RF, Silva LP, Garcia-Teófilo TMN, Silva APFS, Magalhães PJC, Duarte GP, Coelho-De-Souza AN. Antispasmodic effects of essential oil of *Pterodon polygalaeiflorus* and its main constituent β -caryophyllene on rat isolated ileum. *Fundam Clin Pharmacol* 2010; 24: 749–758
- [69] Cho JY, Chang HJ, Lee SK, Kim HJ, Hwang JK, Chun HS. Amelioration of dextran sulfate sodium-induced colitis in mice by oral administration of β -caryophyllene, a sesquiterpene. *Life Sci* 2007; 80: 932–939
- [70] Him A, Ozbek H, Turel I, Oner AC. Antinociceptive activity of alpha-pinene and fenchone. *Pharmacol Online* 2008; 369: 363–369
- [71] El Hadri A, del Río MÁ, Sanz J, Coloma AG, Idaomar M, Ozonias BR, González JB, Reus MI. Cytotoxic activity of α -humulene and *trans*-caryophyllene from *Salvia officinalis* in animal and human tumor cells. *Anal Real Acad Nal Farm* 2010; 76: 343–356
- [72] Miller JA, Lang JE, Ley M, Nagle R, Hsu CH, Thompson PA, Cordova C, Waer A, Chow HHS. Human breast tissue disposition and bioactivity of limonene in women with early-stage breast cancer. *Cancer Prev Res* 2013; 6: 577–584
- [73] Murali R, Karthikeyan A, Saravanan R. Protective effects of D-limonene on lipid peroxidation and antioxidant enzymes in streptozotocin-induced diabetic rats. *Basic Clin Pharmacol Toxicol* 2013; 112: 175–181
- [74] Paumgartten FJ, Delgado IF, Alves EN, Nogueira AC, de-Farias RC, Neubert D. Single dose toxicity study of β -myrcene, a natural analgesic substance. *Brazilian J Med Biol Res* 1990; 23: 873–877
- [75] Ciftci O, Ozdemir I, Tanyildizi S, Yildiz S, Oguzturk H. Antioxidative effects of curcumin, β -myrcene and 1,8-cineole against 2,3,7,8-tetra-

- chlorodibenzo-p-dioxin-induced oxidative stress in rats liver. *Toxicol Ind Health* 2011; 27: 447–453
- [76] Satou T, Kasuya H, Maeda K, Koike K. Daily inhalation of α -pinen in mice: effects on behavior and organ accumulation. *Phyther Res* 2014; 28: 1284–1287
- [77] Guzmán-Gutiérrez SL, Bonilla-Jaime H, Gómez-Cansino R, Reyes-Chilpa R. Linalool and β -pinene exert their antidepressant-like activity through the monoaminergic pathway. *Life Sci* 2015; 128: 24–29
- [78] De Oliveira MGB, Marques RB, De Santana MF, Santos ABD, Brito FA, Barreto EO, De Sousa DP, Almeida FRC, Badauê-Passos D, Antonioli ÂR, Quintans-Júnior LJ. α -Terpineol reduces mechanical hypernociception and inflammatory response. *Basic Clin Pharmacol Toxicol* 2012; 111: 120–125
- [79] Souza R, Cardoso M, Menezes C, Silva J, De Sousa D, Batista J. Gastroprotective activity of α -terpineol in two experimental models of gastric ulcer in rats. *Daru* 2011; 19: 277–281
- [80] Nam JH, Nam DY, Lee DU. Valencene from the Rhizomes of *Cyperus rotundus* Inhibits Skin Photoaging-Related Ion Channels and UV-Induced Melanogenesis in B16F10 Melanoma Cells. *J Nat Prod* 2016; 79: 1091–1096
- [81] Tsoyi K, Jang HJ, Lee YS, Kim YM, Kim HJ, Seo HG, Lee JH, Kwak JH, Lee DU, Chang KC. (+)-Nootkatone and (+)-valencene from rhizomes of *Cyperus rotundus* increase survival rates in septic mice due to heme oxygenase-1 induction. *J Ethnopharmacol* 2011; 137: 1311–1317
- [82] Chavan MJ, Wakte PS, Shinde DB. Analgesic and anti-inflammatory activity of caryophyllene oxide from *Annona squamosa* L. bark. *Phytomedicine* 2010; 17: 149–151
- [83] Park KR, Nam D, Yun HM, Lee SG, Jang HJ, Sethi G, Cho SK, Ahn KS. β -Caryophyllene oxide inhibits growth and induces apoptosis through the suppression of PI3K/AKT/mTOR/S6K1 pathways and ROS-mediated MAPKs activation. *Cancer Lett* 2011; 312: 178–188
- [84] Tyagi AK, Prasad S, Yuan W, Li S, Aggarwal BB. Identification of a novel compound (β -sesquiphellandrene) from turmeric (*Curcuma longa*) with anticancer potential: comparison with curcumin. *Invest New Drugs* 2015; 33: 1175–1186
- [85] Hsu HC, Yang WC, Tsai WJ, Chen CC, Huang HY, Tsai YC. α -Bulnesene, a novel PAF receptor antagonist isolated from *Pogostemon cablin*. *Biochem Biophys Res Commun* 2006; 345: 1033–1038
- [86] Quintans Jde S, Menezes PP, Santos MRV, Bonjardim LR, Almeida JRGS, Gelain DP, Araújo AADS, Quintans-Júnior LJ. Improvement of *p*-cymene antinociceptive and anti-inflammatory effects by inclusion in β -cyclodextrin. *Phytomedicine* 2013; 20: 436–440
- [87] de Oliveira TM, de Carvalho RBF, da Costa IHF, de Oliveira GAL, de Souza AA, de Lima SG, de Freitas RM. Evaluation of *p*-cymene, a natural antioxidant. *Pharm Biol* 2015; 53: 423–428
- [88] Seo MJ, Kim SJ, Kang TH, Rim HK, Jeong HJ, Um JY, Hong SH, Kim HM. The regulatory mechanism of β -eudesmol is through the suppression of caspase-1 activation in mast cell-mediated inflammatory response. *Immunopharmacol Immunotoxicol* 2011; 33: 178–185
- [89] Plengsuriyakarn T, Karbwang J, Na-Bangchang K. Anticancer activity using positron emission tomography-computed tomography and pharmacokinetics of β -eudesmol in human cholangiocarcinoma xenografted nude mouse model. *Clin Exp Pharmacol Physiol* 2015; 42: 293–304
- [90] Kimura Y, Sumiyoshi M. Effects of an *Atractylodes lancea* rhizome extract and a volatile component β -eudesmol on gastrointestinal motility in mice. *J Ethnopharmacol* 2012; 141: 530–536
- [91] Chiou LC, Ling JY, Chang CC. Chinese herb constituent β -eudesmol alleviated the electroshock seizures in mice and electrographic seizures in rat hippocampal slices. *Neurosci Lett* 1997; 231: 171–174
- [92] Ohara K, Fukuda T, Ishida Y, Takahashi C, Ohya R, Katayama M, Uchida K, Tominaga M, Nagai K. β -Eudesmol, an oxygenized sesquiterpene, stimulates appetite via TRPA1 and the autonomic nervous system. *Sci Rep* 2017; 7: 15785
- [93] Peana AT, D'Aquila PS, Chessa ML, Moretti MDL, Serra G, Pippia P. (-)-Linalool produces antinociception in two experimental models of pain. *Eur J Pharmacol* 2003; 460: 37–41
- [94] Kuroda K, Inoue N, Ito Y, Kubota K, Sugimoto A, Kakuda T, Fushiki T. Sedative effects of the jasmine tea odor and (R)-(-)-linalool, one of its major odor components, on autonomic nerve activity and mood states. *Eur J Appl Physiol* 2005; 95: 107–114
- [95] Milde J, Elstner EF, Graßmann J. Synergistic inhibition of low-density lipoprotein oxidation by rutin, γ -terpinene, and ascorbic acid. *Phytomedicine* 2004; 11: 105–113
- [96] Passos F, Lopes E, de Araújo J, de Sousa D, Veras L, Leite J, Almeida F. Involvement of Cholinergic and Opioid System in γ -Terpinene-Mediated Antinociception. *Evid Based Complement Alternat Med* 2015; 2015: 829414
- [97] Giwanon R, Thubthimthed S, Rerk-am U, Sunthorntanasart T. Antimicrobial activity of terpinen-4-ol and sabinene. *Thai J Pharm Sci* 2000; 24: 27
- [98] Lahlou S, Interaminense LF, Leal-Cardoso JH, Duarte GP. Antihypertensive effects of the essential oil of *Alpinia zerumbet* and its main constituent, terpinen-4-ol, in DOCA-salt hypertensive conscious rats. *Fundam Clin Pharmacol* 2003; 17: 323–330
- [99] Tang WT, Fang MF, Liu X, Yue M. Simultaneous Quantitative and Qualitative Analysis of Flavonoids from Ultraviolet-B Radiation in Leaves and Roots of *Scutellaria baicalensis* Georgi Using LC-UV-ESI-Q/TOF/MS. *J Anal Methods Chem* 2014; 2014: 643879
- [100] Ringrose P, Parr M, McLaren M. Effects of anti-inflammatory and other compounds on the release of lysosomal enzymes from macrophages. *Biochem Pharmacol* 1975; 24: 607–614