



## Endoscopic diagnosis and management of chronic relapsing pancreatitis due to eroded embolization coils



► **Fig. 1** Endoscopic ultrasound showed a dilated main pancreatic duct (red arrow) and intraductal filling (white arrow).



► **Fig. 2** Endoscopic retrograde cholangiopancreatography showed coils (black arrow) and intraductal filling (yellow arrow). There was also diffuse main pancreatic duct dilation.



► **Fig. 3** Pancreatic stones seen on SpyGlass examination (Boston Scientific, Marlborough, Massachusetts, USA).

A 59-year-old man with alcohol-induced pancreatitis was referred due to dilated pancreatic duct and pancreatic mass. He had presented 9 years earlier with gastrointestinal bleeding secondary to hemocoelpancreaticus, which was treated by interventional radiology-guided coil and glue application to the superior pancreaticoduodenal artery pseudoaneurysm. He had complained of postprandial upper abdominal pain and a 10-lb weight loss, and had experienced recurrent acute pancreatitis in the preceding 4 months. Contrast-enhanced computed tomography showed dilated pancreatic duct and multiple coils around the head of the pancreas, and extensive shadowing artifact precluded further evaluation.

Endoscopic ultrasound revealed a dilated main pancreatic duct with intraductal filling and a 25×16mm hypoechoic lesion in the head of the pancreas near the coils (► **Fig. 1**). Cytology showed epithelioid cells with abundant debris and no evidence of malignancy.

Endoscopic retrograde cholangiopancreatography was performed. After biliary sphincterotomy, the pancreatic orifice was cannulated with a 3.9-Fr sphincterotome and 0.025-inch angled tip guidewire. A diffuse dilated pancreatic duct and large filling defect was seen on pancreatogram (► **Fig. 2**). Spyglass DS (Boston Scientific, Marlborough, Massachusetts, USA) was passed over the guidewire and multiple large white stones were revealed (► **Fig. 3**). The stones were fragmented using electrohydraulic lithotripsy. Multiple eroded coils were also seen in the proximal duct, from prior embolization (► **Fig. 4**). The coils were removed with SpyBite (Boston Scientific) and rat-tooth forceps (► **Video 1**). Two 7Fr×12cm single-pigtail plastic stents were deployed to maintain duct patency.



► **Fig. 4** SpyGlass examination (Boston Scientific, Marlborough, Massachusetts, USA) showed coils (red arrow) eroded into the pancreatic duct.

The patient tolerated the procedure well and was seen 1 month later, with marked improvement of symptoms and plan to follow up in 3 months.

Coils from prior embolization that have eroded into the gastrointestinal lumen and then either passed spontaneously or been removed endoscopically have been reported [1, 2]. To our knowledge, this is the first report of effective endoscopic management of recurrent pancreatitis caused by coils and glue expelled into the pancreatic duct.

Endoscopy\_UCTN\_Code\_TTT\_1AR\_2AK

### Competing interests

The authors declare that they have no conflict of interest.



**Video 1** Diagnosis and management of chronic relapsing pancreatitis due to eroded embolization coils.

## References

- [1] Soondoos R, Manju DC, Khaled A et al. Vascular coil erosion into hepaticojejunostomy following hepatic arterial embolization. *BMC Surg* 2015; 15: 51
- [2] Bohl JL, Dossett LA, Grau AM. Gastroduodenal artery pseudoaneurysm associated with haemosuccus pancreaticus and obstructive jaundice. *J Gastrointest Surg* 2007; 11: 1752–1754

## Bibliography

*Endoscopy* 2020; 52: E448–E449

**DOI** 10.1055/a-1163-7195

**ISSN** 0013-726X

**published online** 12.5.2020

© 2020. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

## The authors

**Wesam M. Frandah<sup>1</sup>, Nicolas Fiore<sup>2</sup>, Merall Sherif<sup>3</sup>, Saad Emhmed Ali<sup>4</sup>, Ahmed M. Sherif<sup>1</sup>**

- 1 Department of Gastroenterology and Therapeutic endoscopy, Banner health, Greeley, Colorado, United States
- 2 College of Osteopathic Medicine, Rocky Vista University, Parker, Colorado, United States
- 3 Division of Neuroscience, Colorado State University, Fort Collins, Colorado, United States
- 4 Department of Internal Medicine, University of Kentucky, Lexington, Kentucky, United States

## Corresponding author

**Wesam M. Frandah, MD**

Department of Gastroenterology, Banner Health, 2010 16th St. Ste A, Greeley, CO 80631, United States

Fax: +1-970-810-4475

wfrandah@gmail.com

## ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>