



Endoscopic submucosal dissection for gastric duplication cyst with heterotopic pancreas



► **Fig. 1** A subepithelial tumor with severe erythema and hemorrhagic spots was seen in the gastric lower body.



► **Fig. 2** On endoscopic ultrasonography, the lesion was anechoic, mainly confined to the submucosal layer, and a small branch-shaped hypoechoic portion was identified in the cystic wall (arrow).



► **Fig. 3** A clip attached with dental floss was deployed to secure the dissection plane between the tumor and the muscle layer.


A 63-year-old man presented with a gastric subepithelial tumor, incidentally detected on screening endoscopy. A subepithelial tumor, approximately 30 mm in size and with a positive cushion sign, was observed on the greater curvature of the gastric lower body (► **Fig. 1**). Severe erythema and hemorrhagic spots were seen on its surface. On endoscopic ultrasonography, the lesion was anechoic, mainly confined to the submucosal layer, and had a small branch-shaped hypoechoic portion in the cystic wall (► **Fig. 2**). Endoscopic submucosal dissection (ESD) was suggested in order to prevent bleeding (► **Video 1**).

After marking the tumor borders, a 0.9% saline solution with epinephrine and indigo carmine was injected into the submucosal layer. Subsequently, a circumferential mucosal incision was made with a dual knife and submucosal dissection was performed with an insulated-tipped knife. During dissection, the tumor was tightly attached to the proper muscle layer; therefore, a clip attached with dental floss was deployed to secure the dissection plane between the tumor and muscle layer (► **Fig. 3**). The tumor was safely and completely resected (► **Fig. 4**). Histopa-



► **Fig. 4** The inner surface of the resected specimen.

thologically, the cystic lesion was lined with columnar epithelial mucosa and had its own muscle layer, and ectopic pancreatic tissue of acinar structure was present in the cystic wall (► **Fig. 5**). The patient did not develop any adverse events and was discharged after 2 days. Gastric duplication cysts comprise about 2%–9% of gastrointestinal duplication cysts [1]. Most are cystic, occurring along the greater curvature of the stomach without communicating with it [2]. Ectopic pancreatic tissue is found in the



► **Fig. 5** Histopathologically, the cystic lesion was lined with columnar epithelial mucosa and had its own muscle layer (hematoxylin and eosin [H&E] stain, ×12.5). Ectopic pancreatic tissue of acinar structure was also present in the cystic wall (boxed area, H&E stain, ×100).

wall in up to 37% [3]. The conventional treatment for removal is surgical resection; ESD is an alternative modality. Due to the tight attachment of the cyst to the muscle layer, dental floss and clip traction is an efficient and safe method for en bloc resection.

Endoscopy_UCTN_Code_CCL_1AB_2AD_3AF



Video 1 Endoscopic submucosal dissection of a gastric duplication cyst.

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>

Competing interests

The authors declare that they have no conflict of interest.

The authors

Gwang Ha Kim¹, Moon Won Lee¹, Bong Eun Lee¹, Do Youn Park²

- 1 Department of Internal Medicine, Pusan National University School of Medicine and Biomedical Research Institute, Pusan National University Hospital, Busan, Korea
- 2 St. Mary Pathology Laboratory, Busan, Korea

Corresponding author

Gwang Ha Kim, MD, PhD

Department of Internal Medicine, Pusan National University School of Medicine and Biomedical Research Institute, Pusan National University Hospital, 179 Gudeok-ro, Seo-Gu, Busan 49241, Korea
 Fax: +82-51-2448180
 doc0224@pusan.ac.kr

References

- [1] Kim DH, Kim JS, Nam ES et al. Foregut duplication cyst of the stomach. *Pathol Int* 2000; 50: 142–145
- [2] Eom JS, Kim GH, Song GA et al. Gastric duplication cyst removed by endoscopic submucosal dissection. *Korean J Gastroenterol* 2011; 58: 346–349
- [3] Macpherson RI. Gastrointestinal tract duplications: clinical, pathologic, etiologic, and radiologic considerations. *Radiographics* 1993; 13: 1063–1080

Bibliography

Endoscopy 2021; 53: E19–E20

DOI 10.1055/a-1173-7326

ISSN 0013-726X

published online 29.5.2020

© 2020, Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany