


Closure of residual fistula after esophageal atresia repair in a 5-year-old using endoscopic submucosal dissection of surrounding mucosa

Esotracheal fistulas after esophageal atresia repair recur in 5 to 10% of cases [1,2] and lead to recurrent pneumonia or mediastinitis. Several surgical techniques are effective in closing an esotracheal fistula by thoracotomy or cervicotomy [3], but endoscopic success has never been published for this indication. We present the case of a 5-year-old patient with a previous history of type III esophageal atresia neonatal surgery, chronic respiratory congestion, and poor weight gain. She experienced a fistula recurrence with a large tracheoesophageal defect (► **Fig. 1**).

We performed an endoscopic fistula closure after prior endoscopic submucosal dissection (ESD) of the surrounding mucosa as previously reported for a button battery-induced esotracheal fistula [4] or idiopathic chronic fistula [5]. The patient underwent tracheal intubation with balloon placement just under the fistula. ESD was assisted by clip-and-line traction to dissect deeper into the fistula tract (► **Video 1**). Once the mucosa was resected, we closed the area using four clips anchored in the submucosa. A radiological check objectified the tightness of the closure.

The postoperative consequences were favorable, marked by a disappearance of the patient's respiratory symptoms during the following 3 months. A radiological check with opacification carried out 3 months before the gesture shows a tiny residual fistula and pseudo-diverticular scarring (► **Fig. 2**). The patient underwent a second procedure, during which ESD of the surrounding mucosa was done with deep cutting of the diverticular wall. Then, a new closure of the resected area was done. The closure of the residual fistula was confirmed by radiological control after 1 month.


Currently, the management of recurrent esotracheal fistulas after atresia surgery is not well defined. Endoscopic closure after ESD of the surrounding mucosa



► **Fig. 1** First endoscopic submucosal dissection to remove mucosa surrounding the fistula. **a** Appearance of the residual tracheoesophageal fistula (red arrow showing the fistula). **b, c** Deep dissection of the entire fistulous tract using a clip-and-line system. **d** Clips placed in the submucosa to close the orifice.



► **Video 1** Successful endoscopic closure of a residual fistula after esophageal atresia repair in a 5-year-old using endoscopic submucosal dissection of the surrounding mucosa.



► **Fig. 2** Aspect of esophageal transit before and after the second procedure. **a** Radiological opacification after first endoscopic closure: small residual fistula and pseudo-diverticular scarring (red arrow showing the fistula). **b** Radiological opacification after second endoscopic gesture: no residual fistula.

could allow a definitive resolution of the esotracheal fistulas and avoid a second risky surgery.

Endoscopy_UCTN_Code_TTT_1AO_2AI

Competing interests

The authors declare that they have no conflict of interest.

The authors

Grégoire Lavaud^{1,2}, Sophie Heissat¹, Auxane Chauveau², Noël Peretti¹, Alain Lachaux¹, Borathchakra Oung², Mathieu Pioche^{2,3}

- 1 Department of Pediatric Endoscopy and Gastroenterology, Hôpital Femme Mère Enfant, Lyon, France
- 2 Department of Endoscopy and Gastroenterology, Pavillon L, Edouard Herriot Hospital, Lyon, France
- 3 INSERM U1032, LabTau, Lyon, France

Corresponding author

Dr. Mathieu Pioche

Endoscopy Unit – Digestive Disease Department, Pavillon L – Edouard Herriot Hospital, 69437 Lyon Cedex, France
 Fax: +33-472-110-147
 mathieu.pioche@chu-lyon.fr

References

- [1] Kovesi T, Rubin S. Long-term complications of congenital esophageal atresia and/or tracheoesophageal fistula. *Chest* 2004; 126: 915–925
- [2] Bruch SW, Hirschl RB, Coran AG. The diagnosis and management of recurrent tracheoesophageal fistulas. *J Pediatr Surg* 2010; 45: 337–340
- [3] Daniel SJ, Smith MM. Tracheoesophageal fistula: open versus endoscopic repair. *Curr Opin Otolaryngol Head Neck Surg* 2016; 24: 510–515
- [4] Gruner M, Heissat S, Pitiot V et al. Successful endoscopic closure of a refractory button-battery tracheoesophageal fistula in a 3-year child using endoscopic submucosal dissection of the surrounding mucosa. *Endoscopy* 2017; 49: E212–E214
- [5] Bertrand G, Jacques J, Rivory J et al. Deep endoscopic submucosal dissection of a refractory tracheoesophageal fistula using clip-and-line traction: a successful closure. *Endoscopy* 2017; 49: 1278–1280

Bibliography

Endoscopy 2021; 53: E132–E133

DOI 10.1055/a-1216-0271

ISSN 0013-726X

published online 5.8.2020

© 2020. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>