


## Endoscopic ultrasound-guided angiotherapy in bleeding gastrointestinal stromal tumors with coil deployment and cyanoacrylate injection




► **Fig. 1** Endoscopic view of the ulcerated bleeding GI stromal tumor (GIST).



► **Video 1** Endoscopic ultrasound (EUS)-guided angiotherapy in two patients with bleeding GI stromal tumors with EUS-guided coil deployment and cyanoacrylate (CYA) injection targeting the feeding artery.

Hemorrhage is the most frequent (30–40%) and dangerous complication of gastrointestinal stromal tumors (GISTs), with surgery being the treatment of choice and emergency surgery having a significantly higher risk than elective procedures [1]. Conventional endoscopic treatment of tumoral lesions is challenging and efficacy limited because of the complex neoangiogenesis of these lesions, which are beyond the reach of endoscopic contact procedures [2]. Cases of endoscopic ultrasound (EUS)-guided therapy of bleeding GISTs with injection of cyanoacrylate (CYA) [3,4] or ethanol [5] have been published. To the best of our knowledge, EUS-guided coil deployment has not been reported in this setting. We present two cases of bleeding GISTs successfully treated with EUS-guided combined coil deployment and further injection of CYA targeting the feeding artery (► **Video 1**).


**Case 1:** a 72-year-old man was admitted for melena and hemodynamic instability. An ulcerated fundic mass with a necrotic center and a 5-mm intratumoral artery was observed (► **Fig. 1**). EUS-guided injection of CYA (Histoacryl; B. Braun, Melsungen, Germany) into the feeding artery (► **Fig. 2**) with a 22-gauge needle was the initial approach. However, a spurting



► **Fig. 2** Endoscopic ultrasound (EUS) image displaying the feeding artery of the bleeding GI stromal tumor (GIST) (arrows) shown in ► **Fig. 1**.

bleed into the tumor cavity after puncture was observed (► **Fig. 3**). Hemostasis was achieved by deploying a 6-mm × 20-

cm coil (Nester; Cook Medical, Limerick, Ireland) and injecting 2 mL of CYA with a 19-gauge needle (► **Fig. 4**).



► **Fig. 3** Spurting bleed (arrows) after the endoscopic ultrasound (EUS)-guided puncture with a 22-gauge needle.



► **Fig. 4** Endoscopic ultrasound (EUS)-guided deployed coil (arrows) with a 19-gauge needle.

**Case 2:** a 79-year-old woman was admitted for hematemesis and hypovolemic shock. An ulcerated fundic mass with a 4-mm artery was observed. EUS-guided therapy with a 19-gauge needle was performed employing a 6-mm × 14-cm coil, followed by injection of 2 mL of CYA. The clinical outcome in both patients was uneventful, and scheduled surgery took place 4 weeks later. EUS-guided combined angiotherapy targeting the culprit artery of bleeding GISTs maintained hemostasis until scheduled surgery. Moreover, although the follow-up in these two patients was short, EUS-guided angiotherapy may be a therapeutic option in patients with bleeding GISTs deemed unfit for surgery.

Endoscopy\_UCTN\_Code\_TTT\_1AS\_2AG

## Competing interests

The authors declare they have no conflict of interest.

## The authors

**Rafael Romero-Castro<sup>1,2</sup>, Victoria Alejandra Jimenez-Garcia<sup>1,2</sup>, Atsushi Irisawa<sup>3</sup>, Pedro Hergueta-Delgado<sup>1</sup>, Antonio Garrido-Serrano<sup>1</sup>, Reyes Aparcero-Lopez<sup>1</sup>, Angel Caunedo-Alvarez<sup>1</sup>**

- 1 Virgen Macarena University Hospital, Endoscopy Unit, Gastroenterology Division, Seville, Spain
- 2 Vithas Nisa Seville Hospital, Endoscopy Unit, Seville, Spain
- 3 Department of Gastroenterology, Dokkyo Medical University School of Medicine, Tochigi, Japan

## Corresponding author

**Rafael Romero Castro, MD**  
Virgen Macarena University Hospital,  
Endoscopy Unit, Gastroenterology Division,  
Calle Dr. Fedriani, 3, 41009 Seville, Spain  
Fax: +34 955 00 88 05  
rafaromecas@hotmail.com

## References

- [1] Liu Q, Kong F, Zhou J et al. Management of hemorrhage in gastrointestinal stromal tumors: a review. *Cancer Manag Res* 2018; 10: 735–743
- [2] Venezia G, Asnaghi G, Manca A et al. Successful use of Hemospray as a bridge to surgery in life-threatening bleeding from a gastrointestinal stromal tumor of the stomach. *Endoscopy* 2016; 48: E232–E233
- [3] Levy MJ, Wong Kee Song LM, Farnell MB et al. Endoscopic ultrasound (EUS)-guided angiotherapy of refractory gastrointestinal bleeding. *Am J Gastroenterol* 2008; 103: 352–359

- [4] Kumbhari V, Gondal B, Okolo Iii PI et al. Endoscopic ultrasound-guided angiotherapy of a large bleeding gastrointestinal stromal tumor. *Endoscopy* 2013; 45: E326–E327
- [5] Valdivielso Cortazar E, Fernandez-Urien I, Vila Costas JJ et al. Endoscopic ultrasound-guided therapy of a gastrointestinal stromal tumor (GIST). *Endoscopy* 2015; 47: E262–E263

## Bibliography

*Endoscopy* 2021; 53: E124–E125  
**DOI** 10.1055/a-1216-0634  
**ISSN** 0013-726X  
**published online** 24.7.2020  
© 2020. Thieme. All rights reserved.  
Georg Thieme Verlag KG, Rüdigerstraße 14,  
70469 Stuttgart, Germany

### ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at  
<https://mc.manuscriptcentral.com/e-videos>