This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.

A novel endoscopic submucosal dissection training model using skinned chicken meat with gel



► Fig. 1 Setting the skinned chicken meat with an electrode in a 2-L polyethylene terephthalate bottle.



Video 1 Endoscopic submucosal dissection training model using chicken meat with gel.



► Fig. 2 Endoscopic image showing circular markings on the chicken skin.



► **Fig. 4** Endoscopic image showing a dissection view of the gel layer.



► Fig. 5 Chicken skin after resection.



▶ Fig. 3 Endoscopic image showing lifting of the chicken skin by injections of two solutions and creating a gel layer.

Gastric endoscopic submucosal dissection (G-ESD) is performed worldwide. However, some trainees are not offered sufficient opportunities to perform G-ESD. Therefore, hands-on seminars using porcine stomachs are actively held, serving as valuable learning opportunities for trainees [1]. However, preparing porcine stomachs is time-consuming and effort-intensive, and leads to problems related to sanitation and pollution. To address these issues, we developed a new G-ESD training model using skinned chicken meat with gel involving the following steps (**Video 1**). (1) An electrode is attached to the chicken meat on

the skinless slide (▶ Fig. 1). (2) The chicken meat is placed in a 2-L polyethylene terephthalate bottle (▶ Fig. 1). (3) Circular marks are made to mimic a virtual tumor region (▶ Fig. 2). (4) The first injection solution is made by mixing 40 mL of clear-type polyvinyl alcohol glue with 30 mL of water. The second injection solution is made by dissolving 3 g of borax in 50 mL of warm water and adding 3 g of sodium hydrogen carbonate. These two solutions are injected subcutaneously, and skin lifting is observed, similarly to that of the gastric mucosa (▶ Fig. 3). (5)

The skin is cut using a knife, and the cut is gradually widened (▶ Fig. 4, ▶ Fig. 5). The two liquids are mixed and injected under the chicken skin (to mimic the submucosal layer). The greatest advantages of this model are that i) skinned chicken meat can be purchased from any supermarket, ii) there is a low infection risk for hepatitis E virus or other zoonotic infections, and iii) it is a low-cost model (\$1.50 - 1.70). Chicken meat preparation is fast, simple, and inexpensive compared with porcine model preparation. Therefore, G-ESD training using this novel model is feasible, efficient, and may be widely available for trainees who do not have the chance to practice G-ESD or attend hands-on seminars.

Endoscopy_UCTN_Code_TTT_1AU_2AB

Competing interests

The authors declare that they have no conflict of interest.

The authors

Sho Suzuki, Hiroshi Kawakami, Tadashi Miike, Yutaro Ishikawa, Tomoya Hirata, Naoki Ichinari, Anna Komura

Department of Gastroenterology and Hepatology, Center for Digestive Disease and Division of Endoscopy, University of Miyazaki Hospital, Miyazaki, Japan

Corresponding author

Hiroshi Kawakami, MD, PhD

Department of Gastroenterology and Hepatology, Center for Digestive Disease, Division of Endoscopy, University of Miyazaki Hospital, 5200 Kihara, Kiyotake, Miyazaki 889-1692, Japan

Fax: +81-985-859802 hiropon@med.miyazaki-u.ac.jp

Reference

[1] Berr F, Ponchon T, Neureiter D et al. Experimental endoscopic submucosal dissection training in a porcine model: learning experience of skilled Western endoscopists. Dig Endosc 2011; 23: 281–289

Bibliography

Endoscopy 2021; 53: E334–E335

DOI 10.1055/a-1287-8567
ISSN 0013-726X
published online 3.11.2020
© 2020. Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS https://eref.thieme.de/e-videos



Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

https://mc.manuscriptcentral.com/e-videos