


## Endoscopic retrieval of a huge gastric trichobezoar using an electrosurgical knife



► **Fig. 1** Endoscopic image showing a large gastric trichobezoar in a 22-year-old patient.



► **Fig. 2** The trichobezoar was broken up using an electrosurgical knife.




► **Fig. 3** The trichobezoar completely removed after fragmentation.

A trichobezoar is a rare type of bezoar and usually located in the stomach; however, sometimes it extends through the pylorus into the small bowel, even reaching the transverse colon [1]. Trichobezoars may cause potentially life-threatening complications, such as intestinal obstruction, gastric bleeding, and perforation. Epigastric surgical incision is the most common method of large trichobezoar removal [2]. Herein, we report a case of successful endoscopic retrieval of a trichobezoar after its fragmentation using an electrosurgical knife.

A 22-year-old woman visited the primary clinic with a history of upper abdominal pain and early satiety. She had been habitually eating her own hair since childhood. The endoscopic findings showed a large, densely packed intragastric trichobezoar (dark hair with hard mass) approximately 6 cm × 15 cm in size extending through the pylorus, and a shallow ulcer in the body (► **Fig. 1**). We then decided to remove the trichobezoar endoscopically.

With the patient under conscious sedation with midazolam plus propofol, we used a two-channel gastroscope (GIF-2TQ260M; Olympus, Tokyo, Japan) and tried to fragment and remove the trichobezoar using grasping forceps through an overtube. As the lump was huge,



► **Video 1** Endoscopic retrieval of a large gastric trichobezoar after fragmentation into several pieces using an electrosurgical knife.

dense, and tangled in hair, it could not be removed even after repeated attempts. A subsequent attempt to cut the trichobezoar using argon plasma coagulation and a polypectomy snare also failed to fragment it efficiently (► **Fig. 2**). Finally, we used an electrosurgical knife (IT knife 1; Olympus, Tokyo, Japan) to cut the trichobezoar effectively into two pieces (► **Video 1**). The smaller piece was further fragmented and then successfully removed using grasping forceps (► **Video 1**). The remaining piece was successfully removed in two

sessions; each session took about an hour (► **Fig. 3**). The clinical course was tolerable. After trichobezoar removal, the patient was free of pain; she was given a regular diet and discharged.

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AL

### Competing interests

The authors declare that they have no conflict of interest.

## The authors

**Seung Gyun Baek<sup>1</sup>, Chi Hyuk Oh<sup>1</sup>, Ga Young Shin<sup>2</sup>, Jung-Wook Kim<sup>1</sup>, Jae-Young Jang<sup>1</sup>**

- 1 Division of Gastroenterology and Hepatology, Department of Internal Medicine, Kyung Hee University Hospital, Kyung Hee University School of Medicine, Seoul, Republic of Korea
- 2 Department of Medicine, Graduate School, Kyung Hee University, Seoul, Republic of Korea

## Corresponding author

### **Chi Hyuk Oh, MD**

Division of Gastroenterology and Hepatology, Department of Internal Medicine, Kyung Hee University Hospital, 23 Kyungheedaero, Dongdaemun-gu, Seoul 02447, Republic of Korea  
Fax: +82-2-9681848  
ochihyuk@gmail.com

## References

- [1] Vaughan ED Jr, Sawyers JL, Scott HW Jr. The Rapunzel syndrome. An unusual complication of intestinal bezoar. *Surgery* 1968; 63: 339–343
- [2] Benatta MA. Endoscopic retrieval of gastric trichobezoar after fragmentation with electrocautery using polypectomy snare and argon plasma coagulation in a pediatric patient. *Gastroenterol Rep (Oxf)* 2016; 4: 251–253

## Bibliography

*Endoscopy* 2021; 53: E357–E358

**DOI** 10.1055/a-1293-6677

**ISSN** 0013-726X

**published online** 19.11.2020

© 2020. Thieme. All rights reserved.  
Georg Thieme Verlag KG, Rüdigerstraße 14,  
70469 Stuttgart, Germany

## ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>