



Successful endoscopic papillectomy with intrapancreatic ductal radiofrequency ablation for ampulla cancer in surgically altered anatomy

Surgical resection of ampullary neoplasms in patients with surgically altered anatomy is challenging [1]. The recent development of intraductal radiofrequency ablation (RFA) devices and endoscopic techniques enabled endoscopic treatments to be used as viable alternatives for ampullary neoplasm in poor surgical candidates [2–4]. We describe a case of endoscopic papillectomy with intraductal RFA for an ampullary tumor in a patient with Billroth II anastomosis.



A 75-year-old woman was referred for a polypoid mass on the ampulla of Vater. She had undergone Billroth II gastrectomy for gastric cancer 28 years before and bile duct resection with right lobectomy for cholangiocarcinoma 3 years before. We were able to approach the 4.5-cm polypoid mass with conventional duodenoscopy (JF-260V; Olympus Corp., Tokyo, Japan) (► Fig. 1). Biopsy revealed a tubulovillous adenoma. An endoscopic piecemeal resection was performed considering the size of the mass (► Fig. 2). The pathology report revealed a small, multifocal portion of a well-differentiated papillary adenocarcinoma in the background of the tubulovillous adenoma involving the resection margin. We thus performed repetitive endoscopic resection with argon plasma coagulation (APC) for the small remnant adenoma. After 1 month, the endoscopy revealed a remnant tumor at the pancreatic duct orifice. We then performed intrapancreatic ductal RFA for intraductal extension. After pancreatographic evaluation, an RFA catheter (ELRA; Taewoong Medical, Gimpo-si, South Korea) was introduced by a guidewire. RFA (70 W, target temperature 80°C) was applied for 30 seconds under fluoroscopic guidance using a VIVA generator (Taewoong Medical) (► Fig. 3, ► Video 1). After intrapancreatic ductal RFA, we inserted a plastic stent in the pancreatic duct with additional APC around the pancreatic duct orifice.



► Fig. 1 Endoscopic view of the 4.5-cm polypoid ampullary tumor.



► Fig. 2 Endoscopic piecemeal papillectomy for the ampullary tumor.



► Fig. 3 a Endoscopic view of the radiofrequency ablation (RFA) catheter ablating the surrounding mucosa and intraductal lesion. b Fluoroscopic view of the RFA catheter located in the proximal pancreatic duct.

The patient was discharged without complications. No remnant tumor was found at the 7-month follow-up (► Fig. 4). Endoscopic papillectomy with intrapancreatic ductal RFA may be performed for ampullary neoplasm with intraductal extension in patients with surgically altered anatomy.


Endoscopy_UCTN_Code_TTT_1AR_2AK

Competing interests

The authors declare that they have no conflict of interest.



▶ Video 1 Endoscopic papillectomy and intrapancreatic ductal radiofrequency ablation. The large ampullary tumor was successfully managed with repetitive endoscopic resection and intrapancreatic ductal radiofrequency ablation.



▶ Fig. 4 Endoscopic view at follow-up.

The authors

Jung Su Lee¹, Hoonsub So², Jun Seong Hwang², Sung Woo Ko², Tae Jun Song²

- 1 Division of Gastroenterology, Department of Internal Medicine, Ilsan Paik Hospital, Inje University College of Medicine, Goyang, Republic of Korea
- 2 Division of Gastroenterology, Department of Internal Medicine, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Republic of Korea

Corresponding author

Tae Jun Song, MD

Division of Gastroenterology, Department of Internal Medicine, Asan Medical Center, University of Ulsan College of Medicine, 88, Olympic-ro 43-gil, Songpa-gu, Seoul 05505, South Korea
Fax: +82-2-3010-6517
drsong@amc.seoul.kr

References

- [1] Lee D, Lee JH, Choi D et al. Surgical strategy and outcome in patients undergoing pancreaticoduodenectomy after gastric resection: a three-center experience with 39 patients. *World J Surg* 2017; 41: 552–558
- [2] Hintze RE, Adler A, Veltzke W et al. Endoscopic access to the papilla of Vater for endoscopic retrograde cholangiopancreatography in patients with Billroth II or Roux-en-Y gastrojejunostomy. *Endoscopy* 1997; 29: 69–73
- [3] Park TY, Kang JS, Song TJ et al. Outcomes of ERCP in Billroth II gastrectomy patients. *Gastrointest Endosc* 2016; 83: 1193–1201
- [4] Rustagi T, Irani S, Reddy DN et al. Radiofrequency ablation for intraductal extension of ampullary neoplasms. *Gastrointest Endosc* 2017; 86: 170–176

Bibliography

Endoscopy 2022; 54: E73–E74
DOI 10.1055/a-1388-5717
ISSN 0013-726X
published online 15.3.2021
© 2021. Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS
<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>