New Paradigms for Familiar Diseases: Lessons Learned on Circulatory Bacterial Signatures in Cardiometabolic Diseases

Authors

Rima Chakaroun^{1, 2}, Lucas Massier^{1, 3}, Niculina Musat⁴, Peter Kovacs^{1, 5}

Affiliations

- 1 Medical Department III Endocrinology, Nephrology, Rheumatology, University of Leipzig Medical Center, Leipzig, Germany
- 2 Wallenberg Laboratory, Department of Molecular and Clinical Medicine and Sahlgrenska Center for Cardiovascular and Metabolic Research, University of Gothenburg, Gothenburg, Sweden
- 3 Department of Medicine (H7), Karolinska Institutet, Stockholm, Sweden
- 4 Department of Isotope Biogeochemistry, Helmholtz Centre for Environmental Research – UFZ, Leipzig, Germany
- 5 Deutsches Zentrum für Diabetesforschung eV, Neuherberg, Germany

Key words

microbiome, type 2 diabetes, cardiovascular disease, blood microbiome, metabolic disease

received 12.10.2021 revised 15.12.2021 accepted 24.01.2022 published online 23.03.2022

Bibliography

Exp Clin Endocrinol Diabetes 2022; 130: 313–326 DOI 10.1055/a-1756-4509 ISSN 0947-7349 © 2022. Thieme. All rights reserved. Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Correspondence

Rima Chakaroun Leipzig University Hospital Department III – Clinic for Endocrinology Nephrology, and Rheumatology Wallenberg Laboratory Sahlgrenska University Hospital S-413 45 Gothenburg Sweden rima.chakaroun@wlab.gu.se Peter Kovacs Leipzig University Hospital Department III – Clinic for Endocrinology Nephrology, and Rheumatology peter.kovacs@medizin.uni-leipzig.de

ABSTRACT

Despite the strongly accumulating evidence for microbial signatures in metabolic tissues, including the blood, suggesting a novel paradigm for metabolic disease development, the notion of a core blood bacterial signature in health and disease remains a contentious concept. Recent studies clearly demonstrate that under a strict contamination-free environment, methods such as 16 S rRNA gene sequencing, fluorescence in-situ hybridization, transmission electron microscopy, and several more, allied with advanced bioinformatics tools, allow unambiguous detection and quantification of bacteria and bacterial DNA in human tissues. Bacterial load and compositional changes in the blood have been reported for numerous disease states, suggesting that bacteria and their components may partially induce systemic inflammation in cardiometabolic disease. This concept has been so far primarily based on measurements of surrogate parameters. It is now highly desirable to translate the current knowledge into diagnostic, prognostic, and therapeutic approaches.

This review addresses the potential clinical relevance of a blood bacterial signature pertinent to cardiometabolic diseases and outcomes and new avenues for translational approaches. It discusses pitfalls related to research in low bacterial biomass while proposing mitigation strategies for future research and application approaches.

Introduction

In the light of our long co-evolution with microbes [1], the human microbiome can be seen as an accessory genome including diverse bacteria, viruses, archaea, and fungi, expanding over several bodily niches and extending our functional potential. While most bacteria are commensal, many inhabiting our body are mutualistic, and some are detrimental. They are kept in check by the immune system and the surrounding cell communities contributing to the compositional niche, whose survival depends on the livelihood of the entire "human meta-organism". After a period of what can be considered a prolonged cold war with bacteria, brought on by the advent of the hygiene hypothesis [2] and the golden era of the antibiotics, growing evidence of microbial "ecosystem services" [3], the inextricable link between our health and our microbiome, including efficacy and response to medication such as cardiac drugs and chemotherapeutics, have led to a rekindling of interest in mucosa-associated bacterial microbiomes [4-6].

For the last two decades, this area of research has been propelled forward by several consortial efforts, including MetaHit [7], Human Microbiome Project (HMP) [8], Metacardis [9, 10], and Flemish Gut Flora Project [8, 11], in which the intestinal microbiome and, in a minority of cases (HMP), the oral, dermal and vaginal microbiome have been sequenced in large cohorts to define: a) a "healthy microbiome state", b) factors contributing to microbiome composition (including medication, nutrition, and environment)[11-13], and c) the associations of these microbiomes with disease [7, 14]. This has been further facilitated and expedited by the introduction of whole metagenome shotgun sequencing as well as accessible and scalable analyses methods. These population studies have linked dysbiosis - a state of deviation from a core functional and taxonomic composition- with the pathogenesis of several diseases, including obesity, non-alcoholic fatty liver disease, and cardiovascular disease [15]. While a transient presence of bacteria in the blood has been described after teeth brushing or gingival manipulation [16, 17], their consistent presence has been interpreted as indicative of infection or sepsis. This notion changed only recently with the description of low grade or subclinical inflammation in metabolic disease.

Low-grade inflammation is consensually perceived as a potential trigger for metabolic dysfunction, insulin resistance, and diabetes [18] and is often reflected in subtle increases in systemic inflammatory markers such as C-reactive-Protein (CRP). Increased translocation of lipopolysaccharides (LPS), mainly from the gut to the blood, defined as endotoxemia [19], has also been described as a potential trigger for chronic inflammation in metabolic disease, supported by the onset of insulin resistance in healthy subjects after LPS-transfusion [19, 20]. Therefore, it is surprising that the notion of a core blood bacterial signature in health and its divergence in disease with accompanying systemic response remains contentious. However, converging evidence for microbial signatures in metabolic tissues, including the blood, suggests a novel paradigm for metabolic disease development [19-24]. In this review, we address the potential clinical relevance of a blood bacterial signature pertinent to disease development and outcomes and new avenues for translational approaches. We, moreover, discuss pitfalls related to research in low bacterial biomass while suggesting mitigation strategies for future research and application approaches.

Schrödinger's blood bacteria: 'you don't see me, now you do.'

The concept of bacterial presence in the blood has been reported as early as 1969, where living and metabolically active bacteria [25] were localized in the blood of healthy human subjects. In the meantime, a commendable body of work has combined classic molecular and microbiological techniques to evidence the presence of bacterial cells and sequences in the blood of presumably healthy humans, either as a study cohort or as healthy controls for specific disease states (recently reviewed in [26]). These methods include quantitative polymerase chain reaction (qPCR) [27-35] of 16S rRNA or targeted bacterial genes [29, 35], fluorescence in-situ hybridization (FISH) and other employment of fluorescent probes [35], transmission electron microscopy [36], dark field microscopy [36], PCR followed by electrospray ionization-mass spectrometry [37] as well as classical bacterial culture [32, 38, 39]. Several of these results have been met with criticism, considering the longstanding dogma of the blood being an innocuous environment and controversy regarding visual confirmation of bacteria deemed by some authors as L-form bacteria and by others as merely membrane vesicles or aggregated proteins [40].

The advent of efficient and scalable technologies, including next-generation sequencing of the 16 S rRNA gene (present in all bacterial cells), whole metagenome sequencing, RNA-sequencing [38, 40, 41], and the ongoing expansion of microbial reference genomes have facilitated the detection of non-culturable or hard-toculture bacteria across several tissues and diseases. This is especially important considering several blood-borne bacteria can persist in a dormant state [42]. Moreover, the democratization of analytical tools, including the development of microbiome tools with friendly graphic user interfaces [43] and comprehensive documentation [44], have made this research area accessible to a broader scientific community.

Blood bacterial signatures in health and cardiometabolic disease

While the role of systemic inflammation in cardiometabolic disease, partially induced by bacteria, has been widely appreciated, most of the published studies related this observed systemic inflammation to bacterial components and/or selectively analyzed surrogate parameters. This includes measurements of host response patterns (e.g., LPS binding protein (LBP)) or wall components of gram-negative bacteria (LPS) in the circulation to assess bacterial burden indirectly. However, there is scarce but increasing research investigating direct bacterial presence, that is, bacterial cells or bacterial nucleic acids, and its quantitative and qualitative potential to contribute to inflammation and dysfunction of both the immediate microenvironment and the whole system. Several studies have evidenced the presence of bacterial genetic material, whole bacteria, or even bacterial RNA in the blood of healthy subjects [26, 42] (> Fig. 1a), and research dissecting the contribution of circulating bacteria or bacterial genetic material to cardiometabolic disease has been accumulating steadily (> Fig. 1b). From a clinical perspective, it has been noted early on that bacteremia of unknown origin was more prevalent in subjects with type 2 diabetes (T2D) compared to non-diabetic subjects [45], with higher occurrences of Staphylococcus aureus and Klebsiella in the blood along



This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.



Fig. 1 a) Overview of phyla proportions, controls inclusion, and the number of included subjects in selected study cohorts reporting blood bacterial signatures in healthy cohorts and healthy subjects of cohorts with cardiometabolic and cardiovascular disease. Names of the studies are to the left of the bar. Bar length equals 100% (total proportions of reported phyla), and each color in the bar represents a specific phylum (color code specified in the legend to the right). Controls: specifies whether experimental and collection controls were included to account for contamination. A red box corresponds to a lack of controls, a green box to the availability of adequate controls. Ref.: refers to the reference number in the bibliography (Paisse and Gosiewski et al. are reviewed in ref [25]) and N the number of healthy subjects in the cohort. An average distribution for all studies is given in the lowest bar, and N on the right refers to the total number of subjects included. b) Overview of phyla proportions, included controls, and the number of included subjects in selected study cohorts reporting blood bacterial signatures in disease (focusing on the cardiometabolic disease). Names of the studies are to the left of the bar. Bar length equals 100% (total proportions of reported phyla), and each color in the bar represents a specific phylum (color code specified in the legend to the right). Controls: specifies whether experimental and collection controls were included to account for contamination. A red box corresponds to a lack of controls, a green box to the availability of adequate controls. Ref.: refers to the medical conditions of the cases in the studies. An average distribution for all studies is given in the lowest bar, and N on the right). Controls: specifies whether experimental and collection controls were included to account for contamination. A red box corresponds to a lack of controls, a green box to the availability of adequate controls. Ref.: refers to the reference number in the bibliography and N

relatively 'harmless infections' such as urinary tract infections, leading to septic shock more often (OR 1.9) [46] and motivating the early application of targeted empiric antibiotic treatment. A hallmark study was published in 2011 by Amar et al., analyzing blood bacterial load by measuring 16 S rRNA copy numbers in more than 3000 subjects followed over nine years. The authors report an increased abundance of Proteobacteria in visceral obesity as well as onset of T2D [30] and cardiovascular disease at follow-up [47]

In 2014, Sato et al. undertook a nested case-control study reporting that prevalent T2D was associated with an increased inflammatory response with higher levels of LBP independently linked to HbA1c and BMI. Higher abundances of *Clostridium coccoides, Atopobium* clusters [31, 47], *Sediminibacterium* spp.[48], and higher detection of the 16 S rRNA gene was positively associated with the disease [31]. This is particularly important, as increased bacterial DNA detection after bariatric surgery in subjects with T2D and obesity revealed a blunted response to the intervention with minor improvement in glucose tolerance or inflammation [49]. In contrast,

subjects with a high abundance of the genus *Bacteroides* were less likely to develop T2D during a two-year follow-up [48,49].

Likewise, there is increasing evidence for a link between circulating bacteria in liver fibrosis and steatosis, which is associated with an impaired immune system and glucose metabolism and major cardiovascular events [50]. In 2016, Lelouvier et al. reported increased levels of circulating 16 S rDNA and proportions of Proteobacteria (specifically Sphingomonas, Bosea, and Bradyrhizobiaceae taxa) in liver fibrosis. This was further substantiated by other studies showing similar results in liver cirrhosis in several blood compartments [51] and the increased detection of bacterial DNA via in situ hybridization in particularly severe cases of liver decompensation [52]. Chronic kidney disease (CKD), a well-established cardiovascular risk factor [53], has similarly been linked to changes in blood bacterial signature, as it was associated with a decreased alpha diversity - a hallmark of several non-communicable diseases - and the expansion of bacterial taxa expressing uricase as well as indole- and p-cresyl-forming uremic enzymes [54]. These bacte-

rial uremic toxins have been linked to worsening kidney function, endothelial dysfunction, cardiac fibrosis, macrophage activation, and insulin resistance [55]. CKD is also linked to a dysbiosis in the gut, which has been associated with impairment of gut epithelial barrier via depletion of claudin-1, occludin, and zonula occludens-1 proteins in addition to dysfunction of colonic T-regulatory cells [56] as well as further aggravation by direct effects of uremic toxins [56, 57]. Therefore, it is not surprising that CKD is reflected in a decreased bacterial diversity in the blood and associated with a higher abundance of proteobacteria, which were inversely correlated with kidney function [58]. Comprehensive studies on the specific compartmentalization of bacterial DNA in metabolic disease remain rare but promise a more holistic approach. For example, Anhê, Jensen et al. compared the bacterial DNA load and taxonomy in plasma samples, liver, subcutaneous, visceral, and mesenteric adipose tissues, noting a relatively reduced bacterial load in plasma samples compared to other tissues and a preferential presence of two specific genera (Rodoferax and Polaromonas) in the blood, but did not find distinct differences in the tissues between subjects with and without T2D [23]. Results were supported and expanded by findings from our research group, where 16 S rRNA gene content was quantified and sequenced in 75 patients with obesity and with or without T2D. Bacterial composition in the blood was associated with circulating tumor necrosis factor-a, CRP, LBP, and interleukin-6. Moreover, amounts of Lactococcus in the blood correlated negatively with homeostasis model assessment-estimated insulin resistance while Acinetobacter and Tahibacter were found more often in the blood than in several studied adipose tissues and correlated positively with diabetes status [21]. In another recent study, we evidenced reduced bacterial diversity in the blood of subjects with T2D vs. non-T2D, but a reduced richness after bariatric surgery paralleled with an increased bacterial quantity. The latter was surprisingly negatively associated with leukocyte counts. We also evidenced early shifts in taxa after surgery with a reduction in Rhizobacter, Anoxybacillus, Streptococcus, and Ureibacillus. In our cohort, several of these genera were positively correlated with inflammation, fat mass, and body mass index (BMI). Most importantly, 40% of the variance in the blood bacterial composition could be explained by 27 host variables, including medication intake [22]. Similarly, bacterial composition predicted the clinical classification of patients according to metabolic disease robustly. The strength of the last three studies is the extensive experimental and bioinformatic contaminant control, which has since become more common in recent studies [59].

At the other end of the cardiometabolic disease spectrum, infections in the hospital setting as well as outpatient infections very often preceded coronary heart disease (CHD) and stroke in the ARIC study, which analyzed cardiovascular risk in 15792 subjects, showing that in a total of 1312 incident CHD and 727 incident stroke cases, infections were associated with an up to 12.83 OR for CHD (inpatient infections, up to 14 days prior to CHD presentation) [60]. Increased strain can lead to overt manifestations of cardiovascular events in predisposed subjects. This can be exemplified in type 2 myocardial infarction (MI) in infection and anemia, where an imbalance between myocardial oxygen demand and supply becomes apparent. That being said, one can not preclude that systemic inflammation in the realm of infection can further trigger immunological processes in local sites of preexistent vascular lesions in the heart or carotid arteries. In line with a potential mediating effect of blood bacteria and increased bacterial translocation from the gut, Zhou et al. demonstrated higher levels of LPS and D-lactate (bacteria-derived) in the blood of subjects two days post-MI along with an increased alpha diversity in ST-elevation MI (STEMI) and enrichment of gut-derived bacteria. At the same time, chronic CHD presented with lower bacterial diversity in the blood compared to controls. Mediation analysis revealed a mediating role for bacterial translocation in inflammation (CRP and monocytes), left ventricular ejection fraction, and major adverse cardiovascular events. More importantly, in an experimental MI mouse model, the reversal of bacterial translocation via antibiotic treatment reduced serum LPS, alleviated monocytosis, and reduced cardiomyocyte injury post-infarction [61]. More recently, Amar et al. showed that subjects with an acute onset MI displayed a 1.3-fold increase in blood bacterial load compared to control subjects with concurrent metabolic risk factors, which was driven by high LDL cholesterol. In the 99 subjects with MI, several genera known to metabolize cholesterol, such as Nocardiaceae, Aerococcaceae, Gordonia, Propionibacterium, Chryseobacterium, and Rhodococcus, were depleted [62]. In conclusion, most studies found a similar pattern of phyla distribution in the circulation, which was dominated by Proteobacteria (~65%), followed by Actinobacteria (16%), Firmicutes (10%), and Bacteroidetes (4%) (▶ Fig. 1). With some exceptions, changes in cardiovascular disease (CVD) followed the same trend, showing a relative increase of Proteobacteria by 7% and a 10% reduction in Firmicutes on average (> Fig. 2). As depicted, individual studies show more considerable changes but are not always congruent. No objective conclusion can be drawn on lower taxonomic levels, as reported results vary greatly, but for example, Pseudomonadaceae and Streptococcus were enriched in CVD in more than two studies (► Table 1).

Translational avenues: are we there yet?

High throughput sequencing of blood-borne bacteria and other microorganisms is relevant as a potential extension of the repertoire of current assessments in health and disease. It has already shown paradigmal potential for the gut microbiome and promises a similar potential for blood-derived disease markers [15] or liquid biopsies [24] in non-communicable and infectious diseases. Moreover, it will be critical for downstream applications, including novel, cost-effective techniques to detect new pathogens in an everevolving environment [63, 64]. Exploring functional potential using highly granular whole genome sequencing and genome assembly paired with metabolomics and metaproteomics will further underpin disease mechanisms [65], allowing targeted and possibly personalized treatment options.

First things first: traditional applications for non-classical diagnostics

A seemingly immediate application for the analysis of bacterial sequences in the blood, along with other bodily niches, is the potential to precisely identify bloodstream pathogens at the strain level. This helps account for pathogenicity, transmission potential, and antimicrobial resistance [66] and define their source early on in order to inform targeted interventions in clinical infection management. Considering that the latter is in most cases dictated by as-



• Fig. 2 Heatmap referring to the differential abundance [increased (in red) or decreased (in blue)] of phyla in studies with disease vs. healthy controls. Studies are given to the left of the heatmap rows; phyla constitute the columns of the heatmap. A look at differentially abundant genera in CVD in selected studies are shown in the red box (for increased genera) and the blue box (for reduced genera) next to the survey referred. Abbreviation: CVD: cardiovascular disease.

Author	Method	Matrix	Conditions and number	Year	Reference
Rajendhran	16 S V3	Whole blood	133 CVD vs 118 controls, thereof positive for 16S: 31 vs 10 (not matched)	2013	[153]
Dinakaran	Metagenomic, 16S quantification	Plasma	80 CVD vs 40 controls for quantification, 3 vs 3 for sequencing	2014	[29]
Lelouvier	16S V3/V4	Buffy coat	Cohort 1: 11 fibrosis vs 26 controls; Cohort 2: 11 fibrosis vs 60 controls	2016	[154]
Whittle	16 S V 4	Plasma	5 atopic asthmatic vs 5 controls	2018	[38]
Schierwagen	165	Buffy coat	3 variceal bleeding, 4 refractory ascites	2018	[51]
Zhou	16 S V 4	Buffy coat	100 ST-segment elevation myocardial infarction, 50 CHD, 49 controls	2018	[61]
Qiu	16S V5/V6	Whole blood	50 T2D vs 100 healthy (matched for age, sex)	2019	[48]
Alvarez-Silva	165 V3/V4	Whole blood and ascitic fluid	33 cirrhosis	2019	[155]
Shah	16 S V3/V4	Buffy coat	20 CKD vs 20 healthy (not matched)	2019	[58]
Amar	16 S V3/V4	EDTA blood	103 high CVD risk, 99 with myocardial infarction	2019	[62]
Anhê	16 S V3/V4	Plasma	20 T2D, obese vs 20 normal glucose tolerance, obese	2020	[23]
Massier	16 S V4/V5, CARD-FISH	EDTA blood	75 with obesity, 42 no T2D vs 33 T2D	2020	[21]
Chakaroun	16 S V4/V5, CARD-FISH	EDTA blood	64 at baseline with obesity, 24 no T2D vs 24 T2D with 12-month follow-up (matched for age, sex, and BMI)	2021	[22]
CARD-FISH: catalyzed reporter deposition – fluorescence in situ hybridization; CVD: cardiovascular disease; CHD: chronic heart disease; CKD: Chronic kidney disease; EDTA: ethylenediaminetetraacetic acid; T2D: type 2 diabetes					

Table 1 Studies showing evidence of blood bacterial signatures in cardiometabolic disease in human

sumptions of infection source and the fact that culture-based methods are slow and fail to identify potential pathogens in 40% of cases [67], this application seems particularly central in subjects with sepsis [68] and high-risk subjects such as immunocompromised patients and those after hematopoietic cell transplantation (HCT) [69] or subjects with liver cirrhosis [52], as well as patients with diabetes. In the latter, the higher risk of primary bloodstream infections may arise from the translocation of organisms from damaged mucosal sites like the oral cavity or the gut. Additionally, it might be helpful in patients who are already treated with antibiotics, and in whome blood cultures usually fail. A promising approach was published by Tamburini et al., who developed a bioinformatic tool, which utilizes the potential of whole-genome sequencing to profile strain variations between metagenomes to match bloodstream pathogens in HCT to a candidate source. They found concomitant dominant gut colonization with known enteric pathogens a few days prior to bloodstream infection becoming clinically apparent. Interestingly, they found cases where typically non-enteric pathogens (e. g., *Pseudomonas aeruginosa* and *Staphylococcus aureus*) were present in the gut microbiota before infection and as-

certained functional relatedness by showing concordant predicted and clinical antibiotic resistance [70]. While turnaround time in sample preparation and sequencing and the high cost remain hindersome for efficient integration of whole-genome sequencing in clinical care, strain-level identification of pathogens and their reservoirs has the potential to improve infection prevention by reconstructing the trajectories and timing of colonization, the evolution of pathogenicity, and microbial adaptation [71]. This can further enhance management procedures by rapidly identifying resistance and escape mechanisms in real-time [63]. Considering this manuscript was written amid the coronavirus 2 (SARS-CoV-2) pandemic, it would almost be tone-deaf not to stress how metagenomics has been pivotal in helping us develop a holistic ecological understanding of microbial evolution over time and geography. This is particularly true for detecting subclonal mutations (used routinely in cancer research), which have helped delineate the chains of infection and interspecies transmission and the higher frequency of which has been associated with severe acute respiratory distress syndrome in SARS-CoV-2 [72]. The same holds for bacterial infection: bacterial DNA in the bloodstream more effectively identifies subjects at higher risk of death or more severe illness from suspected sepsis [73], regardless of the isolation of viable bacteria via blood culture alone [37]. While immunotherapy is promising but practically nonexistent in cardiometabolic disease [74], the implementation of screening for blood bacterial DNA might be beneficial in critically ill subjects, where adjunctive immunotherapy [75] or treatment with specific toll-like-receptor (TLR) antagonists is an option [76].

Bacteria in the blood: liquid biopsies of the new era?

As liver cirrhosis has been associated with a circulatory bacterial signature, it is not surprising that hepatocellular carcinoma (HCC) displays a specific alteration of the bacterial sequences found in the blood of diseased subjects. Specifically, the blood of subjects with HCC had a lower bacterial diversity. Moreover, significant differences in the relative abundance of Pseudomonas, Streptococcus, Bifidobacterium, Staphylococcus, Acinetobacter, Klebsiella, and Trabulsiella were noted. The latter four bacteria were heavily enriched in HCC, and Staphylococcus had the most significant association with HCC (p = 4.0e-08), showing a 4.3-fold increase compared to controls. The authors then identified a marker-based model depending on five bacterial genera, distinguishing HCC from controls (AUC of 0.879, accuracy 81.6%). Validation in a subgroup of the cohort confirmed that the model accurately differentiated HCC with an AUC of 0.875 and an accuracy of 79.8% [77]. Substantial contributions of local tissue and gut microbiome have been demonstrated for several cancer types [78-85], which further motivates the search for microorganism-derived molecules to diagnose noncommunicable diseases such as cancers. In a recent publication, Poore et al. reassessed whole-genome and whole-transcriptome sequencing studies in The Cancer Genome Atlas accounting for over 30 types of cancer from 104,814 treatment-naive subjects, detecting cancer-specific microbial signatures (viral and bacterial) in the tissues and blood, and benchmarking signatures from microbial DNA from the plasma against cell-free tumor DNA. They, moreover, used deep metagenomic sequencing on plasma samples from 100 tumor patients vs. 69 healthy subjects and showed that cell-free

microbial profiles were discriminatory between healthy and diseased subjects and between different types and stages of cancers. This was only possible for specific cancer types (colon, stomach adenocarcinomas and renal clear cell carcinoma), and the microbial signatures failed to differentiate intermediate cancer stages, suggesting that the microbial community structure might not be associated closely with cancer stages. They further trained a microbial source tracking algorithm and showed that the gut microbiome was most relevant for the bacterial signature in cancers of the gastrointestinal tract, further indicating that Fusobacterium spp. was overabundant in GI- vs. non-GI-tumors. That being said, the authors did not replicate the bacterial signatures found in HCC in other studies and relied on tissues more than blood samples, which were very often used as negative controls and did not generally show a differential microbial signature [24]. While bacterial signatures have been associated with cardiometabolic disease and cancer, this has been done in heterogeneous populations, and studies differed extensively in how they processed samples and what analytical tools were used, making it impossible to benchmark a 'meta-population blood bacterial signature' for cancer or cardiovascular disease. Currently, 'good old' circulatory tumor markers and imaging studies will need to suffice. It is desirable to standardize studies on blood bacterial signatures in several diseases and pair them with traditional disease assessment to apply them in the clinical context of prevention in the future.

Blood bacterial signatures derived therapeutic avenues

Targeting bacterial signatures in disease requires an understanding of how these signatures come about in the first place, whether they are a mirror of a common immunological denominator contributing both to metabolic health and increased susceptibility towards bacterial translocation, or whether translocated bacteria (alive or dead) are at the base of immunological cascades contributing to metabolic disease. Moreover, even if bacteria in the blood do not make up an ecological niche, bacterial functionality related to host-bacterial co-metabolism such as cholesterol degradation might still reflect the severity of cardiometabolic diseases. In this sense, consolidating a new therapeutic strategy might not directly require targeting bacteria in the blood but other upstream mechanisms associated with an increased bacterial translocation. Downstream strategies may ensue over time but are currently inaccessible due to the lack of informative mechanisms linking circulating microbial signatures to disease.

Examples for upstream strategies could be targeted at correcting altered gut microbiota via pro-, pre-, syn- or antibiotics, an impaired gut barrier, or the modulation of the crosstalk between the host and the microbiome by harnessing bacterial metabolites but also by targeting the bacterial metabolism of xenobiotics.

Elimination strategies such as via antibiotics

When it comes to antibiotics, studies in humans have been contradictory, showing that children treated with antibiotics are more likely to develop obesity and diabetes in a dose-dependent manner [86–88], while in mice at least, antibiotics treatment-induced improved gut barrier, insulin sensitivity and weight loss [89]. Furthermore, gram-positive bacteria-targeted antibiotic treatment of mice with systemic lupus erythematosus reduced bacterial growth in mesenteric lymph nodes and the liver [90].

From a clinical perspective, antibiotic treatment not seldom leads to a selection of pathogens such as *C. difficile*. Considering this, the lack of possibility to target only one specific strain/taxon or bacterial function, the increase of multiresistant pathogens, and lack of informative mechanistic studies, we believe the use of antibiotics to treat metabolic disease is currently questionable. Other strategies such as bacteriophage implementation, growth inhibition via bacteriostatic non-antibiotics, or even blood UV-Irradiation [91] might be alternatives but necessitate very nuanced testing for safety and efficacy. Moreover, these rather experimental approaches are hard to justify in metabolic disease, where pharmacotherapies with clinical evidence for hard outcomes are available.

Supplementation strategies via replenishing missing bugs

Fecal microbiota transplants (FMTs) FMTs has proven to be a valuable tool for the discovery of next-generation microbiota targeted therapeutics and has now been successfully implemented in the effective treatment of recurrent C. difficile either via colonic or duodenal infusion or in the form of oral frozen capsules containing fecal material from healthy donors [92, 93]. FMT, on the other hand, appears to only transiently improve insulin sensitivity and insulin secretion without ameliorating obesogenic phenotypes due to the failure of donor's microbiota to colonize the gut of the recipient's long-term [94, 95]. Interestingly, the application of FMT in newly diagnosed type 1 diabetes stabilized residual Beta-cell function mainly seen after autologous as compared to healthy donor FMT [96]. There are no studies proving a reduced translocation of bacterial fragments through the intestinal barrier after FMT. Alas, a case of bacteremia with a multiresistant bacteria with subsequent death has been recently described [97], encouraging the search for more targeted and safe strategies than FMTs in metabolic disease.

Supplementation of single bacteria to alleviate metabolic disease

Given the medical risk of FMTs and their short-lived success, there has been increasing focus on the therapeutical use of single taxa or consortia. Akkermansia muciniphila has received much attention, proving to alleviate obesity, insulin resistance, and increased gut permeability in mice [98]. Supplementation with pasteurized rather than live A. muciniphila mildly improved insulin sensitivity [99] (phase 1-2 Study) and improved gut barrier function via the thermostable outer-membrane protein Amuc_1100's interaction with TLR2 in the host [100]. Similarly, evidence of improved insulin sensitivity being associated with Eubacterium hallii (Anaerobutyricum) [101] after FMT and in mice [102] has led to a pilot study to investigate the effects of direct supplementation of E. hallii in humans, which implicated the supplementation in improving peripheral insulin sensitivity [103]. Similarly, Faecalibacterium prausnitzii is relatively increased in metabolic health [14, 104], which led to further studies to delineate potential beneficial effects on metabolism. While a microbial anti-inflammatory molecule derived from F. prausnitzii reduced inflammation in mice models of colitis via nuclear factor-kappaB inhibition [105], there is, to date, no direct evidence for an anti-hyperglycemic effect. Several bacteria could be considered in the quest for precision medicine in metabolic disease [106] but currently do not embed the notion of bacterial translocation as a clearly defined outcome.

Bacterial metabolites and postbiotics

Several bacterial metabolites have been shown to contribute to metabolic regulation in the host. Secondary bile acids, produced after hydrolysis of primary bile acids by the bacteria in the gut, seem to increase body expenditure and tissue-specific glucose [107–109] uptake as well as pancreas secretion of insulin [110]. But these effects have not been associated with bacterial translocation and are hard to track intracellularly considering the target farnesoid X receptor can be both activated and inhibited in different tissues with similar effects on the metabolism, suggesting further exploration of tissue-specific receptor pathways [109, 111]. On the other hand, short-chain-fatty acids are well-known microbial metabolites with pleiotropic effects on the host produced by bacterial fermentation of fibers [112]. Importantly, butyrate, observed to be ubiquity reduced in T2D (along with a decrease in butyrogenic groups [14, 113]), can directly reduce colonic inflammation by regulating interleukin-18 secretion in the epithelium and immunosuppressive T-cells [114] and hence maintain intestinal homeostasis and integrity via inhibition of histone deacetylase activity [114]. Several other metabolites have been clearly and robustly associated with metabolic diseases such as imidazole propionate in diabetes [115], trimethylamine N-oxide and tyrosine derived metabolites with cardiometabolic disease [116, 117] but also tryptophanderived microbial metabolites, which improve inflammation and the gut barrier function [118, 119] as well as microbially produced polyphenols such as urolithin A, which have been shown to reduce increased gut permeability [120] and improve mitochondrial health in humans [121], motivating the identification and application of postbiotics (bacteria-derived non-living components) or other therapeutic strategies targeting deleterious bacterial metabolites.

In summary, there is a plethora of microbially based next-generation therapeutics which need to be further tested in humans in the specific condition of disease, where their actions could indeed be scrutinized. In which case, it would be helpful to analyze whether these treatments modify bacterial translocation and bacterial blood signatures while changing the course or severity of the metabolic disease. This will further substantiate the contribution of these signatures to disease and place them in a more strategic context for diagnostic or therapeutic implementation.

Important considerations: Contamination, biases in assessment and mitigation strategies

While the technological and analytical developments in the microbiome field can be seen as overall monumental for the scientific landscape, inspiring curiosity, leading the way into personalized medicine of both communicable and non-communicable diseases [121, 122], and helping relinquish established dogmas, low bacterial biomass samples such as the blood require careful considerations of pitfalls and measures to mitigate for several biases [121– 123]. Specifically, these samples are highly prone to contamination from background environmental DNA and to technical biases, not the least of them being over-amplification during PCR [122–124]. The lack of standard operating procedures, benchmarked techniques, and best practices both on the experimental and analytical sides of research in working with low biomass samples has led to sensationalized science and wide controversy [125]. Hence, with increasing bacterial signature mining in low biomass samples, it is vital to unify experimental and analytical procedures aiming to overcome limiting biases, which should be fully understood and accounted for during study design, data interpretation but more importantly, during the development of diagnostic and therapeutic tools based on these results.

Technical aspects

Routinely, microbiome studies will usually consist of the following procedures before any analytical pipeline can be implemented: sample collection, DNA extraction, and sequencing library preparation (possibly with or without bacterial enrichment techniques such as culture [39, 126, 127] or bacterial DNA enrichment), highthroughput sequencing technologies [128] such as amplicon-based sequencing (e.g., 16 S rRNA gene), shotgun metagenomic sequencing [129], as well as RNA sequencing-based approaches [38]. Because procedural controls, which take contamination of samples during collection and cross-contamination during processing and sequencing into account, are rarely implemented, studies in low microbial biomass samples relying on sensitive metagenomic techniques highly confound and inflate the diagnostic implications for microbial contribution to disease [130]. Therefore, it is pivotal to identify the sources of contamination during sampling and laboratory procedures. These are 1) personal (doctors, nurses, doctor assistants, and technicians) and their direct 'personal cloud' [131], 2) the immediate environment of sampling, storage, and processing, and 3) laboratory reagents and equipment.

Accordingly, they can be mitigated: 1) contamination from a sampling/processing person could be reduced by wearing sterile masks, gloves, surgical headdress, surgical robes (for single use) and relying on automation where possible, which can be later better accounted for by correction for batch effects [132]. 2) Environmental contaminants are hard to control and account for, primarily because the sampling usually occurs in a different place from where the samples are stored and processed, and the environmental signatures of these locations can very well depend on the subjects working there [133]. Because these contaminants are most likely to be found in the air and on surfaces, it might be worthwhile to sample these environments during subject inclusion and processing times. 3) low biomass samples are most amenable for contamination during processing with lab reagents and equipment, which might still contain bacterial DNA even when they are 'bacteria free' with even the most promising kits for DNA extraction containing clear DNA signatures [124] and can overwhelm genuine bacterial signatures from the sampled tissue [134]. More interestingly and maybe less intuitively, low input DNA concentration can highly bias population levels because it results in overamplification of the template DNA, increases duplication rates, and favors AT-rich sequences [135], which does not, for example, occur in negative controls, leading hence to the overrepresentation of a particular species or even a contaminant and confounding populationlevel analyses. This requires a multi-pronged approach whereby a) reagents and equipment are subjected to UV radiation, b) controls (extraction blanks) from the sampling and laboratory environments, equipment and reagents are carried through the entire procedures as the actual samples, samples are randomized, and standard operating procedures are consistently followed during lab work with the inclusion of no-template amplification (possibly by the same subjects for a specific project) and c) bioinformatics approaches are included to assess contamination [136, 137], track its source [138, 139], account for it, and remove it [139]. Similarly, reviewers, editors, and readers should be versed when looking at studies analyzing the impact of bacterial signatures in disease. For the interested experimenter and reader, we wholeheartedly recommend reviewing the RIDE-criteria and the recent publication by Eisenhofer et al. [130, 140].

Scientific aspects

While this area of research has immense untapped potential, it is vital to avoid methodological pitfalls, which would lead to an inevitable waste of time, money and highly tarnish the credibility of the research. Moreover, it would be important to address limitations in the design and interpretation of current research. For example, while comparing studies, it is essential to understand the effect of the matrix from which the bacterial signature is derived: is it plasma, whole blood, or peripheral blood mononuclear cells? The distribution of microbial DNA between compartments of the blood is not clear. This information is important because the matrix can highly influence technical procedures such as PCR efficiency. This has overarching implications pertaining to the source and the physiological role of the bacteria/bacterial sequences. Similarly, studies, which controlled and corrected for contamination, did so very differently, making the comparison between them non-conclusive. It will therefore be important to establish an optimal method to control for contaminants that goes beyond complete subtraction of flagged taxa. Specifically, a prevalent risk with stringent decontamination is that real signals reflecting commensal, tissue-specific microbial communities and concomitant predictive microbial profiles may be discarded. Poore et al. showed, for example, that a stringent filtering approach might be desirable with the possible downside, if scalable/universal, to preclude biologically relevant and informative results [24].

It is also safe to say that most works in this area currently fail to address causality and rely on correlation. While correlational observations are an important steppingstone and have often been the beginnings of great stories, they remain exactly that: the beginning. Hence, it is important to address outstanding questions to inform future research and applications. Some of these questions pertain to a) whether sequences are indeed cell-free bacterial DNA or come from bacteria that are either dead, alive, or dormant, b) whether bacteria in the blood occupy a real ecological niche, c) the source of the bacterial DNA or bacteria for which examining primary tissue specimens along matched gut epithelia and sampled orifices will be pivotal, and d) the tissue selectivity of bacteria, which could potentially be informative.

While microbial DNA has the potential of inherent pathogenicity because it activates TLR9 in its unmethylated form resulting in inflammatory cascades [141, 142], the critical question to answer is whether microbial DNA in itself is a pathogenic finding, whether it is an epiphenomenon that reflects overall or specific disease burden or whether both metabolic disease and bacterial signatures are epiphenomena of lurking immunological conditions or active infections (**Fig. 3**), which we yet have to be identified because our tests lack sensitivity.

In moving towards mechanisms and granular interrogation of causality, it will also be crucial to understand whether the entire bacteria, bacterial cell components, or metabolites are most important for the postulated effects of the identified microorganisms. A question that lends itself to further exploration concomitantly is whether the observed taxa are alive or not. A drawback of cultureindependent sequencing methods is their inability to differentiate living from dead bacteria. This is particularly important because DNA, including that from contaminants, can persist in the environment. More importantly, while DNA sequencing and functional annotation can predict bacterial metabolism, only viable bacteria can undertake active metabolic processes. Therefore, methods that do not relay viability information will always overestimate the importance of bacterial metabolism in the host metabolic response, as long as they are not coupled with information on transcription of metabolic modules or direct measurement of metabolites.

Hence, the application of methods that interrogate active metabolism and delineate viability is warranted. Techniques exploring viability can characterize one or several viability aspects, such as the presence of an intact membrane, the replication of genetic material (via RNA detection), and the detection of metabolism or energy. The main challenge being, whether these methods are compatible with next-generation sequencing and whether they are applicable to low-biomass samples.



► Fig. 3 Bacterial translocation can be derived from several sources such as the skin, the oral cavity, the gut, clinical infection sources, medical procedures including dwelling catheters, and contamination/infection from a hospital stay. The translocation can include endotoxins, whole bacteria, or bacterial sequences leading to systemic and local tissue inflammation and culminating in local tissue dysfunction such as insulin resistance in adipose tissue, liver dysfunction, or cardiovascular disease. Similarly, signaling metabolites from the gut can contribute to cardiometabolic disease and specific tissue dysfunctions, which in turn can aggravate bacterial translocation by increasing susceptibility to colonization and infection and increased 'leakiness of the gut.'

Microbial culture caters to both these needs and is the gold standard to prove viability because it displays the ability of bacterial cells to divide and metabolize nutrients. On the other hand, several bacterial strains are painstakingly hard to culture or require immense optimization to determine optimal culturing conditions. Others are viable but non-culturable because they are either damaged or maintaining the structural and metabolic properties observed or postulated in their environment of origin is not possible [143]. The application of propidium iodine, propidium monoazide (PMA), and ethidium monoazide - all dyes that can bind to DNA in membrane compromised cells and extracellular DNA- enable the labeling of dead bacteria and can be coupled with several other techniques, including epifluorescence microscopy, flow cytometry, but also gPCR, digital PCR and, metagenomics among several other downstream techniques [144]. PMA seems to be more selective and less cytotoxic than other methods, and several of its selectivity drawbacks are known (stains viable cells for some species while failing to stain dead cells from other species). More importantly, it can be used for low-biomass samples, assists in the removal of contaminant DNA (environmental or introduced by the reagents including PCR reagents) [124], has the potential to enrich for rare microbial community members [124, 145], and allows for gPCR guantification to distinguish the viable from the total fraction of microbial cells in several settings [146] (reviewed in [144]). Unfortunately, the samples are required to be in an aqueous solution, and ethylenediaminetetraacetic acid (EDTA, found in specific blood collection tubes) can affect dye permeability through membranes [147], limiting the application of this method to other human low biomass samples such as the blood. Other methods such as RNA analyses (e.g., metatranscriptomics) enable quantification and identification of potentially 'active bacterial community members' and can delineate taxa as well as acute responses to stimuli relatively early on because RNA has a very short half-life [148]. Unfortunately, RNA yield is highly dependent on the matrix it derives from and the preservation method employed until RNA extraction is undertaken [148, 149]. Moreover, significant RNA losses have been described during sample preparation, which is particularly problematic in low-biomass samples, where yield is low to begin with [150, 151]. Similarly, metaproteomics can also be applied for the detection of active bacteria. Coupled with metagenomics, it can enable the identification of active metabolic pathways of bacterial communities and connect them to substrate availability and environmental conditions. A drawback is that only proteins with exact matches in the search databases can be identified. Another drawback is the potential persistence of proteins in specific environments.

Various techniques are available and allow their application in conjunction with several other methods to detect viability, phylogeny, and metabolic activity, lending themselves well to studies assessing the contribution of bacterial cells to overall host health. Several bottlenecks remain, most prominently the necessity for extensive optimization in a setting where human tissue is a matrix, the low throughput of these methods paralleled with the high-cost and the need for careful result interpretation, making this a challenging task but an excellent opportunity for collaboration between groups across the scientific and medical spectrum.

Conclusions

Physicians are familiar with cases where viruses induce tumors (such as. Kaposi-sarcoma) or associate with metabolic disease (such as T1D) and are clinically versed in bacteria-induced rheumatic complications (such as heart rheumatism and Reiter syndrome). More importantly, comprehensive contamination-aware analytical approaches employed in large-scale studies have established a robust link between bacterial translocation to human body compartments such as the blood or adipose tissue and cardiometabolic diseases. This prior knowledge and emerging and converging evidence make it difficult to categorically deny the presence of bacteria and bacterial DNA in the blood and the potential of this genetic material (whether alive or dead) to act as an immunostimulatory factor thus triggering or influencing the course of non-communicable cardiometabolic diseases.

A word of caution is still warranted: while the existence of a tissue bacterial signature for the blood seems likely, considering the presence of comparable taxa in independent studies even in health [26], we believe the presence of a stable, non-pathogenic, highly complex microbial community such as the one we see in the gut rather unlikely, because it is neither consistently detectable [152], nor is it highly prevalent [21–23, 125], nor logical based on ecological principles.

Considering recent advances in analytical and bioinformatics tools coupled with observations from clinical studies within intercontinental research consortia, the emerging area of tissue microbiome research is still in its infancy but promises novel prognostic, diagnostic, and therapeutic strategies with paradigm-shifting potential for health care and disease prevention and treatment. To be able to move into translational avenues, our efforts will need to move beyond pure associations towards functional approaches to understand causative chains better, i. e., define causes and consequences in the highly complex interplay between blood bacterial signatures and metabolic diseases.

Conflict of Interest

The authors declare that they have no competing interests.

References

- Moeller AH, Caro-Quintero A, Mjungu D et al. Cospeciation of gut microbiota with hominids. Science 2016; 353: 380–382
- [2] Garn H, Potaczek DP, Pfefferle PI. The hygiene hypothesis and new perspectives-current challenges meeting an old postulate. Front Immunol 2021; 12: 637087
- [3] Sommer F, Bäckhed F. The gut microbiota-masters of host development and physiology. Nat Rev Microbiol 2013; 11: 227–238. doi:10.1038/nrmicro2974
- [4] Haiser HJ, Gootenberg DB, Chatman K et al. Predicting and manipulating cardiac drug inactivation by the human gut bacterium Eggerthella lenta. Science 2013; 341: 295–298
- [5] Alexander JL, Wilson ID, Teare J et al. Gut microbiota modulation of chemotherapy efficacy and toxicity. Nat Rev Gastroenterol Hepatol 2017; 14: 356–365

- [6] McQuade JL, Daniel CR, Helmink BA et al. Modulating the microbiome to improve therapeutic response in cancer. Lancet Oncol 2019; 20: e77–e91
- [7] Pedersen HK, Gudmundsdottir V, Nielsen HB et al. Human gut microbes impact host serum metabolome and insulin sensitivity. Nature 2016; 535: 376–381
- [8] Turnbaugh PJ, Ley RE, Hamady M et al. The human microbiome project. Nature 2007; 449: 804–810. Stand: 30.09.2021
- [9] Vieira-Silva S, Falony G, Belda E et al. Statin therapy is associated with lower prevalence of gut microbiota dysbiosis. Nature 2020; 581: 310–315
- [10] Molinaro A, Bel Lassen P, Henricsson M et al. Imidazole propionate is increased in diabetes and associated with dietary patterns and altered microbial ecology. Nat Commun 2020; 11: 5881
- [11] Falony G, Joossens M, Vieira-Silva S et al. Population-level analysis of gut microbiome variation. Science 2016; 352: 560–564
- [12] Forslund K, Hildebrand F, Nielsen T et al. Disentangling type 2 diabetes and metformin treatment signatures in the human gut microbiota. Nature 2015; 528: 262–266
- [13] Vich Vila A, Collij V, Sanna S et al. Impact of commonly used drugs on the composition and metabolic function of the gut microbiota. Nat Commun 2020; 11: 362
- [14] Qin J, Li Y, Cai Z et al. A metagenome-wide association study of gut microbiota in type 2 diabetes. Nature 2012; 490: 55–60
- [15] Fan Y, Pedersen O. Gut microbiota in human metabolic health and disease. Nat Rev Microbiol 2021; 19: 55–71. doi:10.1038/s41579-020-0433-9
- [16] Lockhart PB, Brennan MT, Sasser HC et al. Bacteremia associated with toothbrushing and dental extraction. Circulation 2008; 117: 3118–3125
- [17] Mougeot FKB, Saunders SE, Brennan MT et al. Associations between bacteremia from oral sources and distant-site infections: Tooth brushing versus single tooth extraction. Oral Surg Oral Med Oral Pathol Oral Radiol 2015; 119: 430–435
- [18] Hotamisligil GS. Inflammation, metaflammation and immunometabolic disorders. Nature 2017; 542: 177–185
- [19] Cani PD, Amar J, Iglesias MA et al. Metabolic endotoxemia initiates obesity and insulin resistance. Diabetes 2007; 56: 1761–1772
- [20] Agwunobi AO, Reid C, Maycock P et al. Insulin resistance and substrate utilization in human endotoxemia. J Clin Endocrinol Metab 2000; 85: 3770–3778
- [21] Massier L, Chakaroun R, Tabei S et al. Adipose tissue derived bacteria are associated with inflammation in obesity and type 2 diabetes. Gut 2020; 69: 1796–1806
- [22] Chakaroun RM, Massier L, Heintz-Buschart A et al. Circulating bacterial signature is linked to metabolic disease and shifts with metabolic alleviation after bariatric surgery. Genome Med 2021; 13: 105
- [23] Anhê FF, Jensen BAH, Varin TV et al. Type 2 diabetes influences bacterial tissue compartmentalisation in human obesity. Nat Metab 2020; 2: 233–242. Stand: 30.09.2021
- [24] Poore GD, Kopylova E, Zhu Q et al. Microbiome analyses of blood and tissues suggest cancer diagnostic approach. Nature 2020; 579: 567–574
- [25] Tedeschi GG, Amici D, Paparelli M. Incorporation of nucleosides and amino-acids in human erythrocyte suspensions: possible relation with a diffuse infection of mycoplasms or bacteria in the L form. Nature 1969; 222: 1285–1286
- [26] Castillo DJ, Rifkin RF, Cowan DA et al. The healthy human blood microbiome: Fact or fiction? Front Cell Infect Microbiol 2019; 9: 148

- [27] Moriyama K, Ando C, Tashiro K et al. Polymerase chain reaction detection of bacterial 16S rRNA gene in human blood. Microbiol Immunol 2008; 52: 375–382. doi:10.1111/j.1348-0421.2008.00048.x
- [28] Dunne WM Jr, Burnham C-AD. eds. Dark art of blood cultures. John Wiley & Sons; 2017
- [29] Dinakaran V, Rathinavel A, Pushpanathan M et al. Elevated levels of circulating DNA in cardiovascular disease patients: Metagenomic profiling of microbiome in the circulation. PLoS ONE 2014; 9: e105221
- [30] Amar J, Serino M, Lange C et al. Involvement of tissue bacteria in the onset of diabetes in humans: Evidence for a concept. Diabetologia 2011; 54: 3055–3061
- [31] Sato J, Kanazawa A, Ikeda F et al. Gut dysbiosis and detection of "live gut bacteria" in blood of Japanese patients with type 2 diabetes. Diabetes Care 2014; 37: 2343–2350
- [32] Damgaard C, Magnussen K, Enevold C et al. Viable bacteria associated with red blood cells and plasma in freshly drawn blood donations. PLoS ONE 2015; 10: e0120826. doi:10.1371/journal. pone.0120826
- [33] Traykova D, Schneider B, Chojkier M et al. Blood microbiome quantity and the hyperdynamic circulation in decompensated cirrhotic patients. PLoS ONE 2017; 12: e0169310
- [34] Li Q, Wang C, Tang C et al. Identification and characterization of blood and neutrophil-associated microbiomes in patients with severe acute pncreatitis using next-generation sequencing. Front Cell Infect Microbiol 2018; 8: 5
- [35] Nikkari S, McLaughlin JJ, Bi W et al. Does blood of healthy subjects contain bacterial ribosomal DNA? J Clin Microbiol 2001; 39: 1956–1959
- [36] McLaughlin RW, Vali H, Lau PCK et al. Are there naturally occurring pleomorphic bacteria in the blood of healthy humans? J Clin Microbiol 2002; 40: 4771–4775
- [37] O'Dwyer MJ, Starczewska MH, Schrenzel J et al. The detection of microbial DNA but not cultured bacteria is associated with increased mortality in patients with suspected sepsis – a prospective multi-centre European observational study. Clin Microbiol Infect 2017; 23: 208.e1–e208.e6. doi:10.1016/j.cmi.2016.11.010
- [38] Whittle E, Leonard MO, Harrison R et al. Multi-method characterization of the human circulating microbiome. Front Microbiol 2018; 9: 3266
- [39] Panaiotov S, Hodzhev Y, Tsafarova B et al. Culturable and nonculturable blood microbiota of healthy individuals. Microorganisms 2021; 9:. doi:10.3390/microorganisms9071464
- [40] Martel J, Wu C-Y, Huang P-R et al. Pleomorphic bacteria-like structures in human blood represent non-living membrane vesicles and protein particles. Sci Rep 2017; 7: 10650
- [41] Olde Loohuis LM, Mangul S, Ori APS et al. Transcriptome analysis in whole blood reveals increased microbial diversity in schizophrenia. Transl Psychiatry 2018; 8: 96
- [42] Potgieter M, Bester J, Kell DB et al. The dormant blood microbiome in chronic, inflammatory diseases. FEMS Microbiol Rev 2015; 39: 567–591
- [43] Chong J, Liu P, Zhou G et al. Using MicrobiomeAnalyst for comprehensive statistical, functional, and meta-analysis of microbiome data. Nat Protoc 2020; 15: 799–821
- [44] Bolyen E, Rideout JR, Dillon MR et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol 2019; 37: 852–857
- [45] Leibovici L, Cohen O, Wysenbeek AJ. Occult bacterial infection in adults with unexplained fever. Validation of a diagnostic index. Arch Intern Med 1990; 150: 1270–1272

- [46] Leibovici L, Samra Z, Konisberger H et al. Bacteremia in adult diabetic patients. Diabetes Care 1991; 14: 89–94
- [47] Amar J, Lange C, Payros G et al. Blood microbiota dysbiosis is associated with the onset of cardiovascular events in a large general population: The D.E.S.I.R. study. PLoS ONE 2013; 8: e54461. doi:10.1371/journal.pone.0054461
- [48] Qiu J, Zhou H, Jing Y et al. Association between blood microbiome and type 2 diabetes mellitus: A nested case-control study. J Clin Lab Anal 2019; 33: e22842
- [49] Ortiz S, Zapater P, Estrada JL et al. Bacterial DNA translocation holds increased insulin resistance and systemic inflammatory levels in morbid obese patients. J Clin Endocrinol Metab 2014; 99: 2575–2583
- [50] Simon TG, Roelstraete B, Hagström H et al. Non-alcoholic fatty liver disease and incident major adverse cardiovascular events: Results from a nationwide histology cohort. Gut 2021. doi:10.1136/ gutjnl-2021-325724
- [51] Schierwagen R, Alvarez-Silva C, Madsen MSA et al. Circulating microbiome in blood of different circulatory compartments. Gut 2019; 68: 578–580
- [52] Usui S, Ebinuma H, Chu P-S et al. Detection of bacterial DNA by in situ hybridization in patients with decompensated liver cirrhosis. BMC Gastroenterol 2017; 17: 106
- [53] Mok Y, Ballew SH, Matsushita K. Chronic kidney disease measures for cardiovascular risk prediction. Atherosclerosis 2021. doi:10.1016/j. atherosclerosis.2021.09.007
- [54] Hobby GP, Karaduta O, Dusio GF et al. Translational physiology: Chronic kidney disease and the gut microbiome. Am J Physiol – Renal Physiol 2019; 316: F1211. Stand: 01.10.2021
- [55] Vanholder R, Schepers E, Pletinck A et al. The uremic toxicity of indoxyl sulfate and p-cresyl sulfate: A systematic review. J Am Soc Nephrol 2014; 25: 1897–1907
- [56] Smith PM, Howitt MR, Panikov N et al. The microbial metabolites, short-chain fatty acids, regulate colonic Treg cell homeostasis. Science 2013; 341: 569–573
- [57] Wong J, Piceno YM, DeSantis TZ et al. Expansion of urease- and uricase-containing, indole- and p-cresol-forming and contraction of short-chain fatty acid-producing intestinal microbiota in ESRD. Am J Nephrol 2014; 39: 230–237
- [58] Shah NB, Allegretti AS, Nigwekar SU et al. Blood microbiome profile in CKD: A pilot study. Clin J Am Soc Nephrol 2019; 14: 692–701
- [59] Luo Z, Alekseyenko AV, Ogunrinde E et al. Rigorous plasma microbiome analysis method enables disease association discovery in clinic. Front Microbiol 2020; 11: 613268
- [60] Cowan LT, Lutsey PL, Pankow JS et al. Inpatient and outpatient infection as a trigger of cardiovascular disease: The ARIC study. J Am Heart Assoc 2018; 7: e009683
- [61] Zhou X, Li J, Guo J et al. Gut-dependent microbial translocation induces inflammation and cardiovascular events after ST-elevation myocardial infarction. Microbiome 2018; 6: 66
- [62] Amar J, Lelouvier B, Servant F et al. Blood microbiota modification after myocardial infarction depends upon low-density lipoprotein cholesterol levels. J Am Heart Assoc 2019; 8: e011797
- [63] Taxt AM, Avershina E, Frye SA et al. Rapid identification of pathogens, antibiotic resistance genes and plasmids in blood cultures by nanopore sequencing. Sci Rep 2020; 10:. doi:10.1038/s41598-020-64616-x
- [64] Ray C, Ming X. Climate change and human health: A review of allergies, autoimmunity and the microbiome. Int J Environ Res Public Health 2020; 17: 4814. doi:10.3390/ijerph17134814
- [65] Heintz-Buschart A, Wilmes P. Human gut microbiome: Function matters. Trends Microbiol 2018; 26: 563–574

- [66] Snitkin ES, Zelazny AM, Montero CI et al. Genome-wide recombination drives diversification of epidemic strains of Acinetobacter baumannii. Proc Natl Acad Sci U S A 2011; 108: 13758–13763
- [67] Phua J, Ngerng W, See K et al. Characteristics and outcomes of culture-negative versus culture-positive severe sepsis. Crit Care 2013; 17: R202
- [68] Dellinger RP, Levy MM, Rhodes A et al. Surviving Sepsis Campaign: International guidelines for management of severe sepsis and septic shock, 2012. Intensive Care Med 2013; 39: 165–228
- [69] Blennow O, Ljungman P, Sparrelid E et al. Incidence, risk factors, and outcome of bloodstream infections during the pre-engraftment phase in 521 allogeneic hematopoietic stem cell transplantations. Transpl Infect Dis 2014; 16: 106–114
- [70] Tamburini FB, Andermann TM, Tkachenko E et al. Precision identification of diverse bloodstream pathogens in the gut microbiome. Nat Med 2018; 24: 1809–1814
- [71] Dingemans J, Ye L, Hildebrand F et al. The deletion of TonBdependent receptor genes is part of the genome reduction process that occurs during adaptation of Pseudomonas aeruginosa to the cystic fibrosis lung. Pathog Dis 2014; 71: 26–38
- [72] Lythgoe KA, Hall M, Ferretti L et al. SARS-CoV-2 within-host diversity and transmission. Science 2021; 372:. doi:10.1126/science.abg0821
- [73] Bloos F, Hinder F, Becker K et al. A multicenter trial to compare blood culture with polymerase chain reaction in severe human sepsis. Intensive Care Med 2010; 36: 241–247
- [74] Esser N, Paquot N, Scheen AJ. Anti-inflammatory agents to treat or prevent type 2 diabetes, metabolic syndrome and cardiovascular disease. Expert Opinion on Investigational Drugs 2015; 24: 283–307. doi:10.1517/13543784.2015.974804
- [75] Leentjens J, Kox M, van der Hoeven JG et al. Immunotherapy for the adjunctive treatment of sepsis: From immunosuppression to immunostimulation. Time for a paradigm change? Am J Respir Crit Care Med 2013; 187: 1287–1293
- [76] Savva A, Roger T. Targeting toll-like receptors: Promising therapeutic strategies for the management of sepsis-associated pathology and infectious diseases. Front Immunol 2013; 4: 387
- [77] Cho EJ, Leem S, Kim SA et al. Circulating microbiota-based metagenomic signature for detection of hepatocellular carcinoma. Sci Rep 2019; 9: 7536
- [78] Bullman S, Pedamallu CS, Sicinska E et al. Analysis of persistence and antibiotic response in colorectal cancer. Science 2017; 358: 1443–1448
- [79] Dejea CM, Fathi P, Craig JM et al. Patients with familial adenomatous polyposis harbor colonic biofilms containing tumorigenic bacteria. Science 2018; 359: 592–597
- [80] Geller LT, Barzily-Rokni M, Danino T et al. Potential role of intratumor bacteria in mediating tumor resistance to the chemotherapeutic drug gemcitabine. Science 2017; 357: 1156–1160
- [81] Jin C, Lagoudas GK, Zhao C et al. Commensal microbiota promote lung cancer development via $\gamma\delta$ T Cells. Cell 2019; 176: 998–1013. e16
- [82] Ma C, Han M, Heinrich B et al. Gut microbiome-mediated bile acid metabolism regulates liver cancer via NKT cells. Science 2018; 360:. doi:10.1126/science.aan5931
- [83] Matson V, Fessler J, Bao R et al. The commensal microbiome is associated with anti–PD-1 efficacy in metastatic melanoma patients. Science 2018; 359: 104–108. doi:10.1126/science.aao3290
- [84] Meisel M, Hinterleitner R, Pacis A et al. Microbial signals drive pre-leukaemic myeloproliferation in a Tet2-deficient host. Nature 2018; 557: 580–584. doi:10.1038/s41586-018-0125-z

- [85] Ye H, Adane B, Khan N et al. Subversion of systemic glucose metabolism as a mechanism to support the growth of leukemia cells. Cancer Cell 2018; 34: 659–673. e6
- [86] Mikkelsen KH, Knop FK, Frost M et al. Use of antibiotics and risk of type 2 diabetes: A population-based case-control study. J Clin Endocrinol Metab 2015; 100: 3633–3640
- [87] Scott FI, Horton DB, Mamtani R et al. Administration of antibiotics to children before age 2 years increases risk for childhood obesity. Gastroenterology 2016; 151: 120–129. e5
- [88] Wernroth M-L, Fall K, Svennblad B et al. Early childhood antibiotic treatment for otitis media and other respiratory tract infections is associated with risk of type 1 diabetes: A nationwide register-based study with sibling analysis. Diabetes Care 2020; 43: 991–999
- [89] Miao Z, Cheng R, Zhang Y et al. Antibiotics can cause weight loss by impairing gut microbiota in mice and the potent benefits of lactobacilli. Biosci Biotechnol Biochem 2020; 84: 411–420
- [90] Manfredo VS, Hiltensperger M, Kumar V et al. Translocation of a gut pathobiont drives autoimmunity in mice and humans. Science 2018; 359: 1156–1161
- [91] Hamblin MR. Ultraviolet irradiation of blood: "The cure that time forgot"? Adv Exp Med Biol 2017; 295–309. doi:10.1007/978-3-319-56017-5_25
- [92] Mattila E, Uusitalo-Seppälä R, Wuorela M et al. Fecal transplantation, through colonoscopy, is effective therapy for recurrent Clostridium difficile infection. Gastroenterology 2012; 142: 490–496. doi:10.1053/j.gastro.2011.11.037
- [93] Savidge T, Spinler J. Faculty opinions recommendation of oral, capsulized, frozen fecal microbiota transplantation for relapsing Clostridium difficile infection. Faculty Opinions 2015. doi:10.3410 /f.722182886.793503668
- [94] Zhang, Zhang, Mocanu et al. Impact of fecal microbiota transplantation on obesity and metabolic syndrome – A systematic review. Nutrients 2019; 11: 2291. doi:10.3390/nu11102291
- [95] de Groot P, de Groot P, Scheithauer T et al. Donor metabolic characteristics drive effects of faecal microbiota transplantation on recipient insulin sensitivity, energy expenditure and intestinal transit time. Gut 2020; 69: 502–512. doi:10.1136/gutjnl-2019-318320
- [96] de Groot P, Nikolic T, Pellegrini S et al. Faecal microbiota transplantation halts progression of human new-onset type 1 diabetes in a randomised controlled trial. Gut 2021; 70: 92–105
- [97] DeFilipp Z, Bloom PP, Soto MT et al. Drug-resistant E. coli bacteremia transmitted by fecal microbiota transplant. N Engl J Med 2019; 381: 2043–2050. doi:10.1056/nejmoa1910437
- [98] Everard A, Belzer C, Geurts L et al. Cross-talk between Akkermansia muciniphila and intestinal epithelium controls diet-induced obesity. Proc Natl Acad Sci U S A 2013; 110: 9066–9071
- [99] Depommier C, Everard A, Druart C et al. Supplementation with Akkermansia muciniphila in overweight and obese human volunteers: A proof-of-concept exploratory study. Nat Med 2019; 25: 1096–1103
- [100] Plovier H, Everard A, Druart C et al. A purified membrane protein from Akkermansia muciniphila or the pasteurized bacterium improves metabolism in obese and diabetic mice. Nat Med 2017; 23: 107–113
- [101] Vrieze A, Van Nood E, Holleman F et al. Transfer of intestinal microbiota from lean donors increases insulin sensitivity in individuals with metabolic syndrome. Gastroenterology 2012; 143: 913–916. e7
- [102] Udayappan S, Manneras-Holm L, Chaplin-Scott A et al. Oral treatment with improves insulin sensitivity in mice. NPJ Biofilms Microbiomes 2016; 2: 16009

- [103] Gilijamse PW, Hartstra AV, Levin E et al. Treatment with Anaerobutyricum soehngenii: A pilot study of safety and dose– response effects on glucose metabolism in human subjects with metabolic syndrome. NPJ Biofilms Microbiomes 2020; 6: 16. doi:10.1038/s41522-020-0127-0
- [104] Koh A, Bäckhed F. From association to causality: The role of the gut microbiota and its functional products on host metabolism. Mol Cell 2020; 78: 584–596
- [105] Quévrain E, Maubert MA, Michon C et al. Identification of an anti-inflammatory protein from Faecalibacterium prausnitzii, a commensal bacterium deficient in Crohn's disease. Gut 2016; 65: 415–425
- [106] Bui TPN, de Vos WM. Next-generation therapeutic bacteria for treatment of obesity, diabetes, and other endocrine diseases. Best Pract Res Clin Endocrinol Metab 2021; 35: 101504
- [107] Broeders EPM, Nascimento EBM, Havekes B et al. The bile acid chenodeoxycholic acid increases human brown adipose tissue activity. Cell Metab 2015; 22: 418–426
- [108] Kars M, Yang L, Gregor MF et al. Tauroursodeoxycholic acid may improve liver and muscle but not adipose tissue insulin sensitivity in obese men and women. Diabetes 2010; 59: 1899–1905. doi:10.2337/db10-0308
- [109] Sun L, Xie C, Wang G et al. Gut microbiota and intestinal FXR mediate the clinical benefits of metformin. Nat Med 2018; 24: 1919–1929. doi:10.1038/s41591-018-0222-4
- [111] Zhang H-M, Wang X, Wu Z-H et al. Beneficial effect of farnesoid X receptor activation on metabolism in a diabetic rat model. Mol Med Rep 2016; 13: 2135–2142. doi:10.3892/mmr.2016.4761
- [112] van der Hee B, Wells JM. Microbial regulation of host physiology by short-chain fatty acids. Trends Microbiol 2021; 29: 700–712
- [113] de Goffau MC, Luopajärvi K, Knip M et al. Fecal microbiota composition differs between children with β -cell autoimmunity and those without. Diabetes 2013; 62: 1238–1244
- [114] Koh A, De Vadder F, Kovatcheva-Datchary P et al. From dietary fiber to host physiology: Short-chain fatty acids as key bacterial metabolites. Cell 2016; 165: 1332–1345
- [115] Koh A, Molinaro A, Ståhlman M et al. Microbially produced imidazole propionate impairs insulin signaling through mTORC1. Cell 2018; 175: 947–961. e17
- [116] Tang WH, Kitai T, Hazen SL. Gut microbiota in cardiovascular health and disease. Circ Res 2017; 120:. doi:10.1161/CIRCRESAHA.117.309715
- [117] Lin CJ, Wu V, Wu PC et al. Meta-analysis of the associations of p-cresyl sulfate (PCS) and indoxyl sulfate (IS) with cardiovascular events and all-Cause mortality in patients with chronic renal failure. PLoS ONE 2015; 10:. doi:10.1371/journal.pone.0132589
- [118] Wlodarska M, Luo C, Kolde R et al. Indoleacrylic acid produced by commensal Peptostreptococcus species suppresses inflammation. Cell Host Microbe 2017; 22: 25–37. e6
- [119] Venkatesh M, Mukherjee S, Wang H et al. Symbiotic bacterial metabolites regulate gastrointestinal barrier function via the xenobiotic sensor PXR and toll-like receptor 4. Immunity 2014; 41: 296–310
- [120] Singh R, Chandrashekharappa S, Bodduluri SR et al. Enhancement of the gut barrier integrity by a microbial metabolite through the Nrf2 pathway. Nat Commun 2019; 10: 89
- [121] Andreux PA, Blanco-Bose W, Ryu D et al. The mitophagy activator urolithin A is safe and induces a molecular signature of improved mitochondrial and cellular health in humans. Nat Metab 2019; 1: 595–603

- [122] Petrosino JF. The microbiome in precision medicine: The way forward. Genome Med 2018; 10: 12
- [123] Selway CA, Eisenhofer R, Weyrich LS. Microbiome applications for pathology: Challenges of low microbial biomass samples during diagnostic testing. Hip Int 2020; 6: 97–106
- [124] Salter SJ, Cox MJ, Turek EM et al. Reagent and laboratory contamination can critically impact sequence-based microbiome analyses. BMC Biol 2014; 12: 87
- [125] Lauder AP, Roche AM, Sherrill-Mix S et al. Comparison of placenta samples with contamination controls does not provide evidence for a distinct placenta microbiota. Microbiome 2016; 4:. doi:10.1186/ s40168-016-0172-3
- [126] Hassan MM, Ranzoni A, Cooper MA. A nanoparticle-based method for culture-free bacterial DNA enrichment from whole blood. Biosens Bioelectron 2018; 99: 150–155
- [127] Weiß CL, Gansauge M-T, Aximu-Petri A et al. Mining ancient microbiomes using selective enrichment of damaged DNA molecules. BMC Genomics 2020; 21: 432
- [128] Kumar R, Eipers P, Little RB et al. Getting started with microbiome analysis: Sample acquisition to bioinformatics. Curr Protoc Hum Genet 2014; 82: 18.8.1–29
- [129] Gu W, Miller S, Chiu CY. Clinical metagenomic next-generation sequencing for pathogen detection. Annu Rev Pathol: Mechanisms of disease 2019; 14: 319–338. doi:10.1146/annurevpathmechdis-012418-012751
- [130] Eisenhofer R, Minich JJ, Marotz C et al. Contamination in low microbial biomass microbiome studies: Issues and recommendations. Trends Microbiol 2019; 27: 105–117
- [131] You R, Cui W, Chen C et al. Measuring the short-term emission rates of particles in the "Personal Cloud" with different clothes and activity intensities in a sealed chamber. Aerosol Air Qual Res 2013; 13: 911–921. doi:10.4209/aaqr.2012.03.0061
- [132] Minich JJ, Sanders JG, Amir A et al. Quantifying and understanding well-to-well contamination in microbiome research. mSystems 2019; 4:. doi:10.1128/mSystems.00186-19
- [133] Kelley ST, Gilbert JA. Studying the microbiology of the indoor environment. Genome Biol 2013; 14: 202
- [134] Tanner MA, Goebel BM, Dojka MA et al. Specific ribosomal DNA sequences from diverse environmental settings correlate with experimental contaminants. Appl Environ Microbiol 1998; 64: 3110–3113
- [135] Chafee M, Maignien L, Simmons SL. The effects of variable sample biomass on comparative metagenomics. Environ Microbiol 2015; 17: 2239–2253
- [136] Llamas B, Valverde G, Fehren-Schmitz L et al. From the field to the laboratory: Controlling DNA contamination in human ancient DNA research in the high-throughput sequencing era. STAR: Sci Technol Archaeol Res 2017; 3: 1–14. doi:10.1080/20548923.2016.1258824
- [137] Minich JJ, Zhu Q, Janssen S et al. KatharoSeq enables high-throughput microbiome analysis from low-biomass samples. mSystems 2018; 3:. doi:10.1128/mSystems.00218-17
- [138] Knights D, Kuczynski J, Charlson ES et al. Bayesian community-wide culture-independent microbial source tracking. Nat Methods 2011; 8: 761–763

- [139] Davis NM, Proctor DM, Holmes SP et al. Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data. Microbiome 2018; 6: 226
- [140] Burnham P, Gomez-Lopez N, Heyang M et al. Separating the signal from the noise in metagenomic cell-free DNA sequencing. Microbiome 2020; 8: 18
- [141] Dalpke A, Frank J, Peter M et al. Activation of toll-like receptor 9 by DNA from different bacterial species. Infect Immun 2006; 74: 940–946
- [142] Bauer S, Kirschning CJ, Häcker H et al. Human TLR9 confers responsiveness to bacterial DNA via species-specific CpG motif recognition. Proc Natl Acad Sci U S A 2001; 98: 9237–9242
- [143] Oliver JD. Recent findings on the viable but nonculturable state in pathogenic bacteria. FEMS Microbiol Rev 2010; 34: 415–425. doi:10.1111/j.1574-6976.2009.00200.x
- [144] Emerson JB, Adams RI, Román CMB et al. Schrödinger's microbes: Tools for distinguishing the living from the dead in microbial ecosystems. Microbiome 2017; 5: 86
- [145] Rogers GB, Cuthbertson L, Hoffman LR et al. Reducing bias in bacterial community analysis of lower respiratory infections. ISME J 2013; 7: 697–706
- [146] Kim SY, Ko G. Using propidium monoazide to distinguish between viable and nonviable bacteria, MS2 and murine norovirus. Lett Appl Microbiol 2012; 55: 182–188
- [147] Vaara M. Agents that increase the permeability of the outer membrane. Microbiol Rev 1992; 56: 395–411. doi:10.1128/ mr.56.3.395-411.1992
- [148] Selinger DW, Saxena RM, Cheung KJ et al. Global RNA half-life analysis in Escherichia coli reveals positional patterns of transcript degradation. Genome Res 2003; 13: 216–223
- [149] Feezor RJ, Baker HV, Mindrinos M et al. Whole blood and leukocyte RNA isolation for gene expression analyses. Physiol Genomics 2004; 19: 247–254
- [150] Freeman WM, Walker SJ, Vrana KE. Quantitative RT-PCR: Pitfalls and potential. Biotechniques 1999; 26: 112–122. 124–125
- [151] Bustin SA. Quantification of mRNA using real-time reverse transcription PCR (RT-PCR): Trends and problems. J Mol Endocrinol 2002; 29: 23–39
- [152] Domingue GJ, Schlegel JU. Novel bacterial structures in human blood: Cultural isolation. Infect Immun 1977; 15: 621–627. doi:10.1128/ iai.15.2.621-627.1977
- [153] Rajendhran J, Shankar M, Dinakaran V et al. Contrasting circulating microbiome in cardiovascular disease patients and healthy individuals. Int J Cardiol 2013; 168: 5118–5120
- [154] Lelouvier B, Servant F, Païssé S et al. Changes in blood microbiota profiles associated with liver fibrosis in obese patients: A pilot analysis. Hepatology 2016; 64: 2015–2027
- [155] Alvarez-Silva C, Schierwagen R, Pohlmann A et al. Compartmentalization of immune response and microbial translocation in decompensated cirrhosis. Front Immunol 2019; 0:. doi:10.3389/fimmu.2019.00069