

## Correction

**Letrozole vs. Placebo Pretreatment in the Medical Management of First Trimester Missed Miscarriage: a Randomized Controlled Trial**

*Haitham A. Torky, Heba Marie, ElSayed ElDesouky, Samy Gebreel, Osama Raslan, Asem A. Moussa, Ali M. Ahmad, Eman Zain, Mohamed N. Mohsen*

*Geburtsh Frauenheilk 2018; 78: 63–69.*

doi:10.1055/s-0043-122499

**“Material and Methods”**

Under the heading “Randomization and blinding” the sentence should read in full: Concealment was achieved using opaque envelopes, to blind the physicians, while to blind the patients, the placebo tablet was methylcellulose resembling letrozole in shape.

**“Results”**

In the 1st paragraph, the last sentence correctly reads: There were no significant differences in age and gestational age between the two groups (► **Table 1**).

Under the heading “Comparison of gravidity”, the p-value has been corrected in the last sentence of the 1st paragraph. Correct is: p = 0.742.

Under the heading “Comparison of previous cesarean sections”, the p-value has been corrected in the penultimate sentence. Correct is: p = 0.117.

► **Table 1** has been corrected. Correct is:

► **Table 1** Comparison of demographic data.

Demographic data	Group 1 (placebo)	Group 2 (letrozole)	p-value
<b>Age</b>			0.863
Mean ± SD	26.62 ± 4.30	26.52 ± 3.87	
Range	19–40	19–35	
<b>Body mass index</b>			0.0163
Mean ± SD	25.81 ± 2.7	25.12 ± 3.3	
Range	16.53–33.33	19.37–37.46	
<b>Gestational age</b>			0.065
Mean ± SD	48.83 ± 8.00	50.19 ± 7.478	
Range	35–62	35–63	
<b>Gravidity</b>			0.742
Gravida 1	46	54	
Gravida 2	61	55	
Gravida 3	22	22	
Gravida 4	4	7	
Gravida 5	86	81	
<b>Parity</b>			0.891
Para 0	59	66	
Para 1	57	52	
Para 2	20	19	
Para 3	3	4	
Para 4	80	78	
<b>Previous cesarean section (CS)</b>			0.117
No previous CS	92	116	
1 previous CS	92	77	
2 previous CS	21	19	
3 previous CS	11	4	
4 previous CS	3	3	

► **Table 2** has been corrected. Correct is:

► **Table 2** Comparison of failure rates between groups.

Failure rate	Group 1 (placebo)		Group 2 (letrozole)		Chi-square test
	n	%	n	%	p-value
No (complete miscarriage)	85	39	171	78	<0.001
Yes (incomplete miscarriage)	134	61	48	22	

► **Table 3** has been corrected. Correct is:

► **Table 3** Comparison of onset and duration of vaginal bleeding and onset of passing products of conception (clinical data).

Clinical findings	Group 1 (placebo)	Group 2 (letrozole)	p-value
<b>Time to 1st vaginal bleeding after administration of misoprostol, days</b>			<0.001
Mean ± SD	2.30 ± 0.64	1.33 ± 0.29	
Range	1–4	1–3	
<b>Time to induction of abortion after administration of misoprostol, days</b>	3.09 ± 0.99	1.42 ± 0.50	<0.001
Mean ± SD	2–7	1–2	
Range			
<b>Time to 1st product of conception passed after administration of misoprostol, days</b>	3.50 ± 1.55	2.09 ± 0.70	<0.001
Mean ± SD	2–9	1–7	
Range			

► **Table 4** has been corrected. Correct is:

► **Table 4** Comparison of side effects of medication.

Side effects	Group 1 (placebo)		Group 2 (letrozole)		Chi-square test
	No	%	No	%	p-value
Nausea and vomiting	7	3	37	17	0.002
Fever	16	7	9	4	0.152
Severe pain	15	7	24	11	0.132
Severe bleeding needing evacuation	48	22	36	16	0.141