



## Using a self-assembling peptide to achieve endoscopic hemostasis for anastomotic bleeding after ileocecal resection ▶


OPEN ACCESS



▶ **Fig. 1** First emergency endoscopy upon presentation with hematochezia. **a** Endoscopy showed a circumferential anastomotic ulcer with active oozing of blood. **b** Endoscopic hemoclips were applied to the visible bleeding vessel.



▶ **Fig. 2** Second emergency endoscopy showing hematochezia. Endoscopy revealed blood oozing from a circumferential anastomotic ulcer.



▶ **Fig. 3** Third emergency endoscopic image. **a** Endoscopy revealed that the anastomotic ulcer was bleeding again. **b** Hemostasis was achieved by applying 3 mL of PuraStat, a viscous transparent gel, to the circumferential anastomotic ulcer.

A novel synthetic self-assembling peptide, PuraStat (3-D Matrix Ltd, Tokyo, Japan), has been introduced as a surgical or endoscopic hemostatic agent [1,2]. The peptide self-assembles into an extracellular scaffold matrix when activated by a pH change associated with exposure to blood. The matrix forms a stable mechanical barrier at the bleeding site, thereby achieving hemostasis in vivo. In addition, the activated matrix promotes tissue proliferation and facilitates effective healing of the mucosa [2]. Herein,


we report a case of endoscopic hemostasis using PuraStat for postoperative anastomotic bleeding.

A 78-year-old man with chronic heart failure and a history of diabetes mellitus underwent ileocecal resection of the cecal volvulus. The patient was not taking anticoagulants. On postoperative day 7, he presented with hematochezia. Emergency endoscopy revealed active oozing of the blood from a circumferential anastomotic ulcer. Endoscopic hemoclips were applied to visible blood vessels

(▶ **Fig. 1**). However, 6 days later, massive hematochezia recurred (▶ **Fig. 2**). Hemostasis was achieved using additional clips and sodium alginate, and the patient was discharged. Seven days after discharge, the patient was readmitted with recurring hematochezia and anemia (hemoglobin level, 6.2 g/dL). Colonoscopy was performed after blood transfusion. Endoscopy showed blood oozing from the anastomotic ulcer and PuraStat was applied to the anastomotic ulcer to stop the bleeding (▶ **Fig. 3**). After application of PuraStat, the patient presented had no hematochezia and was discharged with an improved hemoglobin level of 10.0 g/dL (▶ **Video 1**). He has not presented with hematochezia for 9 months since his last endoscopy. We plan to perform a follow-up endoscopy 1 year after the last endoscopy.

PuraStat is indicated for hemostasis of oozing bleeding in the parenchyma of solid organs, vascular anastomoses, and capillaries of the gastrointestinal tract [2–5]. In the case described here, PuraStat® was effective in achieving hemostasis of bleeding from postoperative anastomotic ulcers.

## ▶ VIDEO



▶ **Video 1** PuraStat was applied over the circumferential anastomotic ulcer using a delivery catheter (Top Corporation, Tokyo, Japan) inserted through the endoscope accessory channel.

Case A 78-year-old man with a history of chronic heart failure and diabetes mellitus underwent ileocecal resection of the cecal volvulus. He presented with hematochezia after surgery. Emergency endoscopy revealed active oozing of blood from a circumferential anastomotic ulcer. Endoscopic hemoclips were applied to the visible bleeding vessel, but recurrent massive hematochezia occurred. Endoscopic hemostasis was attempted using a self-assembling peptide called PuraStat (3-D Matrix Ltd, Tokyo, Japan). When the endoscope was inserted, a large amount of blood accumulated in the intestinal tract. After drainage, active oozing of blood from a circumferential anastomotic ulcer was noted. PuraStat was applied to the circumferential anastomotic ulcer using a delivery catheter (Top Corporation, Tokyo, Japan) inserted through the endoscope accessory channel. Effective hemostasis was achieved after the application of Purastat.

## Bibliography

Endosc Int Open 2022; 10: E921–E922

DOI 10.1055/a-1816-6450

ISSN 2364-3722

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



## Competing interests

The authors declare that they have no conflict of interest.

## The authors

**Ippei Ikoma\***, **Takashi Murakami\***, **Keiichi Haga\***, **Noboru Yatagai\***, **Hiroya Ueyama\***, **Tomoyoshi Shibuya\***, **Akihito Nagahara\***

Department of Gastroenterology, Juntendo University School of Medicine, Tokyo, Japan

## Corresponding author

**Takashi Murakami, MD PhD**

Department of Gastroenterology, Juntendo University School of Medicine, 2-1-1 Hongo, Bunkyo-ku, Tokyo 113-8421, Japan  
Fax: +81-3-3813-8862  
t-murakm@juntendo.ac.jp

## References

- [1] Yoshida M, Goto N, Kawaguchi M et al. Initial clinical trial of a novel hemostat, TDM-621, in the endoscopic treatments of the gastric tumors. *J Gastroenterol Hepatol* 2014; 29: 77–79
- [2] Uraoka T, Ochiai Y, Fujimoto A et al. A novel fully synthetic and self-assembled peptide solution for endoscopic submucosal dissection-induced ulcer in the stomach. *Gastrointest Endosc* 2016; 83: 1259–1264
- [3] Lee MF, Ma Z, Ananda A. A novel haemostatic agent based on self-assembling peptides in the setting of nasal endoscopic surgery, a case series. *Int J Surg Case Rep* 2017; 41: 461–464
- [4] Giritharan S, Salhiyyah K, Tsang GM et al. Feasibility of a novel, synthetic, self-assembling peptide for suture-line haemostasis in cardiac surgery. *J Cardiothorac Surg* 2018; 13: 68
- [5] de Nucci G, Reati R, Arena I et al. Efficacy of a novel self-assembling peptide hemostatic gel as rescue therapy for refractory acute gastrointestinal bleeding. *Endoscopy* 2020; 52: 773–779

\* These authors contributed equally.