

# Practices and Applications of Heart Rate Variability Monitoring in Endurance Athletes

## Authors

Christopher J Lundstrom , Nicholas A Foreman, George Biltz

## Affiliation

School of Kinesiology, University of Minnesota Twin Cities, Minneapolis, United States

## Key words

cardiac autonomic function, training load, performance, wearable technology, athlete monitoring, time series analysis

accepted 10.05.2022

published online 19.07.2022

## Bibliography

Int J Sports Med 2023; 44: 9–19

DOI 10.1055/a-1864-9726

ISSN 0172-4622

© 2022. Thieme. All rights reserved.

Georg Thieme Verlag, Rüdigerstraße 14, 70469 Stuttgart, Germany

## Correspondence

Dr. Christopher J Lundstrom

University of Minnesota Twin Cities

School of Kinesiology

1900 University Ave SE

55455 Minneapolis

United States

Tel.: 6123817970, Fax: 6126267700

lund0982@umn.edu

## ABSTRACT

Heart rate variability reflects fluctuations in the changes in consecutive heartbeats, providing insight into cardiac autonomic function and overall physiological state. Endurance athletes typically demonstrate better cardiac autonomic function than non-athletes, with lower resting heart rates and greater variability. The availability and use of heart rate variability metrics has increased in the broader population and may be particularly useful to endurance athletes. The purpose of this review is to characterize current practices and applications of heart rate variability analysis in endurance athletes. Important considerations for heart rate variability analysis will be discussed, including analysis techniques, monitoring tools, the importance of stationarity of data, body position, timing and duration of the recording window, average heart rate, and sex and age differences. Key factors affecting resting heart rate variability will be discussed, including exercise intensity, duration, modality, overall training load, and lifestyle factors. Training applications will be explored, including heart rate variability-guided training and the identification and monitoring of maladaptive states such as overtraining. Lastly, we will examine some alternative uses of heart rate variability, including during exercise, post-exercise, and for physiological forecasting and predicting performance.

## Introduction

Athletes today have access to previously unimaginable amounts and types of data and metrics to inform their training, competition, and recovery practices. In this rapidly evolving landscape of athlete analytics, heart rate variability (HRV) remains one of the most broadly utilized tools, particularly for endurance athletes. Further, the applications and the nuanced interpretation of HRV data continue to evolve and inform athlete practices. While the opportunities for enhancing athletic practices are abundant, so are the potential pitfalls. This review provides a brief overview of HRV in the context of endurance athletes, addresses technical considerations, factors affecting daily HRV, the implementation of HRV-guided training, and alternative and future uses of HRV for athletes.

## Definition and physiological basis of HRV

By definition, HRV analysis is done by measuring the beat-by-beat fluctuations of the cardiac rhythm, and assessing the patterns of observed variability over a given time period. This is done through measurement of either R-R intervals (as seen on ECG) or interbeat-intervals of pulse rate. Rather than smoothing the data to obtain the average heart rate (HR), HRV analysis assesses the variability within the sequence of heartbeats. Beat-to-beat variability is a physiological phenomenon that humans have been aware of for hundreds of years, which has been described and explored widely since the early 1970s [1]. The physiological basis of HRV was initially understood as a reflection of an individual's autonomic nervous system (ANS), specifically the sympathetic nervous system (SNS) versus parasympathetic nervous system (PNS), with greater HRV being attributed to increased PNS activity [2]. Crudely stated,

SNS and PNS activity embody, respectively, the “flight or fight” and “rest and digest” responses.

A more nuanced understanding of HRV has emerged, as a phenomenon reflective of complex, non-linear integration of multiple physiological processes including both the intrinsic cardiac nervous system and external regulators such as respiration and blood pressure [3, 4]. Modulation of HR is known to be affected in a non-linear manner by the baroreceptor reflex, which responds to pressure modulations with changes in neural output, classically described as a negative feedback loop, with increased blood pressure leading to rapidly decreasing HR [5, 6].

Respiratory sinus arrhythmia (RSA) is a well-characterized physiological phenomenon, wherein the phases of ventilation and HR interact, with HR slowing during expiration and accelerating with inspiration [7]. Slower respiration with larger tidal volumes increases RSA, and the use of prescribed or “paced” breathing yields different values than measurements taken without paced breathing [8] and is considered a common source of discrepant values in the assessment of HRV [9]. During exercise, coupling (or entrainment) of repetitive movement patterns, such as step rate or cycling cadence, with respiratory rate may occur, complicating the interpretation of HRV due to the effect of respiratory modulating effects [10].

While HRV was previously used extensively in a clinical setting to predict morbidity and mortality [11], a 2003 review paper noted the inconsistent findings and relative dearth of what is known about HRV in athletes [12]. Since that time, HRV has been explored in depth as a means of monitoring training stress and adaptation [13, 14], detecting overtraining [15, 16], as a biofeedback tool [17], and to assess recovery from concussion [18], among other applications. Overall, HRV has been established as a reliable tool that can be useful in monitoring athletes and informing their training practices.

### HRV response to exercise

During exercise, cardiac PNS activity decreases (reaching a nadir at 50–60% of  $\text{VO}_2\text{max}$ ) and SNS activity increases, driving further increases in HR [19]. In contrast with the original model proposing that complete PNS withdrawal occurs during exercise, a more nuanced understanding suggests a process of reciprocal antagonism between branches, with diminished PNS influences retained [20, 21]. While SNS activity predominates at higher intensities, there is evidence to support the measurable effects of PNS activity even at high intensities of exercise [22].

### HRV monitoring for athletes

The use of HRV monitoring for endurance athletes has been studied as far back as the 1990s [23]. Endurance athletes generally exhibit greater resting HRV than healthy controls [24, 25]. This has been observed across a wide range of endurance sports, including rowers, cyclists, Nordic skiers, distance runners [12, 24]. Gradual increases in average HRV appear to correspond with improved aerobic fitness. In a study of distance runners, changes in HRV were correlated with both changes in aerobic capacity and 10-km time [26]. A cross-sectional study of male masters runners also found significant correlations between resting HRV and 10-km race time [27]. In recent years, use of HRV has become more common and

widespread, driven by the ease, availability, and portability of monitoring tools. It is now commonplace for athletes, coaches, and sports medicine staff to collect HRV data, with short-term measurements taken first thing in the day frequently recommended [13]. Given the accessibility of tools and technologies, the use of HRV is likely to become more commonplace. Thus, an understanding of the different measures, techniques, applications, and the complex physiological interactions that produce differing patterns of HRV can help guide this practice and improve interpretation of the data.

## Technical considerations

### Analysis techniques

The explosion of HRV research in 1980s and early 1990s led to the establishment of the “Task Force” guidelines for standardization of analysis procedures and considerations [2]. The numerous analysis methods have been broadly categorized into time, frequency, and nonlinear domains [28]. While the Task Force Standards focused on the more traditional time and frequency domain measures, a variety of nonlinear methods of HRV analysis have been developed and widely utilized in more recent years [29, 30].

The most commonly utilized time domain measures are the standard deviation of the normal-normal intervals (SDNN) and the root mean square of the successive differences (RMSSD) [28]. In brief, SDNN characterizes the average difference from the mean, whereas RMSSD captures the average difference from the previous interval. Thus, SDNN is a measure of the overall variability of the time series, whereas RMSSD captures short-term variability. Given the more rapid effect of the PNS than the SNS on HR, RMSSD is considered a reflection of PNS activity [9]. Of all the measures of HRV, RMSSD (or the natural log of RMSSD –  $\ln\text{RMSSD}$ ) is perhaps the most broadly utilized owing to its low typical error, ease of calculation, and robustness to differences in respiratory patterns [13].

Frequency domain (or spectral analysis) measures are calculated by transforming the time domain data, most typically through the fast Fourier method, to a frequency spectrum of cycles per second (Hz), reflecting the contribution of various frequencies to the overall variability of the signal [2]. There are typically three peaks observed: Very low frequency (VLF,  $\leq 0.04$  Hz), low frequency (LF, 0.04–0.15 Hz), and high frequency HF, 0.15–0.4 Hz). Traditional theory proposed HF power as reflective of PNS activity and LF power as capturing a combination of SNS and PNS activity. This theory has been challenged and the notion of LF/HF as a reflection of sympathovagal balance is likely an oversimplification at best [19, 31]. Spectral analysis may be particularly sensitive to the effects of respiration and ectopic beats, leading some to advocate for the use of ECG and requiring a respiratory rate greater than 10 breaths per minute in order to reliably use these measures in athletes [32].

Nonlinear methods of analysis, including detrended fluctuation analysis (DFA) move beyond traditional means of quantifying variability, attempting to capture the complexity of the time series [29]. Complexity, in brief, refers to the amount of information, across multiple time scales, contained within a system or set of information [29]. Complex systems contain multiple interconnected feedback loops, such that the integrated whole contains proper-

ties that cannot be understood simply through examination of the component parts [28].

Sample entropy (SampEn) measures the relative regularity of the data sequence by quantifying template matches across a data sequence [33]. More frequent matches indicates more regularity, and therefore a lower entropy score, whereas a higher entropy score indicates more irregularity and less predictability in the data sequence [28, 33]. One benefit of SampEn is that it has been shown to be relatively reliable with short time series, though a minimum of 200 data points has been recommended [34].

Detrended fluctuation analysis is a nonlinear technique developed to distinguish between internal fluctuations and external perturbations to the system, making it a preferred method for non-static conditions such as exercise [35]. Patterns of variation across multiple measurement scales of differing window lengths are examined, with local trends subtracted. The scaling exponent  $\alpha$  is related to the length of the time window, where  $\alpha_1$  indicates short-range correlations and  $\alpha_2$  specifies long-term correlations. An  $\alpha$  value of 0.5 is indicative of no correlations (random data) and an alpha value of 1.5 is equivalent to highly correlated data. Values of  $\alpha \sim 1.0$  are consistent with complex systems data where self-similarity across time scales is observed. An optimal middle range can make traditional statistical analysis of group data challenging, as initially high and low values may both move toward the middle following training [36].

## Monitoring tools

The gold standard of HRV monitoring is through the use of ECG, though chest strap heart rate monitors have been used extensively and have shown high reliability [37]. Pulse rate variability, as measured through photoplethysmography (PPG) has been used increasingly owing to the ease of use [38]. The accuracy of PPG may depend on the tool and condition in which it is used. Watches with embedded PPG have been shown to be relatively accurate at rest [39]. On the other hand, accuracy during exercise depends on the model of watch and type of exercise, with notable increases in error during exercise that is either vigorous or involves arm movement [40]. Similarly, PPG technology embedded in a finger clip pulse oximeter are reliable but prone to artifact when used during movement [41]. Resting measurement of HRV through a finger placed on the camera of a smartphone is another common use of PPG, with high accuracy reported when done according to suggested practices [38]. The ubiquity of smartphones and the ease of use of this technique makes it appealing to a broad population.

## Stationarity and body position

Stationarity of data refers to the stability of the state of the individual, such that the mean and standard deviation of the data set remain consistent over the recording period [28]. Resting or steady state exercise data satisfy the definition of stationarity, but data collected in the transition from rest to exercise violates the requirement of stationarity [35]. Stationarity is an important consideration for most HRV measures [28]. A notable exception, DFA, corrects for non-stationarity [35], though the efficacy of this correction has been challenged [42].

Given the sensitivity of HRV measures to physiological state, the need for consistency of data collection procedures is clear. Research on body position during data collection illustrates this. Numerous studies have compared resting HRV in different body positions including supine, seated, and standing [15, 43, 44]. The highest HRV values are typically seen in the supine condition, with decreases in the seated condition and further diminishment in the standing condition. Seated measures have been suggested to be more strongly associated with changes in fitness level than supine measures [45]. Thus, when tracking HRV within an individual, or comparing with population norms, consistency of body position during measurement is a critical consideration.

## Timing and duration of recording window

Measurements can be affected by such factors as time of day of the recording and duration of data collection. While 24-hour HRV monitoring has been utilized extensively in clinical and research settings [2], such long-term monitoring may be impractical for athletes. Further, it includes periods of varying activity, and therefore will fluctuate extensively over such an extended window of time [46]. Resting-only measurements of 30 minutes [47] or as short as 5 minutes have been recommended broadly [2] or for athletes in particular [13]. Ultra-short term measurements of one minute have been shown to have acceptable agreement with 5-minute recordings, while 30- and 10-second readings significantly decreased the level of agreement [48].

Time of day may affect HRV measurements. Long-term measurements show the highest levels of HRV during sleep [49]. The lowest levels of HRV are observed during the late morning and early afternoon hours [50]. Thus, use of either overnight readings [51] or at a standardized time of day such as first thing upon waking in the morning [13], are recommended to reduce the confounding effects associated with time of day. Strong correlations have been observed between morning and nocturnal HRV measures in young athletes [52].

## Adjustment for average heart rate

Many measures of HRV are reported in time-specific units, often without adjusting for mean HR. In such cases, a mathematical association between HRV and HR exists, with a slower average HR associated with higher HRV values and vice versa [8, 53]. If no adjustment is made for mean HR, the same absolute fluctuation is indicative of a different magnitude of variability, relative to the mean. Unadjusted HRV, accordingly, includes information not only on the variability of the data but also on the mean. Both time and frequency domain – but not nonlinear measures – are affected by this collinearity [33, 35].

## Sex and age differences

Population norms and applications in HRV have been described in pediatric and adolescent [54, 55], geriatric [56], and in various clinical populations [57, 58]. A large meta-analysis of healthy individuals including all ages showed higher HR and lower SDNN but greater HF power in women than men [59]. Females were shown to have 7% higher HF power in the supine condition than males, but experienced a greater magnitude of reductions in HRV with standing [44]. Men peak in LF power in the morning hours, while women do

so in the afternoon [49]. An analysis of data collected through a commercial app found no sex differences on average, but among females HRV was 3.2% higher during the luteal than follicular phase of the menstrual cycle [60]. The same study found a moderate negative correlation between age and RMSSD. Other studies have found that, with aging, adults decrease in both overall HRV and in magnitude of variation over the course of a 24-hour period [49, 61].

## Factors affecting resting HRV

The most widely used application of HRV for athletes, and increasingly among other health-conscious individuals, is the daily HRV measurement. It is typically taken first thing in the morning in order to minimize confounding effects [13], yet some devices alternatively measure nocturnal HRV during sleep, which has the advantage of less potential for influence of extrinsic factors [51]. Previously available only to researchers and the technically savvy, daily HRV can now be monitored at home using a variety of devices, apps, or software. Some of these options provide raw data while others utilize HRV as part of black box algorithms to provide the user with a recovery score or strain score [62]. Popular brands that may be familiar to the general public include Whoop, Apple Watch, Fitbit, and Oura Ring. Many durations and types of measures are utilized by such devices, which can influence the outcome measurements. For the practical purposes of daily monitoring, consistency of measurement for the specific individual is the most important consideration.

Daily monitoring can provide insight into autonomic response to training. Undoubtedly, there are individual differences in HRV response to training, and longitudinal tracking of individual athletes allows greater insight into how these measures are best utilized for the individual [13, 47]. Population differences such as training status, cardiorespiratory fitness, age, sex, and other factors may all affect not only daily HRV readings but also how they respond to different training and lifestyle stimuli. However, some common observations of daily HRV responses can provide a framework for understanding that may help athletes interpret their responses.

## Overall training load

Chronic training may lead to increased HRV in athletes, especially where training includes a strong aerobic component as is common practice among endurance athletes [12, 24]. However, acute reductions in daily HRV have been observed in response to heavy training [63–65], and days with decreased morning HRV may indicate a reduced performance ability [66]. Athletes with overtraining syndrome have been shown to exhibit reduced HRV compared to healthy, trained athletes [15]. Plotted against training load, a bell-shaped curve for HF power was observed in marathon runners, indicating high parasympathetic activity at moderate levels of training, and lower levels with both abnormally high and low training loads [67]. Similar suppression of HF power has been observed when tracking elite rowers over the course of a year of training and competition [68]. A study of elite-level French Nordic skiers over four years found suppression of HRV parameters in those tested in a fatigued state, as determined by a validated questionnaire [69]. Those in the fatigued state also exhibited greater intra-subject variance, indicating that HRV response may vary between individuals

in response to fatigue [69]. In a population of middle- and long-distance runners and triathletes, an intensive 2-week training camp induced significant reductions in HRV, which rebounded within 3–4 days of rest [47]. This is consistent with an “over-reached” state, in which a heavy training load temporarily alters autonomic function but is reversible within a few days of reduced training.

## Exercise intensity

The intensity of an exercise session appears to have an influence on its impact on HRV measures, either immediately or in some cases the day after exercise. For example, HRV was suppressed in highly trained endurance athletes after exercise above but not below the first ventilatory threshold (VT1), regardless of duration [70]. In the same study, highly trained (national class) athletes returned to baseline more quickly than trained (non-elite) athletes. Similarly, comparing 3,500 m and 7,000 m runs at low (50% of  $\text{VO}_2\text{max}$ ), moderate (63% of  $\text{VO}_2\text{max}$ ), and vigorous (74% of  $\text{VO}_2\text{max}$ ; 3,500 m only) intensity, HRV was higher after the low intensity sessions, regardless of duration [71].

Comparing 30 minutes of exercise at 45%, 60%, and 75% of  $\text{VO}_2\text{max}$ , nocturnal HR was elevated with the higher intensity exercise, but there was no effect on HRV [72]. In a study comparing interval training at 75 and 95% of  $\text{VO}_2\text{max}$ , the higher intensity interval session led to a suppression of HRV at 1-hour post-exercise, but it returned to baseline within the first 24 hours [73]. A high-intensity interval training session (5-km run in 1-min increments at peak velocity attained during a  $\text{VO}_2\text{max}$  test, with 1-min passive recovery) produced a moderate to large effect-size suppression of HRV at 30 min and 1 hour after exercise but small effect size differences from baseline between 4 and 24 hours after exercise [74]. On the other hand, a similar study of ten intervals of 1-min duration at 85% of HR reserve suppressed HRV only immediately after exercise, with a return to baseline by the 1-hour mark, continuing through 8 hours after exercise [75].

In a study comparing 30-minute supine cycling bouts at three intensities (2, 3, and 4 mmol of lactate), HRV suppression was present in all three conditions 5 minutes after exercise, in the 3- and 4-mmol conditions 15 minutes after exercise but had returned to baseline in all three conditions by 30 minutes after exercise [76]. Similarly, participants undergoing a 300-kcal exercise bout at either 50% or 80% of  $\text{VO}_2$  reserve returned to baseline HRV levels within 30 minutes after exercise, though levels remained suppressed longer (up to 25 minutes) after the higher intensity condition [77].

Most studies of a single exercise bout have found minimal disturbance to HRV from the single session, though most of the sessions were not particularly challenging in intensity. In contrast, when comparing constant velocity exercise at VT1 with the same overall workload on a bike, but including 9 × 1 min of maximal intensity intervals, HRV was suppressed for both conditions in the hour immediately afterwards but with greater suppression in the high intensity group [78]. Further, there may be some effect of body position on the persistence of HRV suppression, as decreased HRV was observed in the standing but not supine conditions at 24 and 48 hours after exercise in both conditions [78].

The fitness level of the participant may impact the level of intensity that will lead to prolonged perturbation of HRV. After a

maximal intensity 75-km ski race, for example, HRV was suppressed one but not two days after the race, and the time to return to normal was inversely correlated with individual's  $\text{VO}_2\text{max}$  [79]. However, HRV measured after a marathon and an hour at 70% HRmax in recreational runners found no relationship between  $\text{VO}_2\text{max}$  and nocturnal perturbation of HRV [80].

### Exercise duration

Exercise duration is an important component of overall training load, and increases in overall exercise volume have been shown to decrease HRV [63–65]. However, the amount of training volume necessary to produce HRV suppression may be population-dependent and driven by relative difference rather than absolute volume. For example, in highly trained endurance athletes, HRV measures were unaffected by a 60- and 120-minute runs, provided they were performed below VT1 [70]. Even in the immediate post-exercise period, HRV returns to baseline levels fairly quickly after endurance exercise, provided it is of low to moderate intensity, such as 50 and 63% of  $\text{VO}_2\text{max}$  [71] or below VT1 [70]. On the other hand, in a population of moderately active non-athletes, a 90-minute moderate (60% of  $\text{VO}_2\text{max}$ ) session reduced nocturnal HRV compared to control, whereas 30- and 60-minute sessions at the same intensity did not [72]. In ultra-endurance events, decreased HRV (measured as HF power) and decreased baroreceptor sensitivity have been observed in both elite and recreational athletes [81].

### Exercise modality

Numerous studies have been done examining HRV in response to different types of training interventions, including continuous aerobic endurance, high-intensity interval, resistance, coordinative, and multimodal exercise [82]. However, few studies have compared the effects of different modalities of endurance exercise on athletes. A study of healthy young men performing three modalities of maximal exercise tests (cycling, walking, and running) found that HRV recovered more quickly after the cycling test, which also elicited a lower maximal HR [83]. A few studies have assessed differences in HRV during exercise in different modalities, though the challenges of matching intensity across different activities make it difficult to draw conclusions [19]. Research has focused primarily on land-based sports without restrictions on breathing, which may alter the interpretation of HRV data [84].

### Lifestyle factors

The sensitivity of HRV to lifestyle factors such as life stress, diet, alcohol consumption, smoking, sleep, and body composition has been well-established in literature examining HRV as a means of assessing cardiometabolic risk [85, 86]. It is not unreasonable to expect that endurance athletes may also be subject to the influence of such factors on resting HRV.

Partial sleep deprivation in the form of a reduction of hours of sleep to 3 or 4.5 hours has been shown to decrease HRV [87, 88]. Alcohol use may also impact HRV, with a dose-response relationship. Only small effects were observed from a low dose (0.3 g/kg) [89], and significantly more suppression of HRV was seen with two drinks when compared with one [90]. In addition to the more obvious deleterious effects of smoking on endurance athletes, smoking has been shown to acutely reduce HRV and negatively impact leg

cycling performance [91]. The use of other drugs and medications consumed either acutely or chronically, particularly those acting upon the autonomic nervous system, may also have a marked impact on HRV [92]. Given that the “stress response” is typically characterized by a reduction in parasympathetic activity, it is not surprising that some types of life stress have been shown to reduce HRV [93]. These may include perceived emotional stress [94], anxiety [95], depression [96], and transient stress before bed [97]. Self-perception may modify HRV, as the changes in HRV due to a mental stress task were smaller in students with higher self-esteem [98].

## Training prescription based on daily HRV

### HRV-guided training

The use of HRV-guided training prescription has been investigated in the literature and is being used in the real world to make training decisions [99–103]. Most studies of this approach have compared a training schedule with a set schedule of workouts with one that varies the inclusion of higher intensity work depending on the daily HRV score. Typically, an individual baseline resting HRV is established over a period ranging from 3 days to 4 weeks [99, 100]. Protocols of HRV-guided training vary, but typically moderate and higher intensity workouts are performed only when the HRV score falls within the normal range [99, 104]. Lower intensity workouts and/or rest days are prescribed until HRV returns to normal.

Defining the physiologically normal range is done on an individual basis, using the smallest worthwhile change [100, 105] established over a number of days. A number of applications, such as HRV4Training and EliteHRV [106], iThlete [107], and Welltory [108] have been validated for measurement of daily HRV and can be used by athletes to establish baseline levels and assess daily recovery status. Alternative approaches to the smallest worthwhile change have been studied. For example, vigorous training was prescribed if HRV either increased or stayed the same, while a decreased training load was prescribed when the HRV decreased [101, 103]. The interpretation of increases in HRV above the normal range remains a subject of debate, and one where individual variability of response may come into play [13].

A systematic review on HRV-guided training for runners found greater improvements in performance in the HRV-based groups than in those who followed a predetermined training plan [99]. In most cases, the HRV-guided training led to reduced volume of moderate and high intensity training. Another review including healthy cyclists, runners, and skiers, ranging from untrained to elite, found HRV-guided training to be as effective as but not superior to pre-determined training [104]. A meta-analysis of studies on HRV-guided training for endurance athletes found significantly greater improvements in submaximal exercise measures, whereas performance and maximal exercise measures did not significantly differ between groups [109]. However, there were proportionally more positive responders and fewer negative responders to HRV-guided training groups.

An 8-week intervention in distance runners found greater improvements in 3-km time trial in the HRV-guided training group compared to the traditional training group [100]. In a recreational active population, the use of two different HRV-guided training

approaches compared to a pre-planned program resulted in similar gains in  $\text{VO}_{2\text{peak}}$ , though women in the study gained the same benefits with fewer vigorous intensity sessions [101]. Another study from the same group also favored the HRV-guided training group in maximal running velocity and  $\text{VO}_{2\text{peak}}$  [103].

In endurance trained males, there were greater improvements in peak treadmill velocity, countermovement jump, and resting HRV in the time and frequency domains when compared to a pre-determined block training paradigm, with no difference in 3-km running performance [110]. In a study of professional runners, HRV-guided training improved peak treadmill velocity and peak respiratory exchange ratio [111]. Unlike previous studies, where HRV-guided groups achieve comparable performance through fewer sessions above VT1, the HRV-guided group in this study spent more time between VT1 and VT2, raising the possibility that HRV-guided training may be beneficial for some runners who are able to handle more high-intensity sessions.

When implementing HRV-guided training, practitioners should be aware of the divergent changes in HRV seen with modifications in training volume or intensity [112]. Further, it is worth noting that all of these studies compared a pre-determined training plan with HRV-guided training, whereas athletes often work with an exercise professional such as a coach who may adapt training according to athlete response.

## HRV and maladaptive states

In addition to its use in guiding training, HRV has been suggested as a means of detecting and assessing severity of maladaptive states, including illness [113, 114], overtraining [15, 115], and concussion [18]. Concussion is an area of great interest in sport science research in recent years, but is much more prevalent among team sport athletes. Illness and overtraining, on the other hand, are common among endurance athletes.

Overtraining is thought to be a result of an imbalance between training and other stresses and recovery, leading to symptoms such as disturbed sleep, fatigue, mood disturbances, altered hormone levels, prolonged muscle soreness, and a reduction in exercise capacity [116]. Detection of overtraining syndrome is challenging, and with no clear diagnostic criteria, it often becomes a diagnosis of exclusion [117]. Overtrained athletes have been observed to have reduced HRV in both supine and upright positions [15]. On the other hand, a case study of a junior cross-country skiers experiencing overtraining symptoms showed an increase in HF and total power in the supine position [118]. One study showed an increased strength of relationship between RR interval length and HF power in overtrained endurance athletes [115]. In a group of elite canoeists engaged in a 6-day intensive training camp,  $\text{VO}_{2\text{max}}$  and maximal lactate decreased, but HRV measures did not change [119]. This suggests the possible use of HRV to distinguish between overtraining and short-term training fatigue.

A recent meta-analysis found a negative correlation between HRV and inflammatory processes, such as illness [113]. In a critical care setting, HRV suppression has been used to predict severity of illness and mortality risk [120, 121]. Less severe illness may also impact HRV. In elite-level swimmers, HRV was suppressed during weeks characterized by upper respiratory or pulmonary infections [114]. However, the predictive ability of HRV in this study was un-

clear; during the week prior to infection, HRV was higher in the supine position but lower in the upright position.

The potential for HRV monitoring to successfully identify early-stage illness and/or overtraining remains to be determined. In the face of such health issues, it is unclear whether HRV could be utilized to successfully adapt and adjust in order to reduce severity or duration of illness and other medical issues. However, HRV-guided training and monitoring of health status using HRV and other biometric markers represent potentially useful tools to assist athletes and coaches in navigating such challenges.

## Alternative HRV monitoring options

### Exercise HRV

Exercise leads to an increase in heart rate to meet increased metabolic demands, while time and frequency HRV measures decrease dramatically [19, 122–125]. Although there are differences between the methods as well as possibly between modes of exercise, the general pattern is that HRV decreases in a dose-response relationship with exercise intensity. At moderate intensities, HRV reaches a relatively stable, low level, and there are minimal changes with further increases in exercise intensity [19]. As a result, these measures are difficult to interpret in the context of exercise at the levels of intensity at which athletes often train. Evaluation of HRV during exercise is not as common as resting HRV at rest. It is, however, fairly reliable and reproducible, though reliability may be compromised at very high intensities [126].

Nonlinear measures of exercise HRV have been used to monitor and adjust exercise intensity during a discrete workout session. Specifically, a DFA $\alpha$ 1 value of 0.75 has been suggested to be equivalent to the aerobic threshold as measured using ventilatory data [127]. Monitoring applications such as Heart Rate Variability Logger and FatMaxxer use DFA $\alpha$ 1 calculated in real time to adjust exercise intensity [127, 128]. Though it can “detrend” and adjust for changes in exercise intensity, staying at the same exercise intensity for a minimum of two minutes is recommended for these applications.

### Post-exercise HRV

As noted in the section above on factors affecting daily HRV, recovery of HRV upon cessation of exercise may differ between exercise sessions. The monitoring of HR and HRV recovery after exercise may provide valuable insight into the stress of the session. Recovery back to baseline HRV levels commonly occurs within the hour after exercise [19, 70, 71, 75, 77, 78, 129–133]. Low-intensity exercise may even increase HRV immediately after exercise if the intensity is appreciably low [71, 134]. However, exercise at intensities above VT1 appears to contribute to greater or longer periods of HRV suppression [70, 78, 133, 135]. Nighttime monitoring may also be of value, as Hynynen et al. [80] showed moderate to strong correlations between nocturnal HRV and post-exercise suppression of HRV. Post-exercise HRV may also be correlated with improvements in anaerobic fitness, as post-exercise RMSSD was correlated with improvements in repeated sprint ability after a nine-week training program [136]. The monitoring of post-exercise HRV as a tool for HRV-guided training has not been explored to our



knowledge, but could serve as a useful tool for endurance athletes, particularly those engaged in high training volumes and/or multiple training sessions per day.

### Future directions

Given the abundance of data and metrics available to coaches and athletes, there is increased interest in physiological forecasting, or predicting outcomes or performances based on an array of variables that may be collected using wearable technology and other means. These technologies may be able to quantify stress and recovery outside of the training period to improve quantification of training load [137]. It seems clear that HRV alone is not linearly predictive of performance, as it has been observed that HRV may diminish slightly during the peaking phase [13]. Additionally, changes in endurance performance are not always coupled with changes in HRV or other metrics used to monitor training such as the countermovement jump [138]. However, observation of individual patterns and responses to training, carried out longitudinally, allow for an understanding of individual physiological response to training and other stimuli. In one study, correlations between HRV and measures of training load were not significant between individuals, but were significant within individuals when using repeated-measures correlations [139]. This and other studies support the notion that exercise prescription may be improved with longitudinal HRV monitoring. Some recreational runners may improve more with high volumes of endurance exercise instead of greater intensity, and these differences are correlated with changes in nocturnal HRV [140]. Baseline HRV at the start of a training period may also be related to improvements in training in a recreational population [141]. These factors should be considered when attempting to guide training using HRV in combination with physiological and psychological variables. Exploration of practices such as HRV-biofeedback training, which have been used successfully used to increase HRV [8], could also be assessed as a training tool.

Previous research applying HRV to sport has focused largely on endurance sport. While that population is the focus of this review, it is worth noting that endurance athletes often also engage in resistance exercise and anaerobic activities. Little is currently known regarding the use of HRV for guiding resistance training or exercise prescription in team sport environments. A study of recreationally active adults engaged in high-intensity functional training found similar improvements in body composition, strength, and  $\text{VO}_2$  peak when training was guided by HRV, despite a 20% reduction in high-intensity sessions over a 9-week period [142]. Resistance exercise suppresses HRV but may [143] or may not [144] effectively differentiate differences in fatigue from varying exercise bouts. The timeline of HRV recovery may differ from other variables, including countermovement jump and psychological variables. Nuutila et al. [135] observed different fatigue recovery patterns for countermovement jump and post-exercise HRV depending on exercise intensity. Similarly, Vacher et al. [145] noted linear changes in HRV despite nonlinear changes in perceived stress and recovery. Thus, HRV data can be viewed as one piece of a puzzle, contributing to improvements in predicting performance, enhancing training and competition strategies, and refining the science of endurance sport.

## Conclusion

Many endurance athletes at all levels are using HRV in a variety of applications. Increased availability and ease of access to tools for measuring and processing HRV data carry the promise and possibility for further enhancement of training and competition practices. However, athletes and coaches are cautioned to use a rigorous approach to controlling for the quality of data being collected and to be aware of key variables that can affect HRV measurements. Practices such as daily HRV monitoring and HRV-guided training are well-established, promising applications that should be considered by endurance athletes. Finally, the use of HRV should be understood as one piece in the complex network of physiological regulation and adaptation. Future research integrating a variety of metrics – including HRV – promise to further our understanding of endurance performance and training, and to aid individual endurance athletes in developing a nuanced understanding of their own physiology.

### Conflict of Interest

The authors declare that they have no conflict of interest.

### References

- [1] Billman GE. Heart rate variability – a historical perspective. *Front Physiol* 2011; 2: 86. doi:10.3389/fphys.2011.00086
- [2] Heart rate variability: standards of measurement, physiological interpretation and clinical use. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Circulation* 1996; 93: 1043–1065
- [3] Shaffer F, McCraty R, Zerr CL. A healthy heart is not a metronome: an integrative review of the heart's anatomy and heart rate variability. *Front Psychol* 2014; 5: 1040. doi:10.3389/fpsyg.2014.01040
- [4] Ernst G. Heart-rate variability – more than heart beats? *Front Public Health* 2017; 5: 240. doi:10.3389/fpubh.2017.00240
- [5] Rea RF, Eckberg DL. Carotid baroreceptor-muscle sympathetic relation in humans. *J Appl Physiol* 1987; 253: R929–R934
- [6] Eckberg DL. Nonlinearities of the human carotid baroreceptor-cardiac reflex. *Circ Res* 1980; 47: 208–216. doi:10.1161/01.RES.47.2.208
- [7] Denver JW, Reed SF, Porges SW. Methodological issues in the quantification of respiratory sinus arrhythmia. *Biol Psychol* 2007; 74: 286–294. doi:10.1016/j.biopsycho.2005.09.005
- [8] Shaffer F, Ginsberg JP. An overview of heart rate variability metrics and norms. *Front Public Health* 2017; 5: 258. doi:10.3389/fpubh.2017.00258
- [9] Nunan D, Sandercock GRH, Brodie DA. A quantitative systematic review of normal values for short-term heart rate variability in healthy adults. *Pacing Clin Electrophysiol* 2010; 33: 1407–1417. doi:10.1111/j.1540-8159.2010.02841.x
- [10] Blain G, Meste O, Blain A et al. Time-frequency analysis of heart rate variability reveals cardiocomotor coupling during dynamic cycling exercise in humans. *Am J Physiol Heart Circ Physiol* 2009; 296: H1651–H1659. doi:10.1152/ajpheart.00881.2008
- [11] Goldberger AL, West BJ. Fractals in physiology and medicine. *Yale J Biol Med* 1987; 60: 421–435
- [12] Aubert AE, Seps B, Beckers F. Heart rate variability in athletes. *Sports Med* 2003; 33: 889–919. doi:10.2165/00007256-200333120-00003

- [13] Plews D, Laursen P, Stanley J et al. Training adaptation and heart rate variability in elite endurance athletes: Opening the door to effective monitoring. *Sports Med* 2013; 43: 773–781. doi:10.1007/s40279-013-0071-8
- [14] Schneider C, Hanakam F, Wiewelshove T et al. Heart rate monitoring in team sports – a conceptual framework for contextualizing heart rate measures for training and recovery prescription. *Front Physiol* 2018; 9: 639. doi:10.3389/fphys.2018.00639
- [15] Mourot L, Bouhaddi M, Perrey S et al. Decrease in heart rate variability with overtraining: assessment by the Poincaré plot analysis. *Clin Physiol Funct Imaging* 2004; 24: 10–18
- [16] Hynynen E, Uusitalo A, Konttinen N et al. Heart rate variability during night sleep and after awakening in overtrained athletes. *Med Sci Sports Exerc* 2006; 38: 313–317. doi:10.1249/01.mss.0000184631.27641.b5
- [17] Pagaduan JC, Chen Y-S, Fell JW et al. A preliminary systematic review and meta-analysis on the effects of heart rate variability biofeedback on heart rate variability and respiration of athletes. *J Complement Integr Med* 2021. Online ahead of print. doi:10.1515/jcim-2020-0528
- [18] Senthinathan A, Mainwaring LM, Hutchison M. Heart rate variability of athletes across concussion recovery milestones: a preliminary study. *Clin J Sport Med* 2017; 27: 288–295. doi:10.1097/JSM.0000000000000337
- [19] Michael S, Graham KS, Davis GMO. Cardiac autonomic responses during exercise and post-exercise recovery using heart rate variability and systolic time intervals—a review. *Front Physiol* 2017; 8: 301. doi:10.3389/fphys.2017.00301
- [20] White DW, Raven PB. Autonomic neural control of heart rate during dynamic exercise: revisited. *J Physiol* 2014; 592: 2491–2500. doi:10.1113/jphysiol.2014.271858
- [21] Levy MN. Sympathetic-parasympathetic interactions in the heart. *Circ Res* 1971; 29: 437–445. doi:10.1161/01.res.29.5.437
- [22] Kannankeril PJ, Le FK, Kadish AH et al. Parasympathetic effects on heart rate recovery after exercise. *J Investig Med* 2004; 52: 394–401. doi:10.1136/jim-52-06-34
- [23] Aubert A, Ramaekers D, Cuche Y et al. Effect of long term physical training on heart rate variability. *Comput Cardiol* 1996; 22: 17–20. doi:10.1109/CIC.1996.542462
- [24] da Silva VP, de Oliveira NA, Silveira H et al. Heart rate variability indexes as a marker of chronic adaptation in athletes: a systematic review. *Ann Noninvasive Electrocardiol* 2015; 20: 108–118. doi:10.1111/anec.12237
- [25] Bellenger CR, Fuller JT, Thomson RL et al. Monitoring athletic training status through autonomic heart rate regulation: a systematic review and meta-analysis. *Sports Med* 2016; 46: 1461–1486. doi:10.1007/s40279-016-0484-2
- [26] Buchheit M, Chivot A, Parouty J et al. Monitoring endurance running performance using cardiac parasympathetic function. *Eur J Appl Physiol* 2010; 108: 1153–1167. doi:10.1007/s00421-009-1317-x
- [27] Cataldo A, Bianco A, Paoli A et al. Resting sympatho-vagal balance is related to 10 km running performance in master endurance athletes. *Eur J Transl Myol* 2018; 28: 1025–1033. doi:10.4081/ejtm.2018.7051
- [28] Seely AJE, Macklem PT. Complex systems and the technology of variability analysis. *Crit Care* 2004; 8: R367–R384. doi:10.1186/cc2948
- [29] Henriques T, Ribeiro M, Teixeira A et al. Nonlinear methods most applied to heart-rate time series: a review. *Entropy (Basel)* 2020; 22: 309
- [30] Sassi R, Cerutti S, Lombardi F et al. Advances in heart rate variability signal analysis: joint position statement by the e-Cardiology ESC Working Group and the European Heart Rhythm Association co-endorsed by the Asia Pacific Heart Rhythm Society. *Europace* 2015; 17: 1341–1353. doi:10.1093/europace/euv015
- [31] Billman GE. The LF/HF ratio does not accurately measure cardiac sympatho-vagal balance. *Front Physiol* 2013; 4: 26. doi:10.3389/fphys.2013.00026
- [32] Lucini D, Marchetti I, Spataro A et al. Heart rate variability to monitor performance in elite athletes: criticalities and avoidable pitfalls. *Int J Cardiol* 2017; 240: 307–312. doi:10.1016/j.ijcard.2017.05.001
- [33] Richman JS, Moorman JR. Physiological time-series analysis using approximate entropy and sample entropy. *Am J Physiol Heart Circ Physiol* 2000; 278: H2039–H2049. doi:10.1152/ajpheart.2000.278.6.H2039
- [34] Yentes JM, Hunt N, Schmid KK et al. The appropriate use of approximate entropy and sample entropy with short data sets. *Ann Biomed Eng* 2013; 41: 349–365. doi:10.1007/s10439-012-0668-3
- [35] Peng CK, Havlin S, Stanley HE et al. Quantification of scaling exponents and crossover phenomena in nonstationary heartbeat time series. *Chaos* 1995; 5: 82–87. doi:10.1063/1.166141
- [36] Heffernan KS, Sosnoff JJ, Fahs CA et al. Fractal scaling properties of heart rate dynamics following resistance exercise training. *J Appl Physiol* (1985) 2008; 105: 109–113. doi:10.1152/jappphysiol.00150.2008
- [37] Giles D, Draper N, Neil W. Validity of the Polar V800 heart rate monitor to measure RR intervals at rest. *Eur J Appl Physiol* 2016; 116: 563–571. doi:10.1007/s00421-015-3303-9
- [38] Plews DJ, Scott B, Altini M et al. Comparison of heart-rate-variability recording with smartphone photoplethysmography, Polar H7 chest strap, and electrocardiography. *Int J Sports Physiol Perform* 2017; 12: 1324–1328. doi:10.1123/ijssp.2016-0668
- [39] Wang R, Blackburn G, Desai M et al. Accuracy of wrist-worn heart rate monitors. *JAMA Cardiol* 2017; 2: 104–106
- [40] Gillinov S, Etiwy M, Wang R et al. Variable accuracy of wearable heart rate monitors during aerobic exercise. *Med Sci Sports Exerc* 2017; 49: 1697–1703. doi:10.1249/MSS.0000000000001284
- [41] Lu G, Yang F. Limitations of oximetry to measure heart rate variability measures. *Cardiovasc Eng* 2009; 9: 119–125. doi:10.1007/s10558-009-9082-3
- [42] Bryce RM, Sprague KB. Revisiting detrended fluctuation analysis. *Sci Rep* 2012; 2: 315. doi:10.1038/srep00315
- [43] Pomeranz B, Macaulay RJ, Caudill MA et al. Assessment of autonomic function in humans by heart rate spectral analysis. *Am J Physiol* 1985; 248: H151–H153. doi:10.1152/ajpheart.1985.248.1.H151
- [44] Hnatkova K, Šišáková M, Smetana P et al. Sex differences in heart rate responses to postural provocations. *Int J Cardiol* 2019; 297: 126–134. doi:10.1016/j.ijcard.2019.09.044
- [45] Rabbani M, Agha-Alinejad H, Gharakhanlou R et al. Monitoring training in women's volleyball: supine or seated heart rate variability? *Physiol Behav* 2021; 240: 113537. doi:10.1016/j.physbeh.2021.113537
- [46] Saul JP, Albrecht P, Berger RD et al. Analysis of long term heart rate variability: methods, 1/f scaling and implications. *Comput Cardiol* 1988; 14: 419–422
- [47] Voss A. Longitudinal analysis of heart rate variability. *J Electrocardiol* 2007; 40: S26–S29. doi:10.1016/j.jelectrocard.2006.10.024
- [48] Esco MR, Flatt AA. Ultra-short-term heart rate variability indexes at rest and post-exercise in athletes: evaluating the agreement with accepted recommendations. *J Sports Sci Med* 2014; 13: 535–541
- [49] Yamasaki Y, Kodama M, Matsuhisa M et al. Diurnal heart rate variability in healthy subjects: effects of aging and sex difference. *Am J Physiol* 1996; 271: H303–H310. doi:10.1152/ajpheart.1996.271.1.H303
- [50] Jarczok MN, Guendel H, McGrath J et al. Circadian rhythms of the autonomic nervous system: scientific implication and practical implementation. In: Svorc P (Ed). *Chronobiology - The Science of Biological Time Structure*. London: IntechOpen; 2019. doi:10.5772/intechopen.86822



- [51] Sapplier N, Bauer A. Nocturnal heart rate variability: a population-based screening tool? *Heart Rhythm* 2022; 19: 640–641
- [52] Mishica C, Kyröläinen H, Hynynen E et al. Evaluation of nocturnal vs. morning measures of heart rate indices in young athletes. *PLoS One* 2022; 17: e0262333
- [53] Sacha J. Interaction between heart rate and heart rate variability. *Ann Noninvasive Electrocardiol* 2014; 19: 207–216. doi:10.1111/anec.12148
- [54] Gąsior JS, Sacha J, Pawłowski M et al. Normative values for heart rate variability parameters in school-aged children: simple approach considering differences in average heart rate. *Front Physiol* 2018; 9: 1495. doi:10.3389/fphys.2018.01495
- [55] Sharma VK, Subramanian SK, Arunachalam V et al. Heart rate variability in adolescents – normative data stratified by sex and physical activity. *J Clin Diagn Res* 2015; 9: CC08–CC13. doi:10.7860/JCDR/2015/15373.6662
- [56] Ogliari G, Mahinrad S, Stott DJ et al. Resting heart rate, heart rate variability and functional decline in old age. *CMAJ* 2015; 187: E442–E449. doi:10.1503/cmaj.150462
- [57] Stys A, Stys T. Current clinical applications of heart rate variability. *Clin Cardiol* 1998; 21: 719–724. doi:10.1002/clc.4960211005
- [58] Tsai C-H, Ma H-P, Lin Y-T et al. Usefulness of heart rhythm complexity in heart failure detection and diagnosis. *Sci Rep* 2020; 10: 14916. doi:10.1038/s41598-020-71909-8
- [59] Koenig J, Thayer JF. Sex differences in healthy human heart rate variability: a meta-analysis. *Neurosci Biobehav Rev* 2016; 64: 288–310. doi:10.1016/j.neubiorev.2016.03.007
- [60] Altini M, Plews D. What is behind changes in resting heart rate and heart rate variability? A large-scale analysis of longitudinal measurements acquired in free-living. *Sensors (Basel)* 2021; 21: 7932. doi:10.3390/s21237932
- [61] Bonnemeier H, Wiegand UKH, Brandes A et al. Circadian profile of cardiac autonomic nervous modulation in healthy subjects. *J Cardiovasc Electrophysiol* 2003; 14: 791–799. doi:https://doi.org/10.1046/j.1540-8167.2003.03078.x
- [62] Kaewkannate K, Kim S. A comparison of wearable fitness devices. *BMC Public Health* 2016; 16: 433. doi:10.1186/s12889-016-3059-0
- [63] Buchheit M. Monitoring training status with HR measures: do all roads lead to Rome? *Front Physiol* 2014; 5: 73. doi:10.3389/fphys.2014.00073
- [64] Pichot V, Busso T, Roche F et al. Autonomic adaptations to intensive and overload training periods: a laboratory study. *Med Sci Sports Exerc* 2002; 34: 1660–1666. doi:10.1097/00005768-200210000-00019
- [65] Pichot V, Roche F, Gaspoz J-M et al. Relation between heart rate variability and training load in middle-distance runners. *Med Sci Sports Exerc* 2000; 32: 1729–1736. doi:10.1097/00005768-200010000-00011
- [66] Deblauw JA, Crawford DC, Stein JA et al. Association of heart rate variability and simulated cycling time trial performance. *J Sci Cycling* 2022; 10: 25–33. doi:10.28985/1221.jsc.09
- [67] Manzi V, Castagna C, Padua E et al. Dose-response relationship of autonomic nervous system responses to individualized training impulse in marathon runners. *Am J Physiol Heart Circ Physiol* 2009; 296: 1733–1740. doi:10.1152/ajpheart.00054.2009
- [68] Iellamo F, Pigozzi F, Spataro A et al. Autonomic and psychological adaptations in Olympic rowers. *J Sports Med Phys Fitness* 2006; 46: 598–604
- [69] Schmitt L, Regnard J, Desmarests M et al. Fatigue shifts and scatters heart rate variability in elite endurance athletes. *PLoS One* 2013; 8: e71588. doi:10.1371/journal.pone.0071588
- [70] Seiler S, Haugen O, Kuffel E. Autonomic recovery after exercise in trained athletes: intensity and duration effects. *Med Sci Sports Exerc* 2007; 39: 1366–1373. doi:10.1249/mss.0b013e318060f17d
- [71] Kaikkonen P, Nummela A, Rusko H. Heart rate variability dynamics during early recovery after different endurance exercises. *Eur J Appl Physiol* 2007; 102: 79–86. doi:10.1007/s00421-007-0559-8
- [72] Myllymäki T, Rusko H, Syväoja H et al. Effects of exercise intensity and duration on nocturnal heart rate variability and sleep quality. *Eur J Appl Physiol* 2012; 112: 801–809. doi:10.1007/s00421-011-2034-9
- [73] James DVB, Munson SC, Maldonado-Martin S et al. Heart rate variability: effect of exercise intensity on postexercise response. *Res Q Exerc Sport* 2012; 83: 533–539. doi:10.1080/02701367.2012.10599142
- [74] Panissa VLG, Cal Abad CC, Julio UF et al. High-intensity intermittent exercise and its effects on heart rate variability and subsequent strength performance. *Front Physiol* 2016; 7: 81. doi:10.3389/fphys.2016.00081
- [75] Burma JS, Copeland P, Macaulay A et al. Effects of high-intensity intervals and moderate-intensity exercise on baroreceptor sensitivity and heart rate variability during recovery. *Appl Physiol Nutr Metab* 2020; 45: 1156–1164. doi:10.1139/apnm-2019-0810
- [76] Gladwell VF, Sandercock GRH, Birch SL. Cardiac vagal activity following three intensities of exercise in humans. *Clin Physiol Funct Imaging* 2010; 30: 17–22. doi:10.1111/j.1475-097X.2009.00899.x
- [77] Parekh A, Lee CM. Heart rate variability after isocaloric exercise bouts of different intensities. *Med Sci Sports Exerc* 2005; 37: 599–605. doi:10.1249/01.mss.0000159139.29220.9a
- [78] Mourot L, Bouhaddi M, Tordi N et al. Short- and long-term effects of a single bout of exercise on heart rate variability: comparison between constant and interval training exercises. *Eur J Appl Physiol* 2004; 92: 508–517. doi:10.1007/s00421-004-1119-0
- [79] Hautala A, Tulppo MP, Mäkikallio TH et al. Changes in cardiac autonomic regulation after prolonged maximal exercise. *Clin Physiol* 2001; 21: 238–245. doi:10.1046/j.1365-2281.2001.00309.x
- [80] Hynynen E, Vesterinen V, Rusko H et al. Effects of moderate and heavy endurance exercise on nocturnal HRV. *Int J Sports Med* 2010; 31: 428–432. doi:10.1055/s-0030-1249625
- [81] Foulds HJA, Cote AT, Phillips AA et al. Characterisation of baroreflex sensitivity of recreational ultra-endurance athletes. *Eur J Sport Sci* 2014; 14: 686–694. doi:10.1080/17461391.2014.884169
- [82] Grässler B, Thielmann B, Böckelmann I et al. Effects of different training interventions on heart rate variability and cardiovascular health and risk factors in young and middle-aged adults: a systematic review. *Front Physiol* 2021; 12: 657274
- [83] Cunha FA, Midgley AW, Gonçalves T et al. Parasympathetic reactivation after maximal CPET depends on exercise modality and resting vagal activity in healthy men. *Springerplus* 2015; 4: 100. doi:10.1186/s40064-015-0882-1
- [84] Solana-Tramunt M, Morales J, Buscà B et al. Heart-rate variability in elite synchronized swimmers. *Int J Sports Physiol Perform* 2019; 14: 464–471. doi:10.1123/ijspp.2018-0538
- [85] Marsac J. Heart rate variability: a cardiometabolic risk marker with public health implications. *Bull Acad Natl Med* 2013; 197: 175–186
- [86] Jandackova VK, Scholes S, Britton A et al. Healthy lifestyle and cardiac vagal modulation over 10 years: Whitehall II cohort study. *J Am Heart Assoc* 2019; 8: e012420. doi:10.1161/JAHA.119.012420
- [87] Dettoni JL, Consolim-Colombo FM, Drager LF et al. Cardiovascular effects of partial sleep deprivation in healthy volunteers. *J Appl Physiol* (1985) 2012; 113: 232–236. doi:10.1152/jappphysiol.01604.2011
- [88] Bourdillon N, Jeanneret F, Nilchian M et al. Sleep deprivation deteriorates heart rate variability and photoplethysmography. *Front Neurosci* 2021; 15: 642548. doi:10.3389/fnins.2021.642548

- [89] Gonzalez Gonzalez J, Mendez Llorens A, Mendez Novoa A et al. Effect of acute alcohol ingestion on short-term heart rate fluctuations. *J Stud Alcohol* 1992; 53: 86–90. doi:10.15288/jsa.1992.53.86
- [90] Spaak J, Tomlinson G, McGowan CL et al. Dose-related effects of red wine and alcohol on heart rate variability. *Am J Physiol Heart Circ Physiol* 2010; 298: H2226–H2231. doi:10.1152/ajpheart.00700.2009
- [91] Chen C-L, Tang J-S, Li P-C et al. Immediate effects of smoking on cardiorespiratory responses during dynamic exercise: arm vs. leg ergometry. *Front Physiol* 2015; 6: 376. doi:10.3389/fphys.2015.00376
- [92] Vanoli E, Cerati D, Pedretti RF. Autonomic control of heart rate: pharmacological and nonpharmacological modulation. *Basic Res Cardiol* 1998; 93: 133–142. doi:10.1007/s003950050239
- [93] Verkuil B, Brosschot JF, Tollenaar MS et al. Prolonged non-metabolic heart rate variability reduction as a physiological marker of psychological stress in daily life. *Ann Behav Med* 2016; 50: 704–714. doi:10.1007/s12160-016-9795-7
- [94] Dishman RK, Nakamura Y, Garcia ME et al. Heart rate variability, trait anxiety, and perceived stress among physically fit men and women. *Int J Psychophysiol* 2000; 37: 121–133. doi:10.1016/S0167-8760(00)00085-4
- [95] Chalmers JA, Quintana DS, Abbott MJA et al. Anxiety disorders are associated with reduced heart rate variability: a meta-analysis. *Front Psychiatry* 2014; 5: 80. doi:10.3389/fpsyg.2014.00080
- [96] Kemp AH, Quintana DS, Gray MA et al. Impact of depression and antidepressant treatment on heart rate variability: a review and meta-analysis. *Biol Psychiatry* 2010; 67: 1067–1074. doi:10.1016/j.biopsych.2009.12.012
- [97] Hall M, Vasko R, Buysse D et al. Acute stress affects heart rate variability during sleep. *Psychosom Med* 2004; 66: 56–62. doi:10.1097/01.PSY.0000106884.58744.09
- [98] O'Donnell K, Brydon L, Wright CE et al. Self-esteem levels and cardiovascular and inflammatory responses to acute stress. *Brain Behav Immun* 2008; 22: 1241–1247. doi:10.1016/j.bbi.2008.06.012
- [99] Düking P, Zinner C, Reed JL et al. Predefined vs. data-guided training prescription based on autonomic nervous system variation: a systematic review. *Scand J Med Sci Sports* 2020; 30: 2291–2304. doi:10.1111/sms.13802
- [100] Vesterinen V, Nummela A, Heikura I et al. Individual endurance training prescription with heart rate variability. *Med Sci Sports Exerc* 2016; 48: 1347–1354. doi:10.1249/MSS.0000000000000910
- [101] Kiviniemi AM, Hautala AJ, Kinnunen H et al. Daily exercise prescription on the basis of HR variability among men and women. *Med Sci Sports Exerc* 2010; 42: 1355–1363
- [102] Tornberg J, Ikaheimo TM, Kiviniemi A et al. Physical activity is associated with cardiac autonomic function in adolescent men. *PLoS One* 2019; 14: e0222121. doi:10.1371/journal.pone.0222121
- [103] Kiviniemi AM, Hautala AJ, Kinnunen H et al. Endurance training guided individually by daily heart rate variability measurements. *Eur J Appl Physiol* 2007; 101: 743–751. doi:10.1007/s00421-007-0552-2
- [104] Ruiz JPM, Rubio-Arias J, Clemente-Suarez VJ et al. Effectiveness of training prescription guided by heart rate variability versus predefined training for physiological and aerobic performance improvements: a systematic review and meta-analysis. *Appl Sci* 2020; 10: 8532. doi:10.3390/app10238532
- [105] Plews DJ, Laursen PB, Kilding AE et al. Heart rate variability in elite triathletes, is variation in variability the key to effective training? A case comparison. *Eur J Appl Physiol* 2012; 112: 3729–3741. doi:10.1007/s00421-012-2354-4
- [106] Stone JD, Ulman HK, Ulman HK et al. Assessing the accuracy of popular commercial technologies that measure resting heart rate and heart rate variability. *Front Sport Act Living* 2021; 3: 585870. doi:10.3389/fspor.2021.585870
- [107] Flatt AA, Esco MR. Validity of the ithlete smart phone application for determining ultra-short-term heart rate variability. *J Hum Kinet* 2013; 39: 85–92. doi:10.2478/hukin-2013-0071
- [108] Moya-Ramon M, Mateo-March M, Peña-González I et al. Validity and reliability of different smartphones applications to measure HRV during short and ultra-short measurements in elite athletes. *Comput Methods Programs Biomed* 2022; 217: 106696. doi:10.1016/j.cmpb.2022.106696
- [109] Düking P, Zinner C, Trabelsi K et al. Monitoring and adapting endurance training on the basis of heart rate variability monitored by wearable technologies: a systematic review with meta-analysis. *J Sci Med Sport* 2021; 24: 1180–1192
- [110] Nuutila O-P, Nikander A, Polomoshnov D et al. Effects of HRV-guided vs. predetermined block training on performance, HRV and serum hormones. *Int J Sports Med* 2017; 38: 909–920. doi:10.1055/s-00043-115122
- [111] Carrasco-Poyatos M, González-Quílez A, Altini M et al. Heart rate variability-guided training in professional runners: effects on performance and vagal modulation. *Physiol Behav* 2022; 244. doi:10.1016/j.physbeh.2021.113654
- [112] Nuutila O-P, Ari N, Heikki K et al. Physiological, perceptual, and performance responses to the 2-wk block of high- versus low-intensity endurance training. *Med Sci Sports Exerc* 2022; 54: 851–860. doi:10.1249/mss.0000000000002861
- [113] Williams DP, Koenig J, Carnevali L et al. Heart rate variability and inflammation: a meta-analysis of human studies. *Brain Behav Immun* 2019; 80: 219–226. doi:10.1016/j.bbi.2019.03.009
- [114] Hellard P, Guimaraes F, Avalos M et al. Modeling the association between HR variability and illness in elite swimmers. *Med Sci Sports Exerc* 2011; 43: 1063–1070. doi:10.1249/MSS.0b013e318204de1c
- [115] Kiviniemi AM, Tulppo MP, Hautala AJ et al. Altered relationship between R-R interval and R-R interval variability in endurance athletes with overtraining syndrome. *Scand J Med Sci Sports* 2014; 24: e77–e85. doi:10.1111/sms.12114
- [116] Lehmann M, Foster C, Keul J. Overtraining in endurance athletes: a brief review. *Med Sci Sports Exerc* 1993; 25: 854–862. doi:10.1249/00005768-199307000-00015
- [117] Meeusen R, Duclos M, Foster C et al. Prevention, diagnosis, and treatment of the overtraining syndrome: joint consensus statement of the European College of Sport Science and the American College of Sports Medicine. *Med Sci Sports Exerc* 2013; 45: 186–205. doi:10.1249/MSS.0b013e318279a10a
- [118] Hedelin R, Wiklund U, Bjerle P et al. Cardiac autonomic imbalance in an overtrained athlete. *Med Sci Sports Exerc* 2000; 32: 1531–1533
- [119] Hedelin R, Kenttä G, Wiklund U et al. Short-term overtraining: effects on performance, circulatory responses, and heart rate variability. *Med Sci Sports Exerc* 2000; 32: 1480–1484. doi:10.1097/00005768-200008000-00017
- [120] Norris PR, Ozdas A, Cao H et al. Cardiac uncoupling and heart rate variability stratify ICU patients by mortality: a study of 2088 trauma patients. *Ann Surg* 2006; 243: 804–814. doi:10.1097/01.sla.0000219642.92637.fd
- [121] Winchell RJ, Hoyt DB. Spectral analysis of heart rate variability in the ICU: a measure of autonomic function. *J Surg Res* 1996; 63: 11–16. doi:10.1006/jsre.1996.0214
- [122] Tulppo MP, Makikallio TH, Takala TE et al. Quantitative beat-to-beat analysis of heart rate dynamics during exercise. *Am J Physiol Heart Circ Physiol* 1996; 271: H244–H252. doi:10.1152/ajpheart.1996.271.1.h244
- [123] Pichon AP, De Bisschop C, Roulaud M et al. Spectral analysis of heart rate variability during exercise in trained subjects. *Med Sci Sports Exerc* 2004; 36: 1702–1708. doi:10.1249/01.MSS.0000142403.93205.35

- [124] Perini R, Orizio C, Baselli G et al. The influence of exercise intensity on the power spectrum of heart rate variability. *Eur J Appl Physiol Occup Physiol* 1990; 61: 143–148. doi:10.1007/BF00236709
- [125] Sandercock GRH, Brodie DA. The use of heart rate variability measures to assess autonomic control during exercise. *Scand J Med Sci Sports* 2006; 16: 302–313
- [126] McNarry MA, Lewis MJ. Heart rate variability reproducibility during exercise. *Physiol Meas* 2012; 33: 1123–1133. doi:10.1088/0967-3334/33/7/1123
- [127] Rogers B, Giles D, Draper N et al. A new detection method defining the aerobic threshold for endurance exercise and training prescription based on fractal correlation properties of heart rate variability. *Front Physiol* 2021; 11: 1806
- [128] Gronwald T, Rogers B, Hoos O. Fractal correlation properties of heart rate variability: a new biomarker for intensity distribution in endurance exercise and training prescription? *Front Physiol* 2020; 11: 1152
- [129] Cabral-Santos C, Giacon TR, Campos EZ et al. Impact of high-intensity intermittent and moderate-intensity continuous exercise on autonomic modulation in young men. *Int J Sports Med* 2016; 37: 431–435. doi:10.1055/s-0042-100292
- [130] Castrillon CIM, Miranda RAT, Cabral-Santos C et al. High-intensity intermittent exercise and autonomic modulation: effects of different volume sessions. *Int J Sports Med* 2017; 38: 468–472. doi:10.1055/s-0042-121898
- [131] Martinmäki K, Rusko H. Time-frequency analysis of heart rate variability during immediate recovery from low and high intensity exercise. *Eur J Appl Physiol* 2008; 102: 353–360. doi:10.1007/s00421-007-0594-5
- [132] Terziotti P, Schena F, Gulli G et al. Post-exercise recovery of autonomic cardiovascular control: a study by spectrum and cross-spectrum analysis in humans. *Eur J Appl Physiol* 2001; 84: 187–194. doi:10.1007/s004210170003
- [133] Perkins SE, Jelinek HF, Al-Aubaidy HA et al. Immediate and long term effects of endurance and high intensity interval exercise on linear and nonlinear heart rate variability. *J Sci Med Sport* 2017; 20: 312–316. doi:10.1016/j.jsams.2016.08.009
- [134] Raczak G, Pinna GD, La Rovere MT et al. Cardiovascular response to acute mild exercise in young healthy subjects. *Circ J* 2005; 69: 976–980
- [135] Nuutila O-P, Kyröläinen H, Häkkinen K et al. Acute physiological responses to four running sessions performed at different intensity zones. *Int J Sports Med* 2021; 42: 513–522. doi:10.1055/a-1263-1034
- [136] Buchheit M, Millet GP, Parisy A et al. Supramaximal training and postexercise parasympathetic reactivation in adolescents. *Med Sci Sports Exerc* 2008; 40: 362–371. doi:10.1249/mss.0b013e31815aa2ee
- [137] Treff G, Leppich R, Winkert K et al. The integration of training and off-training activities substantially alters training volume and load analysis in elite rowers. *Sci Rep* 2021; 11: 17218. doi:10.1038/s41598-021-96569-0
- [138] Nuutila OP, Nummela A, Häkkinen K et al. Monitoring training and recovery during a period of increased intensity or volume in recreational endurance athletes. *Int J Environ Res Public Health* 2021; 18: 2401. doi:10.3390/ijerph18052401
- [139] Davletyarova K, Vacher P, Nicolas M et al. Associations between heart rate variability-derived indexes and training load. *J Strength Cond Res* 2022; 36: 2005–2010. doi:10.1519/JSC.0000000000003760
- [140] Vesterinen V, Häkkinen K, Laine T et al. Predictors of individual adaptation to high-volume or high-intensity endurance training in recreational endurance runners. *Scand J Med Sci Sports* 2016; 26: 885–893. doi:10.1111/sms.12530
- [141] Vesterinen V, Häkkinen K, Hynynen E et al. Heart rate variability in prediction of individual adaptation to endurance training in recreational endurance runners. *Scand J Med Sci Sports* 2013; 23: 171–180. doi:10.1111/j.1600-0838.2011.01365.x
- [142] Deblauw JA, Drake NB, Kurtz BK et al. High-intensity functional training guided by individualized heart rate variability results in similar health and fitness improvements as predetermined training with less effort. *J Funct Morphol Kinesiol* 2021; 6: 102. doi:10.3390/jfmk6040102
- [143] Pareja-Blanco F, Rodríguez-Rosell D, Sánchez-Medina L et al. Acute and delayed response to resistance exercise leading or not leading to muscle failure. *Clin Physiol Funct Imaging* 2017; 37: 630–639. doi:10.1111/cpf.12348
- [144] De Souza JC, Tibana RA, Cavaglieri CR et al. Resistance exercise leading to failure versus not to failure: effects on cardiovascular control. *BMC Cardiovasc Disord* 2013; 13: 105. doi:10.1186/1471-2261-13-105
- [145] Vacher P, Filaire E, Mourot L et al. Stress and recovery in sports: effects on heart rate variability, cortisol, and subjective experience. *Int J Psychophysiol* 2019; 143: 25–35. doi:10.1016/j.ijpsycho.2019.06.011