

Effect of Aerobic Exercise on Arterial Stiffness in Individuals with Different Smoking Statuses



Authors

Xianxuan Wang^{1#}, Guanzhi Chen^{2#}, Zegui Huang¹, Yiran Zang³, Zefeng Cai¹, Xiong Ding⁴, Zekai Chen⁵, Yulong Lan¹, Weijian Li¹, Wei Fang¹, Weiqiang Wu¹, Zhichao Chen¹, Shouling Wu⁶, Youren Chen¹

Affiliations

- 1 Department of Cardiology, Second Affiliated Hospital of Shantou University Medical College, Shantou, China
- 2 Second Clinical College, China Medical University, Shenyang, China
- 3 Graduate School, North China University of Science and Technology, Tangshan, China
- 4 School of Public Health, North China University of Science and Technology, Tangshan, China
- 5 Department of Epidemiology, University Medical Centre Groningen, Groningen, Netherlands
- 6 Department of Cardiology, Kailuan General Hospital, Tangshan, China

Key words

aerobic exercise, pulse wave velocity, arterial stiffness, smoking

accepted 28.07.2022

published online 04.11.2022

Bibliography

Int J Sports Med 2023; 44: 48–55

DOI 10.1055/a-1925-7588

ISSN 0172-4622

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

Correspondence

Mr. Youren Chen
Department of Cardiology
Second Affiliated Hospital of Shantou University
Medical College
No.69, Dongxia North Road
515000 Shantou
China
Tel.: +86-15507883729
13902779840@139.com



Supplementary Material is available under <https://doi.org/10.1055/a-1925-7588>

ABSTRACT

This study aimed to investigate the immediate effects of acute bout of aerobic exercise on arterial stiffness in individuals with different smoking statuses. A total of 940 male individuals (mean age of 36.82 ± 7.76 years) in the Kailuan study cohort were selected to participate in the fifth National Physical Fitness Monitoring. All participants completed measurements of brachial – ankle pulse wave velocity (baPWV) before and after twice-quantitative cycle ergometer exercise. Four groups were defined: (1) non-smokers ($n = 231$), (2) former smokers ($n = 165$), (3) light smokers (1–10 cigarettes/day, $n = 254$), (4) heavy smokers (> 10 cigarettes/day, $n = 290$). Generalized linear models were established to analyze between-group differences in the change in baPWV before and after acute aerobic exercise in individuals with different smoking statuses. Overall, after acute aerobic exercise, baPWV was immediately decreased significantly (-33.55 cm/s [95%CI, -39.69 to -27.42]). Compared with non-smokers, former smokers, light smokers, and heavy smokers showed a greater decrease in baPWV (-12.17 cm/s [95%CI, -30.08 to 5.75], -18.43 cm/s [95%CI, -34.69 to -2.16], and -22.46 cm/s [95%CI, -38.39 to -6.54]) respectively. There is a transient decrease in baPWV in individuals with different smoking statuses. Compared with non-smokers, baPWV decreased more significantly in light and heavy smokers.

These authors have contributed equally to this work

ABBREVIATIONS

BP	blood pressure
SBP	systolic blood pressure
DBP	diastolic blood pressure
baPWV	brachial – ankle pulse wave velocity
HR	heart rate,
MAP	mean arterial pressure
CI	confidence interval
TG	Triglyceride
FBG	fasting blood glucose
VO ₂ max	maximum oxygen consumption
CV	coefficients of variation

Introduction

Arterial stiffness is not only a sign of vascular aging, but is also an independent risk factor for cardiovascular diseases, renal failure, cognitive dysfunction, and all-cause mortality [1–4]. Observational and longitudinal cohort studies have shown that age and blood pressure (BP) are the main risk factors for arterial stiffness [5, 6]. However, traditional risk factors, such as diabetes, smoking, and chronic inflammation, are also risk factors for arterial stiffness [7–9]. Arterial stiffness is significantly increased after smoking compared with before smoking in healthy young people [10]. Additionally, Tomiyama et al. (2010) found that the progression of arterial stiffness was significantly faster in smokers than in non-smokers of the same age [11].

In 2010, the American Heart Association suggested that an ideal healthy lifestyle, such as being a non-smoker, moderate exercise, and a healthy diet, can reduce the risk of cardiovascular disease and all-cause mortality [12]. In particular, long-term regular aerobic exercise is an effective lifestyle intervention that reduces the damage to blood vessels caused by risk factors, such as smoking, hypertension, and diabetes [13–15]. However, the results of research on the immediate effect of acute bout of aerobic exercise on arterial stiffness in individuals with different smoking statuses are inconsistent [16, 17]. Doonan et al. (2011) studied 24 smokers and found that arterial stiffness increased immediately after aerobic exercise [16]. A study of 50 healthy non-smokers showed that arterial stiffness decreased immediately after aerobic exercise [17]. These studies were small-sample studies on a Caucasian population.

To the best of our knowledge, few studies have investigated and compared the effect of acute aerobic exercise-induced changes on arterial stiffness in Asian men with different smoking statuses. Therefore, we used data from the Kailuan Study to evaluate short-term changes in arterial stiffness following an acute aerobic exercise test in men with different smoking statuses.

Materials and Methods

Study design and participants

The data of this study were derived from the subjects in the Kailuan Study population who were selected to participate in the fifth National Fitness Monitoring. The Kailuan Study began in 2006 and

participants were followed up every 2 years thereafter in 2008–2009, 2010–2011, 2012–2013, 2014–2015, 2016–2017, and 2018–2019 [18, 19]. To systematically assess the nationwide physical fitness of individuals, the State Sports General Administration initiated the fifth National Physique Monitoring in 2020. In addition to the general population, special monitoring also had been carried out on the public security, finance, construction and coal industries of typical workers [20].

All the participants in this study are workers engaged in the coal industry. According to the Labor Law, few female employees were underground coal miners or workers in dangerous occupation [21]. Therefore, 1,200 male coal miners aged 20–49 years were selected by random methods as subjects of the fifth National Physique Monitoring, using the employee information database reported and established by the four affiliated companies of the Kailuan Group. All participants were asked to perform these tests, including body morphology, physical function, and physical fitness. We also measured brachial–ankle pulse wave velocity (baPWV) before and immediately after acute aerobic exercise.


This study was performed in accordance with the Declaration of Helsinki and was approved by the Ethics Committee of the Kailuan General Hospital. All participants provided written informed consent.

Inclusion and exclusion criteria

In this study, a total of 1,200 employees of the Kailuan Group participated in the fifth National Physique Monitoring, of whom 1,138 participated in the 2018–2019 follow-up of the Kailuan Study. Those who did not have baPWV measured twice before and after exercise ($n = 185$) and those without a twice-quantitative cycle ergometer exercise ($n = 13$) were excluded. Finally, 940 individuals were included for analysis (► Fig. 1).

Exercise protocol

The procedure of performing cycling exercises complies with the Exercise Standards for Testing and Training released by the American Heart Association (AHA) and is briefly described below [22]. All participants and measurements were performed in a quiet, temperature-controlled room (22 °C–25 °C). Before cycling exercise, the participants had avoided alcohol, tobacco, ingest caffeine or perform vigorous activities within 12 hours. After a preliminary pe-



► Fig. 1 Inclusion/exclusion flowchart for study participants. Caption: The flowchart of 940 participants included in the final analyses.

riod of 15 minutes of supine rest, pre-exercise measurements of baPWV were made in the supine position. Cycling exercise was then performed with individually tailored ramp protocols via upright cycle ergometry (GMCS-GLC3, Beijing, China). Following a brief warm-up by cycling against unloaded exercise (0 watts) for 30 seconds, according to individual's fitness conditions, participants started to cycle at a moderate intensity (50 watts–80 watts) for 3 minutes so that the heart rate reached 60% ~ 80% of their estimated maximum heart (calculated as 220 minus age in years), and the resistance was increased by 25watts in next 3 minutes. Throughout the exercise, participants were instructed to maintain a pedaling cadence of 60 revolutions per minute (rpm). The cycling exercise finished with 30 seconds of unloaded (0 watts). HR was monitored and recorded continuously. Maximum oxygen consumption (VO₂max) was indirectly estimated based on the cycling exercise and calculated by the ergometry according to a previously described equation [23].

BaPWV measurement

All participants underwent baPWV measurement after 12 h abstinence from smoking and fasting. We used the BP-203 RPE III networked arterial stiffness detection device (Omron Health Medical [China] Co., Ltd.) automation to measure baPWV. We used the BP-203 RPE III networked arterial stiffness detection device (Omron Health Medical [China] Co., Ltd.) automation to measure baPWV. BaPWV is determined by measuring the pulse waves of the brachial artery and ankle, and dividing the distance between the upper arm and ankle (DAB), which is calculated using linear regression of body height, by the time difference between the two pulses (TAB) [24]. Briefly, the calculation formula is expressed as $baPWV = DAB / TAB$. BaPWV was recorded before and immediately (or as soon as possible) after cycling exercise, which simultaneously records BP, HR following standard operation procedures [25]. For the first measurement, participants in light clothes had not been smoking and were seated for at least 15 minutes in a room with the temperature controlled between 22 °C and 25 °C. The participants were asked to lie down on an examination couch in the supine position and remain quiet during the measurement. Cuffs were wrapped on both arms and ankles. The lower edges of the arm cuffs were positioned 2–3 cm above the transverse striation of the cubital fossa, while the lower edges of the ankle cuffs were positioned 1–2 cm above the superior aspect of the medial malleolus. One heart sound detector was placed at the left and right edges of the sternum. Measurements were repeated twice for each subject, and the result of the second time was used as the final result.

The maximum of the left and right sides of baPWV was used for analysis.

Smoking status

The smoking status of the subjects (i. e., non-smokers, former smokers, and current smokers) was assessed with a questionnaire. We ensured the accuracy of smoking information through a telephone follow-up survey. Non-smokers were defined as having no history of smoking, former smokers [26] as not smoking for at least 6 months, and current smokers [27] as any who had smoked at least one cigarette per day for 1 year. Additionally, to calculate pack-

years in current smokers, data on the average number of cigarettes smoked per day and the years of cigarette smoking were collected. Pack-years of smoking were defined as the average number of packs of cigarettes smoked per day multiplied by the duration of smoking in years [28]. On the basis of the number of cigarettes smoked per day, current smokers were classified into the two following groups: light smokers (1–10 cigarettes/day) and heavy smokers (> 10 cigarettes/day) [29]. To further confirm our findings, we divided the smokers into light smokers (≤ 4 pack-years) and heavy smokers (> 4 pack-years) in accordance with the pack-years as described in Celermajer et al.'s (1993) [30] study as well.

Data collection

Information on demographic data and lifestyle factors (e. g., age, sex, alcohol consumption) and medical history (e. g., diabetes, hypertension) were collected using a self-reported questionnaire. Anthropometric indicators, including height, weight were derived from the fifth National Physique Monitoring data. BMI was calculated as weight (kg) divided by height squared (m²). The physical exercise was defined as taking exercise ≥ 3 times/week with at least 30 mins each time [31]. All participants underwent blood tests after 12 h fasting at each physical examination. The blood samples were analyzed using an auto-analyzer (Hitachi 747, Hitachi, Tokyo, Japan) on the day of the blood draw. The biochemical indicators tested included fasting blood glucose (FBG), triglycerides (TG) levels. Hypertension was defined as (i) systolic blood pressure (SBP) ≥ 140 mmHg or diastolic blood pressure (DBP) ≥ 90 mmHg, or (ii) formerly diagnosed hypertension or using antihypertensive drugs [32]. Diabetes was defined as either a fasting blood glucose (FBG) level ≥ 7.0 mmol/L, self-reporting of a physician's diagnosis, or self-reported use of antidiabetic medication [33].

Statistical analysis

Continuous variables that conformed to a normal distribution are presented as the mean \pm standard deviation ($\bar{x} \pm s$). Data with a non-normal distribution are expressed as the median (interquartile range). Categorical variables are expressed as the number of samples (percentage). Demographic factors and hemodynamic parameters were compared between the groups using the one-way analysis of variance or the Wilcoxon rank sum test, as appropriate. Categorical variables were assessed using the χ^2 test for independence. The paired-sample t test was used to compare baPWV, HR, and BP values in each group post-exercise and pre-exercise.

Linear regression models were established to evaluate group changes in baPWV, HR and BP post-exercise and pre-exercise. In the linear regression models, we adjusted for age (continuous variable), the pre-exercise value of BMI (continuous variable), and TG level (continuous variable), education level (categorical variable), alcohol consumption (categorical variable), physical activity (categorical variable) and the timing of post-exercise measurements (continuous variable).

Generalized linear models were used to analyze differences in the change in baPWV between the groups. In generalized linear models, we adjusted for age and the pre-exercise values of baPWV in model 1. In model 2, we further adjusted for the pre-exercise values of BMI, HR, MAP, and TG level, the timing of post-exercise measurements, education level, alcohol consumption, physical activity.

In model 3, we further adjusted for the change in MAP and HR. Between-group differences are described by β -value and 95% confidence interval (CI).

Stratified analyses were performed based on the different physical activity levels among participants. To test the robustness of our findings, the following sensitivity analyses were performed: 1) We repeated the above-mentioned generalized linear model for sensitivity analysis of smokers classified by pack-years. 2) We excluded participants with diabetes mellitus, which is associated with a high baPWV and is a potential risk factor for arterial stiffness. 3) We also excluded participants with hypertension, which is associated with a high baPWV and is a potential risk factor for arterial stiffness. 4) To reduce the effect of the timing of post-exercise measurements, we also conducted another sensitivity analysis by excluding participants with the timing of post-exercise measurements exceeding 95% percent (26.7 minutes).

A P value < 0.05 was considered statistically significant with a two-sided test. SAS 9.4 (Version 9.4; SAS Institute, Cary, NC) was used for data analysis.

Results

Baseline characteristics of the study population

The study population had a mean age of 36.82 ± 7.76 years (► **Table 1**). On the basis of smoking status, the study population was divided into non-smokers ($n = 231$), former smokers ($n = 165$), light smokers ($n = 254$), and heavy smokers ($n = 290$). Participants with heavy smoke had higher pre-exercise SBP, pre-exercise DBP, pre-exercise MAP, pre-exercise baPWV, TG level, and rate of alcohol consumption, lower education level than those in the other three groups (all $P < 0.05$). The heavy smoker group had higher rate of taking exercise than those in the other three groups, but there were no statistical difference among groups ($P = 0.64$). After the cycling exercise, baPWV of all participants significantly decreased (1375.08 ± 209.09 cm/s vs. 1341.53 ± 208.04 cm/s, $P < 0.05$), and their HR increased (80.24 ± 12.54 bpm vs. 85.73 ± 13.56 bpm, $P < 0.05$). There were no significant changes in SBP, DBP, or MAP before and after exercise (► **Table 1**).

Comparison of changes in BP, HR, and baPWV between the groups

In adjusted analyses, the changes in baPWV of all participants post-exercise were -33.55 cm/s (95% CI, -39.69 to -27.42), and the change in the heavy smoking group was more significantly than other three groups. The changes in HR of all participants were 5.59 bpm (95% CI, 4.94 to 6.24). The changes in MAP of all participants post-exercise were 0.33 mmHg (95% CI, -0.19 to 0.85) (► **Table 2**).

Generalized linear model estimation of between-group differences in the change in baPWV

We used the generalized linear model to compare between-group differences in the change in baPWV. In Model 3, with the non-smokers as the control, the β -values and 95% CI of former smokers, light smoking smokers, and heavy smoking smokers were -12.17 cm/s (95% CI, -30.08 to 5.75), -18.43 cm/s (95% CI, -34.69 to -2.16), and -22.46 cm/s (95% CI, -38.39 to -6.54), respectively (► **Table 3**). For every increase in one standard deviation (6.26 cigarettes/

day) in the number of cigarettes smoked, the β -value and 95% CI were -18.11 cm/s (95% CI, -114.16 to 77.95) (► **Table 3**).

Stratified analysis

In the stratified analysis, after adjustment for covariates, the result was consistent with the main analysis among participants without regular exercise habit. But for those with regular exercise habit, there were no significant difference in the change of baPWV in former smoker, light smoker, and heavy smoker, compared with non-smokers (**Table S1**).

Sensitivity analysis

The results of the sensitivity analysis were consistent with those of the main analysis, which indicated that the main results were robust (**Table S2**).

Discussion

Using a functional community population, this study showed that aerobic exercise immediately improved arterial stiffness in individuals with different smoking statuses. Remarkably, the beneficial effects of exercise on arterial stiffness appeared to be more evident in light and heavy smoker than in non-smoker.

Following a 7-minute bout of exercise, we observed a significant reduction in baPWV of 33.55 cm/s from resting values in young and middle-aged (36.82 ± 7.76 years) subjects. This finding is similar to that by Yamato et al. (2016) [34]. They found that baPWV in non-smoking healthy men ($n = 26$, age: 21.0 ± 1.0 years) was decreased by 33.5 cm/s after 15 minutes of aerobic exercise for 40 minutes. However, Doonan et al.'s (2011) [16] study of 24 healthy young smoking men and 53 non-smokers showed that carotid-femoral pulse wave velocity in smokers and non-smokers increased by 80 cm/s and 60 cm/s, respectively, after 5 minutes of exhausting exercise. There are several possible reasons for this discrepancy between our study and Doonan et al.'s (2011) study. The sample size of Doonan et al.'s study was small, the population was relatively young (26.06 ± 6.70 years), and the exercise style was more intense compared with our study. Additionally, the difference in the type of pulse wave velocity could be the reason for the inconsistent conclusion between the studies. A meta-analysis suggested that aerobic exercise had different effects on arterial stiffness of different segments. In this meta-analysis, after aerobic exercise, arterial stiffness in the peripheral segments was significantly improved, but it did not significantly improve arterial stiffness in the central segment [35].

Our study showed that there was a difference in the decrease in baPWV after aerobic exercise in individuals with different smoking statuses. After adjusting for related confounding factors, baPWV was more significantly immediately decreased in low-intensity and high-intensity smokers than in never smokers (by 18.43 and 22.46 cm/s, respectively). A more likely explanation is that short-term aerobic exercise may have ameliorated the adverse effects of oxidative chemicals contained in cigarette smoke on vascular endothelial function and sympathetic nervous system activity [36]. Oxidative chemicals in cigarette smoke can affect vascular endothelial function through inflammatory response [37] and oxidative stress [38], which reduce the activity of endothelial nitric oxide syn-

► **Table 1** Baseline participant characteristics.

	All n=940	Never n=231	Former n=165	Light Smoker n=254	Heavy Smoker n=290	P
Demographic factors						
Age (year)	36.82 ± 7.76	37.18 ± 8.11	39.20 ± 6.97	34.36 ± 7.70	37.50 ± 7.64	<0.01
BMI (kg/m ²)	26.25 ± 3.89	26.45 ± 3.61	26.28 ± 3.50	25.82 ± 4.15	26.45 ± 4.05	0.14
Laboratory parameters						
TG (mmol/L)	1.20(0.83,1.91)	1.15(0.77,1.88)	1.19(0.79,1.78)	1.09(0.84,1.72)	1.35(0.90,2.12)	<0.01
FBG (mmol/L)	5.51 ± 0.94	5.54 ± 0.76	5.52 ± 0.81	5.47 ± 0.79	5.51 ± 1.22	0.89
Current drinker, n (%)	523(55.64)	108(46.75)	83(50.30)	144(56.69)	188(64.83)	<0.01
Physical activity, n (%)	352(37.45)	79(34.20)	63(38.18)	96(37.80)	114(39.31)	0.64
Senior college or above, n (%)	184(19.57)	62(26.84)	36(21.82)	51(20.08)	35(12.07)	<0.01
Diabetes mellitus, n (%)	66(7.02)	20(8.66)	14(8.48)	15(5.91)	17(5.86)	0.46
Hypertension, n (%)	319(33.94)	80(34.63)	50(30.30)	86(33.86)	103(35.52)	0.22
Arterial stiffness and hemodynamics (measured supine)						
Pre-exercise SBP (mmHg)	131.84 ± 15.97	132.38 ± 17.61	130.16 ± 15.46	129.99 ± 14.81	133.99 ± 15.66	0.01
Post-exercise SBP (mmHg)	132.34 ± 14.83	133.09 ± 14.45	131.20 ± 15.18	130.88 ± 14.57	133.67 ± 15.05	0.10
Pre-exercise DBP (mmHg)	78.99 ± 11.27	78.65 ± 12.04	78.87 ± 11.13	77.27 ± 11.51	80.84 ± 10.26	<0.01
Post-exercise DBP (mmHg)	79.35 ± 10.92	79.79 ± 10.75*	79.34 ± 11.12	77.51 ± 11.43	80.63 ± 10.30	<0.01
Pre-exercise MAP (mmHg)	99.09 ± 12.91	98.35 ± 13.88	98.28 ± 12.84	97.54 ± 12.32	101.50 ± 12.38	<0.01
Post-exercise MAP (mmHg)	99.42 ± 11.95	99.74 ± 11.76*	99.24 ± 12.45	97.71 ± 11.67	100.77 ± 11.94	0.03
Pre-exercise HR (bpm)	80.24 ± 12.54	79.04 ± 13.97	77.85 ± 12.54	81.82 ± 12.51	81.18 ± 11.03	<0.01
Post-exercise HR (bpm)	85.73 ± 13.56*	84.13 ± 14.46*	82.67 ± 13.59*	87.23 ± 13.71*	86.64 ± 12.40*	<0.01
Pre-exercise baPWV (cm/s)	1375.08 ± 209.09	1384.32 ± 220.35	1368.85 ± 218.41	1345.80 ± 194.43	1396.92 ± 204.75	0.03
Post-exercise baPWV (cm/s)	1341.53 ± 208.04*	1361.32 ± 216.35*	1341.83 ± 221.97*	1305.55 ± 195.90*	1357.10 ± 200.32*	<0.01
VO ₂ max (ml/kg/min)	3.19 ± 0.58	3.27 ± 0.68	3.18 ± 0.54	3.16 ± 0.52	3.17 ± 0.57	0.58
Peak heat rate (bpm)	139.39 ± 14.47	138.42 ± 13.90	137.33 ± 13.95	142.19 ± 14.60	138.88 ± 14.80	<0.01
Time interval (min)	7.13(4.93,11.16)	6.98(5.13,10.43)	7.57(4.95,12.60)	6.89(5.00,11.05)	7.02(4.73,11.15)	0.42
CVbaPWV	4.00(1.57,5.53)	3.95(1.58,5.55)	3.81(1.24,5.23)	3.77(1.30,5.27)	4.35(2.00,5.80)	0.04
CVSBP	3.90(1.40,5.43)	4.13(1.55,4.98)	3.74(1.55,5.46)	3.92(1.29,5.66)	3.80(1.26,5.42)	0.94
CVDBP	4.40(1.49,5.85)	4.61(1.45,6.01)	4.39(1.31,5.64)	4.44(1.44,5.64)	4.22(1.63,5.91)	0.93
CVMAP	4.24(1.45,6.07)	4.89(1.70,6.39)	4.02(1.42,5.22)	4.08(1.49,6.03)	3.99(1.19,6.20)	0.50
CVHR	6.57(2.49,9.10)	6.98(2.74,9.64)	6.49(1.96,9.98)	7.04(2.64,9.82)	5.87(2.45,8.22)	0.13
Note: BMI, body mass index; TG, triglyceride; FBG, fasting blood glucose; SBP, systolic blood pressure; DBP, diastolic blood pressure; MAP, mean arterial pressure; HR, heart rate; baPWV, brachial-ankle pulse wave velocity; VO ₂ max: maximum oxygen consumption; Time interval: the timing of post-exercise measurements; CV: coefficients of variation; *P<0.05 against pre-exercise value.						

► **Table 2** Adjusted values of changes in arterial stiffness and hemodynamics.

	All n=940	Never n=231	Former n=165	Light Smoker n=254	Heavy Smoker n=290
ΔMAP (mmHg)	0.33 (-0.19 to 0.85)	1.15 (0.28 to 2.01)	0.50 (-0.54 to 1.53)	0.20 (-0.64 to 1.04)	-0.20 (-0.98 to 0.57)
ΔHR (bpm)	5.59 (4.94 to 6.24)	5.10 (4.96 to 5.24)	4.79 (4.52 to 5.06)	5.36 (5.11 to 5.61)	5.50 (5.38 to 5.62)
Δ baPWV (cm/s)	-33.55 (-39.69 to -27.42)	-23.53 (-35.89 to -11.16)	-29.46 (-44.20 to -14.71)	-37.21 (-49.18 to -25.24)	-40.67 (-51.72 to -29.63)
Note: Adjusted means (95% CI) are presented; Adjusted for Age, the pre-exercise value BMI, and TG level; VO ₂ mx; Education level; Alcohol consumption; Physical activity; and Time interval; Abbreviation: MAP, mean arterial pressure; HR, heart rate; baPWV, brachial-ankle pulse wave velocity; VO ₂ max: Maximum oxygen consumption; Time interval: the timing of post-exercise measurements; ΔMAP/ΔHR/ΔbaPWV, post-exercise value minus pre-exercise value.					

thase and the production of nitric oxide [39]. Because aerobic exercise can stimulate the expression of antioxidant enzymes [40] and vasodilatory substances [41]. We speculate that the augmented antioxidant defense mechanism in current smokers is the reason why baPWV improvement is more significant than that in non-smokers after exercise.

Although we found that there was a difference in the decrease in baPWV after aerobic exercise in individuals with different smoking statuses, and this decrease in heavy smokers was more obvious, we did not examine the specific mechanism(s). Previous studies have shown that arterial stiffness is mainly determined by two factors. One factor is that the structure of arteries affects arterial

► **Table 3** Generalized linear model analysis results of the between-group difference of Δ baPWV with different smoking statuses.

	Smoking status	N	β	95% CI
Model 1	Never	231	Ref	
	Former	165	- 11.32	- 29.72 to 7.08
	Light smoker, ≤ 10 cigarettes/day	254	- 15.80	- 32.27 to - 0.68
	Heavy smoker, > 10 cigarettes/day	290	- 15.77	- 31.61 to - 0.07
	P for trend	-	0.05	
	Daily cigarette consumption, per SD (6.26)	-	6.99	- 118.27 to 75.59
Model 2	Never	231	Ref	
	Former	165	- 13.67	- 30.73 to 4.40
	Light smoker, ≤ 10 cigarettes/day	254	- 20.76	- 37.17 to - 4.36
	Heavy smoker, > 10 cigarettes/day	290	- 24.80	- 40.86 to - 8.73
	P for trend	-	< 0.01	
	Daily cigarette consumption, per SD (6.26)	-	- 21.34	- 118.27 to 75.59
Model 3	Never	231	Ref	
	Former	165	- 12.17	- 30.08 to 5.75
	Light Smoker, ≤ 10 cigarettes/day	254	- 18.43	- 34.69 to - 2.16
	Heavy Smoker, > 10 cigarettes/day	290	- 22.46	- 38.39 to - 6.54
	P for trend	-	< 0.01	
	Daily Cigarette Consumption, per SD (6.26)	-	- 18.11	- 114.16 to 77.95

Model 1: included Age, the pre-exercise values of baPWV; Model 2: included Model 1 covariates plus BMI, the pre-exercise values of HR, MAP, and TG level; VO2max, Time interval; Alcohol consumption; Education level; Model 3: included Model 2 covariates plus Δ HR; Δ MAP; Abbreviation: MAP, mean arterial pressure; HR, heart rate; baPWV, brachial-ankle pulse wave velocity, BMI, body mass index; TG, triglyceride; VO2max: maximum oxygen consumption; Time interval: the timing of post-exercise measurements; Δ HR/ Δ MAP, post-exercise value minus pre-exercise value.

stiffness. A decrease in elastin in the middle layer of the arterial wall and an increase in fibrin cause changes in vascular structure, which increases arterial stiffness [42–45]. The other factor is that the functional status of arteries also affects arterial stiffness. Vascular endothelial dysfunction and nervous system activity disorders can affect vascular function, resulting in increased arterial stiffness [46, 47]. Short-term aerobic exercise increases blood flow by stimulating the production of endothelial nitric oxide, prostaglandin, and other vasodilator factors, and decreases vasoconstrictors (e. g., endothelin) [41, 48–50], or vascular modifications [51]. These processes can improve vascular endothelial function and arterial stiffness. In addition, an improvement in arterial stiffness after aerobic exercise may also be related to changes in the levels of circulatory metabolites. The Framingham study showed a significant reduction in circulating blood metabolites after aerobic exercise, which resulted in insulin resistance, body inflammation, and oxidative stress. Metabolites related to cardio-cerebral vascular protection were significantly increased [52].

Improving lifestyles and preventing and controlling multiple risk factors are still primary prevention measures for cardiovascular disease, which can effectively reduce the probability of adverse cardiovascular events [53]. Hamdy et al. (2005) found that long-term regular aerobic exercise delayed and reversed the process of vascular aging [54]. Our study showed that arterial stiffness in individuals with different smoking statuses were significantly immediately decreased after exercise compared with before exercise. However, this might not be a continuous change. If the exercise time is prolonged and the exercise frequency is increased, arterial stiffness may be maintained. Therefore, smokers can adopt a healthy life-

style such as aerobic exercise to improve arterial stiffness, delay vascular aging, and reduce the damage to blood vessels caused by smoking.

This study has some limitations: (1) This was a cross-sectional study that was performed to examine the causal relationships between various factors and the difference in baPWV before and after exercise. (2) We did not find a dose–response relationship of the change in baPWV after aerobic exercise, possibly because we had difficulty in determining the specific number of cigarettes smoked per day. The number of cigarettes smoked was treated as a classification variable (1–10, 11–20, 21–30, 31–40, and > 40 cigarettes/day). Therefore, it is difficult to observe a dose–response relationship. (3) We used baPWV as a measurement of arterial stiffness instead of carotid–femoral pulse wave velocity, which is the gold standard. The validity and accuracy of baPWV against carotid–femoral pulse wave velocity have been demonstrated in previous studies [55], and the American Heart Association has also recommended baPWV as a common indicator for arterial stiffness [56]. (4) The study was conducted only in men, and findings could not be readily generalizable to female working populations or to workers in other settings or other countries. (5) Because of space and personnel constraints, the timing of baPWV measurement after aerobic exercise could not be unified. But baPWV was measured immediately (or as soon as possible) after aerobic exercise. In addition, when evaluating the short-term effect of acute aerobic exercise on arterial stiffness, we adjusted for the timing of baPWV measurement.

In summary, this study shows that a single aerobic exercise can immediately decrease baPWV in individuals with different smok-

ing statuses. Additionally, compared with non-smokers, baPWV decreased more significantly in light and heavy smokers.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- [1] Mitchell GF, Hwang SJ, Vasan RS et al. Arterial stiffness and cardiovascular events: the Framingham Heart Study. *Circulation* 2010; 121: 505–511
- [2] Safar ME, London GM, Plante GE. Arterial stiffness and kidney function. *Hypertension* 2004; 43: 163–168
- [3] Tsao CW, Seshadri S, Beiser AS et al. Relations of arterial stiffness and endothelial function to brain aging in the community. *Neurology* 2013; 81: 984–991
- [4] Sharif S, Visseren FLJ, Spiering W et al. Arterial stiffness as a risk factor for cardiovascular events and all-cause mortality in people with Type 2 diabetes. *Diabet Med* 2019; 36: 1125–1132
- [5] Said MA, Eppinga RN, Lipsic E et al. Relationship of arterial stiffness index and pulse pressure with cardiovascular disease and mortality. *J Am Heart Assoc* 2018; 7: e007621. doi: 10.1161/JAHA.117.007621
- [6] Cavalcante JL, Lima JA, Redheuil A et al. Aortic stiffness: current understanding and future directions. *J Am Coll Cardiol* 2011; 57: 1511–1522
- [7] Fu S, Chen W, Luo L et al. Roles of fasting and postprandial blood glucose in the effect of type 2 diabetes on central arterial stiffness: a 5-year prospective community-based analysis. *Diabetol Metab Syndr* 2017; 9: 33
- [8] Vlachopoulos C, Aznaouridis K, Stefanadis C. Prediction of cardiovascular events and all-cause mortality with arterial stiffness: a systematic review and meta-analysis. *J Am Coll Cardiol* 2010; 55: 1318–1327
- [9] Camplain R, Meyer ML, Tanaka H et al. Smoking behaviors and arterial stiffness measured by pulse wave velocity in older adults: the Atherosclerosis Risk in Communities (ARIC) study. *Am J Hypertens* 2016; 29: 1268–1275
- [10] Kubozono T, Miyata M, Ueyama K et al. Acute and chronic effects of smoking on arterial stiffness. *Circ J* 2011; 75: 698–702
- [11] Tomiyama H, Hashimoto H, Tanaka H et al. Continuous smoking and progression of arterial stiffening: a prospective study. *J Am Coll Cardiol* 2010; 55: 1979–1987
- [12] Lloyd-Jones DM, Hong Y, Labarthe D et al. Defining and setting national goals for cardiovascular health promotion and disease reduction: the American Heart Association's strategic Impact Goal through 2020 and beyond. *Circulation* 2010; 121: 586–613
- [13] Vaitkevicius PV, Fleg JL, Engel JH et al. Effects of age and aerobic capacity on arterial stiffness in healthy adults. *Circulation* 1993; 88: 1456–1462
- [14] Son WM, Sung KD, Cho JM et al. Combined exercise reduces arterial stiffness, blood pressure, and blood markers for cardiovascular risk in postmenopausal women with hypertension. *Menopause* 2017; 24: 262–268
- [15] Hamdy O, Ledbury S, Mullooly C et al. Lifestyle modification improves endothelial function in obese subjects with the insulin resistance syndrome. *Diabetes Care* 2003; 26: 2119–2125
- [16] Doonan RJ, Scheffler P, Yu A et al. Altered arterial stiffness and subendocardial viability ratio in young healthy light smokers after acute exercise. *PLoS One* 2011; 6: e26151
- [17] Naka KK, Tweddel AC, Parthimos D et al. Arterial distensibility: acute changes following dynamic exercise in normal subjects. *Am J Physiol Heart Circ Physiol* 2003; 284: H970–H978
- [18] Jin C, Chen S, Vaidya A et al. Longitudinal change in fasting blood glucose and myocardial infarction risk in a population without diabetes. *Diabetes Care* 2017; 40: 1565–1572
- [19] Wang XX, Chen GZ, Huang ZG et al. Short-Term Effect of Acute Aerobic Exercise On Arterial Stiffness in People with Different Smoking Statuses (self-citations). *research square* 2021
- [20] The State Council. the State Sports General Administration on the fifth National Physique Monitoring notice. Accessed on Feb 28, 2022 http://www.gov.cn/xinwen/2019-02/02/content_5363384.htm
- [21] The State Council. Labor Law of the People's Republic of China 2018. Accessed on Feb 28, 2022 http://english.www.gov.cn/archive/laws_regulations/2014/08/23/content_281474983042473.htm
- [22] Fletcher GF, Ades PA, Kligfield P et al. Exercise standards for testing and training: a scientific statement from the American Heart Association. *Circulation* 2013; 128: 873–934
- [23] Terry JW, Tolson H, Johnson DJ et al. A workload selection procedure for the Astrand-Rhyming test. *J Sports Med Phys Fitness* 1977; 17: 361–366
- [24] Munakata M. Brachial-ankle pulse wave velocity in the measurement of arterial stiffness: recent evidence and clinical applications. *Curr Hypertens Rev* 2014; 10: 49–57
- [25] Yamashina A, Tomiyama H, Takeda K et al. Validity, reproducibility, and clinical significance of noninvasive brachial-ankle pulse wave velocity measurement. *Hypertens Res* 2002; 25: 359–364
- [26] Eng L, Su J, Qiu X et al. Second-hand smoke as a predictor of smoking cessation among lung cancer survivors. *J Clin Oncol* 2014; 32: 564–570
- [27] Jatoti NA, Jerrard-Dunne P, Feely J et al. Impact of smoking and smoking cessation on arterial stiffness and aortic wave reflection in hypertension. *Hypertension* 2007; 49: 981–985
- [28] Kianoush S, Yakoob MY, Al-Rifai M et al. Associations of cigarette smoking with subclinical inflammation and atherosclerosis: elsa-brasil (the brazilian longitudinal study of adult health). *J Am Heart Assoc* 2017; 6: e005088. doi: 10.1161/JAHA.116.005088
- [29] Biener L, Albers AB. Young adults: vulnerable new targets of tobacco marketing. *Am J Public Health* 2004; 94: 326–330
- [30] Celermajer DS, Sorensen KE, Georgakopoulos D et al. Cigarette smoking is associated with dose-related and potentially reversible impairment of endothelium-dependent dilation in healthy young adults. *Circulation* 1993; 88: 2149–2155
- [31] Zhao M, Song L, Sun L et al. Associations of type 2 diabetes onset age with cardiovascular disease and mortality: the kailuan study. *Diabetes Care* 2021; 44: 1426–1432
- [32] deRuiter W, Faulkner G. Tobacco harm reduction strategies: the case for physical activity. *Nicotine Tob Res* 2006; 8: 157–168
- [33] Vaidya A, Cui L, Sun L et al. A prospective study of impaired fasting glucose and type 2 diabetes in China: The Kailuan study. *Medicine (Baltimore)* 2016; 95: e5350
- [34] Yamato Y, Hasegawa N, Sato K et al. Acute effect of static stretching exercise on arterial stiffness in healthy young adults. *Am J Phys Med Rehabil* 2016; 95: 764–770
- [35] Saz-Lara A, Cervero-Redondo I, Alvarez-Bueno C et al. The acute effect of exercise on arterial stiffness in healthy subjects: a meta-analysis. *J Clin Med* 2021; 10: 291
- [36] Eleutério-Silva MA, Sá da Fonseca LJ, Velloso EP et al. Short-term cardiovascular physical programme ameliorates arterial stiffness and decreases oxidative stress in women with metabolic syndrome. *J Rehabil Med* 2013; 45: 572–579

- [37] Barua RS, Ambrose JA, Eales-Reynolds LJ et al. Dysfunctional endothelial nitric oxide biosynthesis in healthy smokers with impaired endothelium-dependent vasodilatation. *Circulation* 2001; 104: 1905–1910
- [38] Kamceva G, Arsova-Sarafinovska Z, Ruskovska T et al. Cigarette smoking and oxidative stress in patients with coronary artery disease. *Open Access Maced J Med Sci* 2016; 4: 636–640
- [39] Siasos G, Tsigkou V, Kokkou E et al. Smoking and atherosclerosis: mechanisms of disease and new therapeutic approaches. *Curr Med Chem* 2014; 21: 3936–3948
- [40] Ji LL. Exercise-induced modulation of antioxidant defense. *Ann N Y Acad Sci* 2002; 959: 82–92
- [41] Rådegran G, Calbet JA. Role of adenosine in exercise-induced human skeletal muscle vasodilatation. *Acta Physiol Scand* 2001; 171: 177–185
- [42] McEniery CM, McDonnell BJ, So A et al. Aortic calcification is associated with aortic stiffness and isolated systolic hypertension in healthy individuals. *Hypertension* 2009; 53: 524–531
- [43] O'Rourke MF, Nichols WW. Aortic diameter, aortic stiffness, and wave reflection increase with age and isolated systolic hypertension. *Hypertension* 2005; 45: 652–658
- [44] McEniery CM, Yasmin, Maki-Petaja KM et al. The impact of cardiovascular risk factors on aortic stiffness and wave reflections depends on age: the Anglo-Cardiff Collaborative Trial (ACCT III). *Hypertension* 2010; 56: 591–597
- [45] Leone A. Interactive effect of combined exposure to active and passive smoking on cardiovascular system. *Recent Pat Cardiovasc Drug Discov* 2011; 6: 61–69
- [46] Lee DH, Ihm SH, Youn HJ et al. Age is an independent risk factor for the early morning blood pressure surge in patients never-treated for hypertension. *Korean Circ J* 2009; 39: 322–327
- [47] Chia S, Wilson R, Ludlam CA et al. Endothelial dysfunction in patients with recent myocardial infarction and hyperhomocysteinaemia: effects of vitamin supplementation. *Clin Sci* 2005; 108: 65–72
- [48] Tinken TM, Thijssen DH, Hopkins N et al. Shear stress mediates endothelial adaptations to exercise training in humans. *Hypertension* 2010; 55: 312–318
- [49] Sarelius I, Pohl U. Control of muscle blood flow during exercise: local factors and integrative mechanisms. *Acta Physiol (Oxf)* 2010; 199: 349–365
- [50] Wray DW, Nishiyama SK, Donato AJ et al. Endothelin-1-mediated vasoconstriction at rest and during dynamic exercise in healthy humans. *Am J Physiol Heart Circ Physiol* 2007; 293: H2550–H2556
- [51] Siasos G, Athanasiou D, Terzis G et al. Acute effects of different types of aerobic exercise on endothelial function and arterial stiffness. *Eur J Prev Cardiol* 2016; 23: 1565–1572
- [52] Nayor M, Shah RV, Miller PE et al. Metabolic architecture of acute exercise response in middle-aged adults in the community. *Circulation* 2020; 142: 1905–1924
- [53] Arnett DK, Blumenthal RS, Albert MA et al. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: a report of the american college of cardiology/american heart association task force on clinical practice guidelines. *Circulation* 2019; 140: e596–e646
- [54] Hamdy O. Lifestyle modification and endothelial function in obese subjects. *Expert Rev Cardiovasc Ther* 2005; 3: 231–241
- [55] Tanaka H, Munakata M, Kawano Y et al. Comparison between carotid-femoral and brachial-ankle pulse wave velocity as measures of arterial stiffness. *J Hypertens* 2009; 27: 2022–2027
- [56] Townsend RR, Wilkinson IB, Schiffrin EL et al. Recommendations for improving and standardizing vascular research on arterial stiffness: a scientific statement from the american heart association. *Hypertension* 2015; 66: 698–722