#### E-Videos

# Unique double guidewire technique using a guidewire perforating the diverticulum for a difficult biliary cannulation





► **Fig.1** Computed tomography showing a common bile duct stone.



► Fig.2 Endoscopic image: The orifice of the papilla could be identified after cleaning food remnants in the diverticulum and insertion of the tip of the side-viewing endoscope into the diverticulum.

Biliary cannulation of an intradiverticular papilla, classified type 1 as per the Li-Tanaka classification, is sometimes challenging [1]. Although in cases with difficult biliary cannulation the pancreatic duct guidewire placement technique is useful [2–3], placing the guidewire in the pancreatic duct is sometimes difficult. We report a successful case of biliary cannulation in a patient with an intradiverticular papilla using an unusual double guidewire technique taking advantage of a guidewire perforating the diverticulum. An 80-year-old woman presented to another hospital with high fever and abdominal pain. She was diagnosed with



**Fig.3** The guidewire perforating through the papilla into the diverticulum during several attempts of wire-guided cannulation.

cholangitis due to common bile duct stones (> Fig. 1). Three attempts of endoscopic retrograde cholangiopancreatography (ERCP) failed to identify the papilla. Therefore, she was referred to our hospital. Identification of the papilla was difficult; the orifice of the papilla was identified after cleaning food remnants in the diverticulum and inserting the tip of the side-viewing endoscope (TJF-Q290V; Olympus Medical Systems, Tokyo, Japan) into the diverticulum (**Fig.2**). Contrast medium injection was possible, and the bile duct stones and pancreatic duct were visualized. However, guidewire insertion into the bile and pancreatic ducts was challenging. During several attempts of wireguided cannulation, the guidewire perforated the diverticulum through the papilla (> Fig. 3). We attempted to perform biliary cannulation using the double guidewire technique with the guidewire perforating the diverticulum instead of placing the guidewire in the pancreatic duct to help stabilize and visualize the orifice of the papilla. Wire-guided cannulation of the bile duct was successful (> Fig. 4), and the stone was removed using a basket and balloon catheter



► Fig. 4 Fluoroscopic image: Using the double guidewire technique with the guidewire perforating the diverticulum, the wire-guided cannulation of the bile duct was successful.

following endoscopic balloon dilation (> Video 1).

The guidewire sometimes perforates the papilla during ERCP; however, most cases improve conservatively [4–5]. Therefore, instead of removing the guidewire that perforated the diverticulum, we used it for the double guidewire method to cannulate the bile duct.



**Video 1** Biliary cannulation was successfully performed using a unique double guidewire technique with a guidewire that had perforated the diverticulum.

## Endoscopy\_UCTN\_Code\_TTT\_1AR\_2AH

## **Competing interests**

The authors declare that they have no conflict of interest.

## The authors

Kazuya Koizumi 🧠 Kento Shionoya 🔍 Sakue Masuda 🧐 Jun Kubota, Karen Kimura Gastroenterology Medicine Center, Shonan Kamakura General Hospital, Kamakura, Kanagawa, Japan

### Corresponding author

#### Kazuya Koizumi, MD, PhD

Gastroenterology Medicine Center, Shonan Kamakura General Hospital, 1370-1 Okamoto, Kamakura, Kanagawa 247-8533, |apan Fax: +81-467-45-0190 kizm2010@gmail.com

### References

- [1] Yue P, Zhu KX, Wang HP et al. Clinical significance of different periampullary diverticulum classifications for endoscopic retrograde cholangiopancreatography cannulation. World | Gastroenterol 2020; 26: 2403-2415
- [2] Dumonceau JM, Devière J, Cremer M. A new method of achieving deep cannulation of the common bile duct during endoscopic retrograde cholangiopancreatography. Endoscopy 1998; 30: S80
- [3] Gyökeres T, Duhl J, Varsányi M et al. Double guide wire placement for endoscopic pancreaticobiliary procedures. Endoscopy 2003; 35: 95-96
- [4] Kawakami H, Maguchi H, Mukai T et al. A multicenter, prospective, randomized study of selective bile duct cannulation performed by multiple endoscopists: the BIDMEN study. Gastrointest Endosc 2012; 75: 362-372
- [5] Takano Y, Nagahama M, Yamamura E et al. Perforation of the papilla of Vater in wireguided cannulation. Can J Gastroenterol Hepatol 2016; 2016: 5825230

## **Bibliography**

Endoscopy 2023; 55: E734-E735 DOI 10.1055/a-2086-2153 ISSN 0013-726X © 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



# **ENDOSCOPY E-VIDEOS** https://eref.thieme.de/e-videos



section of the journal Endoscopy, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high-guality video and are published with a Creative Commons CC-BY license. Endoscopy E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: https:// www.research4life.org/access/eligibility/).

E-Videos is an open access online

This section has its own submission website at https://mc.manuscriptcentral.com/e-videos