


# Copper Sulfate (CuSO<sub>4</sub>): An Efficient Reagent in Organic Synthesis

Houseberg D. S. Ferreira\* 

Euzébio G. Barbosa

Alessandro K. Jordão 

Universidade Federal do Rio Grande do Norte, Departamento de Farmácia, Programa de Pós-Graduação em Ciências Farmacêuticas, Rua General Gustavo Cordeiro de Faria S/N, Petrópolis, Centro de Ciências da Saúde, Natal, RN, CEP 59012-570, Brasil  
houseberg.donanvam@ufrn.edu.br

Received: 11.05.2023



Accepted after revision: 19.07.2023

Published online: 21.07.2023 (Accepted Manuscript),

15.08.2023 (Version of Record)

DOI: 10.1055/a-2134-9007; Art ID: SO-2023-05-0034-SPOT



License terms:  

© 2023. The Author(s). This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution and reproduction, so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)


**Key words** synthesis, copper sulfate, reagents, catalysis, organic chemistry

Copper sulfate (CuSO<sub>4</sub>) occurs in nature as anhydrous form (hydrocyanite) and as mono- and pentahydrate. It is a simple, inexpensive, and commercially available salt synthesized by the treatment of cupric oxide with sulfuric acid.<sup>1</sup>

As monohydrate, also known as dried cupric sulfate, is hygroscopic, off-white powder, soluble in water, and practically insoluble in alcohol (MeOH, EtOH). As pentahydrate, also known as blue vitriol, Salzburg vitriol among others is a large, blue or ultramarine, triclinic crystals or blue granules or light-blue powder. It can be obtained from the anhydrous form by dissolving in deionized water and allowing the solvent to evaporate at room temperature, forming blue crystals in approximately one week.<sup>2</sup> Loses 2 H<sub>2</sub>O at 30 °C becomes anhydrous by 250 °C. It is very soluble in water, soluble in methanol, glycerol, and slightly soluble in ethanol and shows acid characteristic with pH 0.2 molar.<sup>1</sup>

The anhydrous salt is used for detecting and removing trace amount of water from alcohols and other organic compounds. It is also used in agricultural as fungicide, algicide, bactericide, herbicide, and fertilizer additive.<sup>1</sup>

Besides, copper pentahydrate is an eye irritant but not a skin irritant, it may induce allergic dermatitis in sensitive people.<sup>3</sup> Copper is also an essential trace element and an important catalyst for heme synthesis and iron absorption on biological organisms. His dysregulation has been studied



**Houseberg Donanvam da Silva Ferreira** received his pharmacy degree from the Federal University of Rio Grande do Norte in 2020. Currently, he is an MSc pharmaceutical sciences student at the same institution under the supervision of Prof. A. K. Jordão and Prof. E. G. Barbosa. His work involves the synthesis and biological evaluation of hydrazones.

**Euzébio Guimarães Barbosa** received his PhD in chemistry from Campinas University (UNICAMP) in 2011 under the supervision of Prof. Marcia Miguel Castro Ferreira. Currently he is a professor at the Federal University of Rio Grande do Norte. His research interests focus on medicinal chemistry and computer-aided drug design.

**Alessandro Kappel Jordão** received his PhD in chemistry from the Fluminense Federal University (UFF) in 2010 under the supervision of Prof. Vitor Francisco Ferreira and Prof. Anna Claudia. Currently he is a professor at the Federal University of Rio Grande do Norte. His research interests focus on the synthesis of heterocyclic compounds.

with a focus on neurodegenerative diseases, such as Wilson's, Alzheimer, and Parkinson disease.<sup>4</sup>

In chemistry, this compound is normally used as a catalyst for reactions, due to its low cost, possibility of use at low temperatures and ecological advantages.<sup>5</sup> Several reactions have been reported using this compound since 1944 when Hann and Hudson<sup>6</sup> reported the activity of CuSO<sub>4</sub> as a catalyst in triazole formation reactions. Copper sulfate stands out for its application in catalysis reactions, for example, in the synthesis of aryl/vinyl halides and azides of vinyl/aryl boronate esters/boronic acids.<sup>7</sup> Table 1 presents a series of recent applications of this reagent.

**Table 1** Recent Applications of Cupric Sulfate (CuSO<sub>4</sub>)

<p>(A) Daniel Gonzaga <i>et al.</i> synthesized 2-phenyl-triazole-carbaldehyde using a three-step sequence for obtaining the Fisher osazone (D-glucose adduct with substituted phenylhydrazines) followed by oxidative cyclization by Hudson's method (refluxed in aqueous solution of CuSO<sub>4</sub>), generating the derivate phenyl-D-glucosotriazol, followed by the oxidative cleavage of the glycotriazole in aqueous NaIO<sub>4</sub> to obtain the carboxaldehyde.<sup>8</sup></p>	<p>1) D-glucose, H<sub>2</sub>O, reflux, sodium acetate, 2 h, 63%  2) CuSO<sub>4</sub> (3.52 mol%), rt, 2 h, 67%  3) NaIO<sub>4</sub>, rt, 24 h, &gt;98%  4) NaBH<sub>4</sub>, MeOH, rt, 24 h</p>
<p>(B) Elodie Decuypere <i>et al.</i> synthesized a series of 1,2,3-triazoles using the CuSO<sub>4</sub> as a copper salt catalyst in the copper-catalyzed aza-iminosydnone-alkyne cycloaddition reactions. It is added to the reaction mixture along with sodium ascorbate and one of the selected ligands to form the copper complex used in the reaction.<sup>9</sup></p>	<p>CuSO<sub>4</sub> (0.02 mol%)-Ligand A or E  Et<sub>3</sub>N  Na ascorbate  tBuOH/H<sub>2</sub>O  30 °C, overnight  29-78%</p> <p>R = <i>n</i>Pr, <i>i</i>Pr, <i>t</i>Bu, Ph, COPh  R' = (CH<sub>2</sub>)<sub>2</sub>Ph, CH<sub>2</sub>OH, (CH<sub>2</sub>)<sub>2</sub>OH, (CH<sub>2</sub>)<sub>4</sub>OH</p>
<p>(C) Mario Ficker <i>et al.</i> prepared an ester by reduction of alkene using a copper-cobalt catalytic system, resulting in a good product yield.<sup>10</sup></p>	<p>CuSO<sub>4</sub> (3 mol%)  CoCl<sub>2</sub>  NaBH<sub>4</sub>  H<sub>2</sub>O  rt, 10 min  91%</p>
<p>(D) Salhah D. Al-Qahtani <i>et al.</i> synthesized metallic complexes using EtOH/H<sub>2</sub>O mixtures with a concentration of 0.001 mol dm<sup>-3</sup> of copper sulfate for the preparation of the metallic binding solution with the proposed naphthohydrazone.<sup>11</sup></p>	<p>CuSO<sub>4</sub> (0.28 mol%)  EtOH-H<sub>2</sub>O  rt  58%</p>
<p>(E) Pankaj R Chaudhari <i>et al.</i> published a six-step synthesis for the preparation of a sutezolid compound where one of the intermediates is synthesized by the reaction involving copper sulfate catalysis for 12 h in ethanol over -0.5 °C in 79% yield.<sup>12</sup></p>	<p>sodium triacetoxyborohydride  DCM, CuSO<sub>4</sub> (0.003 mol%), MeOH  -0.5 °C, 12 h  79%</p>
<p>(F) David Vrbata <i>et al.</i> performed a coupling of reaction with azide precursor under microwave irradiation at 40 °C for 1 h with a binary mixture of <i>t</i>-BuOH/H<sub>2</sub>O (proportions related to the polarity of the azide) involving copper sulfate and sodium ascorbate.<sup>13</sup></p>	<p>CuSO<sub>4</sub> (8.4 mol%), sodium ascorbate  40 °C, 2 h, MW irradiation  23-36%</p> <p>R =     X = H, Ac, Br, NO<sub>2</sub>, CN, <i>t</i>Bu, F</p>
<p>(G) Huafang Fan <i>et al.</i> synthesized vinyl lactams using a three-step reaction involving oxidative cleavage of D-isoascorbic acid, followed by acetonide formation of the resulting D-erythro lactone catalyzed by copper sulfate and, per last, ring opening of the lactone with sodium azide in 39% yield.<sup>14</sup></p>	<p>1) H<sub>2</sub>O<sub>2</sub>, Na<sub>2</sub>CO<sub>3</sub>  2) Acetone, CuSO<sub>4</sub> (9.46 mol%)  3) NaN<sub>3</sub>, DMF  110 °C, 12 h  23%  39%</p>
<p>(H) Jeffrey R. Groch, Nicholas R. Lauta, and Jon T. Njardarson synthesized two chiral sulfonimines using the copper sulfate to drive imine formation in the final product with varying yields of 60–89%.<sup>15</sup></p>	<p>H<sub>2</sub>N-S(=O)-<i>t</i>Bu  CuSO<sub>4</sub> (4.5 mol%), DCM  rt, 24 h  60-89%  R = CH<sub>2</sub>CH<sub>3</sub>  2 examples</p>

<p>(I) Alex A. Hunt-Painter <i>et al.</i> proposed a synthetic route for the synthesis of an iminosugar using a method of protection of isopropylidene groups using acetone, sulfuric acid, and copper sulfate for a subsequent iodination with 78% yield over two steps.<sup>16</sup></p>	
<p>(J) Ankush Banerjee, Shuvendu Saha, and Modhu Sudan Maji proposed an economical approach to the use of biologically active marine alkaloids, where hyellazole and 6-chlorohyellazole were first synthesized and were subjected to a dimerization reaction in the presence of CuSO<sub>4</sub>/Al<sub>2</sub>O<sub>3</sub> under an O<sub>2</sub> atmosphere at 140 °C to produce the dimer known as sorazolone E2 in 67% yield.<sup>17</sup></p>	
<p>(K) Seki and Takahashi reported a catalytic system with copper sulfate for regioselective C–H azidation of methyl anthranilate. This reaction is performed using NaN<sub>3</sub> and Na<sub>2</sub>S<sub>2</sub>O<sub>8</sub> to provide the corresponding <math>\alpha</math>-azidated product in 80% yield.<sup>18</sup></p>	
<p>(L) Shan and co-workers published the synthesis of indole-fused tetracyclic heteroacene via CuSO<sub>4</sub>-catalyzed carbanion radical redox relay, yielding benzofuro[3,2-b]indole in 98%.<sup>19</sup></p>	

In summary, copper sulfate enables a variety of diverse reactions and functionalizations that include: triazole formation, reduction of alkenes and alkynes, complexes formation, and catalysis among others. The support reagent offers advantages for the reactions, and in the most it is used as catalyst due to all advantages such increase reagent stability, ease of workup, low cost, and ecological advantages.

## Conflict of Interest

The authors declare no conflict of interest.

## References

- (1) *The Merck Index, 11th ed*; Budavari, S., Ed.; Merck & Co: Rahway, **1989**.
- (2) Ruggiero, M. T.; Erba, A.; Orlando, R.; Kortner, T. M. *Phys. Chem. Chem. Phys.* **2015**, *17*, 31023.
- (3) Aquilina, G.; Bach, A.; Bampidis, V.; Bastos, M. L.; Flachowsky, G.; Gasa-Gasó, J.; Gralak, M.; Hogstrand, C.; Leng, L.; López-Puente, S.; Martelli, G.; Mayo, B.; Renshaw, D.; Rychen, D.; Saarela, M.; Sejrsen, K.; Beelen, P. V.; Wallace, R. J.; Westendorf, J. *EFSA J.* **2012**, *10*, 12.
- (4) Manto, M. *Toxics* **2014**, *2*, 327.
- (5) Güner, E. K.; Kancan, D.; Naktiyok, J.; Özer, A. *Asia-Pac. J. Chem. Eng.* **2021**, *16*, e2583.
- (6) Hann, R. M.; Hudson, C. S. *American Chemical Society* **1944**, *66*, 735.
- (7) Tao, C.-Z.; Cui, X.; Li, J.; Liu, A.-X.; Liu, L.; Guo, Q.-X. *Tetrahedron Lett.* **2007**, *48*, 3525.
- (8) Gonzaga, D.; Da Silva, F. C.; Ferreira, V. F.; Wardell, J. L.; Wardell, S. M. S. V. *J. Chem. Crystallogr.* **2016**, *46*, 67.
- (9) Decuyper, E.; Bernard, S.; Feng, M.; Porte, K.; Riomet, M.; Thuéry, P.; Audisio, D.; Taran, F. *ACS Catal.* **2018**, *8*, 11882.
- (10) Ficker, M.; Svenningsen, S. W.; Larribeau, T.; Christensen, J. B. *Tetrahedron Lett.* **2018**, *59*, 1125.
- (11) Al-Qahtani, S. D.; Alsoliemy, A.; Almeahadi, S. J.; Alkhamis, K.; Alrefaei, A. F.; Zaky, R.; El-Metwaly, N. *J. Mol. Struct.* **2021**, 1244.
- (12) Chaudhari, P. R.; Bhise, N. B.; Singh, G. P.; Bhat, V.; Shenoy, G. G. *J. Chem. Sci.* **2022**, *134*, 56.
- (13) Vrbata, D.; Filipová, M.; Tavares, M. R.; Cerveny, J.; Vlachova, M.; Sirova, M.; Pelantova, H.; Petraskova, L.; Bumba, L.; Konefał, R.; Etrych, T.; Kren, V.; Chytil, P.; Bojarová, P. *J. Med. Chem.* **2022**, *65*, 3866.
- (14) Fan, H.; Tong, Z.; Ren, Z.; Mishra, K.; Morita, S.; Edouarzin, E.; Gorla, L.; Averkiev, B.; Day, V. W.; Hua, D. H. *J. Org. Chem.* **2022**, *87*, 6742.
- (15) Groch, J. R.; Lauta, N. R.; Njardarson, J. T. *Org. Lett.* **2023**, *25*, 395.
- (16) Hunt-Painter, A. A.; Deeble, B. M.; Stocker, B. L.; Timmer, M. S. M. *ACS Omega* **2022**, *7*, 28756.
- (17) Banerjee, A.; Saha, S.; Maji, M. S. *J. Org. Chem.* **2022**, *87*, 4343.
- (18) Seki, M.; Takahashi, Y. *J. Org. Chem.* **2021**, *86*, 7842.
- (19) Shan, X.-H.; Yang, B.; Qu, J.-P.; Kang, Y.-B. *Chem. Commun.* **2020**, *56*, 4063.