



Trends of Microsurgical Head and Neck Free Flap Reconstruction and Safety during the COVID-19 Pandemic

Valeria P. Bustos, MD, MS, MPH¹ Nicholas Elmer, MD¹ Anamika Veeramani, MD¹ Carly D. Comer, MD¹
Jacquelyn R. Kinney, MD¹ Samuel M. Manstein, MD¹ Natalie Hassell, MD¹
Bernard T. Lee, MD, MBA, MPH, FACS¹ Samuel J. Lin, MD, MBA, FACS^{1,2}

¹ Division of Plastic Surgery, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts

² Division of Otolaryngology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts

Address for correspondence Samuel J. Lin, MD, MBA, Division of Plastic Surgery, Beth Israel Deaconess Medical Center, Harvard Medical School, 110 Francis St, Suite 5A, Boston, MA 02215 (e-mail: sjlin@bidmc.harvard.edu).

J Reconstr Microsurg Open 2024;9:e69–e77.

Abstract

Background The coronavirus disease 2019 (COVID-19) pandemic poses unprecedented challenges among patients with head and neck (HN) cancer that require oncological and reconstructive surgeries. This study aims to identify differences in postoperative outcomes for patients who underwent microsurgical HN free flap reconstruction prior to versus during the COVID-19 pandemic.

Methods A retrospective observational study using the American College of Surgeons National Surgical Quality Improvement Program 2019 to 2020 database to identify patients with HN cancer who underwent a vascularized free tissue transfer was undertaken. Two cohorts were created: pre- and during COVID-19. Fisher's exact test and the unpaired Student's *t*-test were used to evaluate differences in sociodemographic and clinical characteristics between the cohorts. Multivariable logistic regression was used to assess differences in reoperation rates between groups as well as to identify potential risk factors for reoperation.

Results A total of 763 patients were analyzed. The mean age of patients in the overall cohort was 63.6 (standard deviation: 11.5) years. Most patients were white (62.7%). Overall, no statistically significant difference was evidenced between cohorts in terms of immediate postoperative outcomes. Similarly, reoperation rates were similar between groups ($p > 0.05$). Dependent functional status ($p = 0.021$) and postoperative infection ($p < 0.001$) were found to be risk factors for reoperation after holding other factors constant.

Conclusion HN flap reconstruction can be performed safely during the COVID-19 era. Standardized protocols for patient selection must be strictly followed to avoid disease progression and optimize surgical outcomes. Further studies assessing long-term outcomes during the pandemic are of utmost importance to elucidate the true impact of the COVID-19 pandemic on this population.

Keywords

- ▶ reconstruction
- ▶ microsurgery
- ▶ head and neck
- ▶ COVID-19

received

November 24, 2023

accepted after revision

February 16, 2024

accepted manuscript online

March 14, 2024

DOI <https://doi.org/10.1055/a-2287-6606>.

ISSN 2377-0813.

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Medical Publishers, Inc., 333 Seventh Avenue, 18th Floor, New York, NY 10001, USA

Introduction

The coronavirus disease 2019 (COVID-19) pandemic has had far-ranging effects on every health care system in the world. Hospitals and clinics have been overwhelmed by the deluge of patient care needs directly resulting from COVID-19 infections, while many other patients have been unable to access other types of health care.¹ The toll of this pandemic, which includes widespread difficulties in providing health care for a myriad of reasons, has led to a need for resource stewardship. This has resulted in some patients not receiving timely and critical care, particularly patients with cancer.² Reconstructive surgery has been no exception to this strain, where temporary holds on elective surgery have delayed care for many patients.^{3,4}

Many surgeons have devised innovative safety protocols to help these patients receive necessary procedures.^{5–7} Studies have examined the effects of these altered protocols on patient course and complication rates, specifically in the context of microsurgical head and neck (HN) reconstruction.⁸ Preliminary center-specific studies have demonstrated that, with adjusted safety protocols and careful patient selection, there has been significant difference in complications when these measures to mitigate viral exposure are taken.^{8,9} To further these efforts and protect access to care for HN malignancy, specifically that of free tissue transfer reconstruction, there is a need to assess the efficacy and safety of these new protocols comprehensively.

Using data from the American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP) database, this study aims to identify any differences in postoperative outcomes for patients who underwent microsurgical HN free flap reconstruction prior to the COVID-19 pandemic compared with those who underwent reconstruction during the pandemic.

Methods

Patient Identification

After obtaining approval from our institution's IRB (Protocol# 2021D001052), we performed a retrospective study using the ACS-NSQIP database. All patients undergoing HN free flap reconstruction from the ACS-NSQIP 2019 to 2020 were prospectively collected and analyzed.^{10–12} CPT (Current Procedural Terminology) codes were first used to identify patients who underwent a vascularized free tissue transfer (► **Supplementary Table S1**, available in the online version). Identified cases were then cross-referenced with patients with an ICD-10 (International Classification of Diseases 10th Revision) code associated with a HN malignancy (► **Supplementary Table S2**, available in the online version).

Aims and Outcomes of Interest

The primary aim of this study was to compare differences in 30-day postoperative outcomes in patients who underwent HN free flap reconstruction during the COVID-19 pandemic (Q2–Q4 of 2020) compared with those who underwent HN free flap reconstruction in the year prior to the pandemic

(Q1–Q4 of 2019 and Q1 of 2020). Furthermore, we compared the percentage of unplanned reoperations for each day through the first postoperative month (postoperative day [POD]: 30) to provide a detailed postoperative timeline for the two groups in an effort to capture potential delays in care linked to the COVID-19 pandemic. We also aimed to analyze potential risk factors for reoperation.

We included HN free flap patients from the entire year of 2019 and the first quarter of 2020 as the prepandemic control in this study for several reasons. First, most institutional practices regarding surgical logistics, surgeon technique, and postoperative enhanced recovery pathways were likely most similar between these 2 years. Utilizing data from years prior to 2019 may have introduced important differences related to these factors, which could confound the data. Therefore, we surmised that comparing these two cohorts offers the best opportunity to elucidate the potential effects of the COVID-19 pandemic on patient selection, hospital course and severity, and postoperative outcomes.

Risk Adjustment/Statistical Analysis

Patient demographics and clinical characteristics were summarized. Rates of wound-related, pulmonary, cardiac, renal/genitourinary, hematologic, and systemic complications were compared between the 2 years. For descriptive analysis, frequencies and percentages were used to present categorical variables. Following the Central Limit Theorem, the data were considered normally distributed, and therefore, means and standard deviations (SDs) were used to present continuous variables.

For inferential analysis, the Fisher's exact test and the unpaired *t*-test were used to assess differences in sociodemographic, clinical characteristics, and complications between groups in categorical and continuous data, respectively. A multivariable logistic regression was used to assess differences in reoperation between groups as well as to identify potential risk factors for reoperation. Statistical significance was set up for a *p*-value less than 0.05. Analysis was performed using STATA statistical software, version 16.1 (STATA Corp., College Station, TX).

Results

Demographics: Prepandemic and Coronavirus Disease 2019 Cohorts

A total of 763 HN free flaps were analyzed between 2019 and 2020. The mean (SD) age of patients in the overall cohort was 63.6 (11.5) years, and the mean (SD) BMI was 27.0 (6.5). Among patients in the study, 62.7% of patients were White, 4.8% Black or African American, and 32.5% represented other races (Asian, American Indian, Alaska or Hawaiian native, Pacific Islander) or were unknown. Overall, 27.7% of patients were smokers at the time of surgery, and 79.6% had an ASA (American Society of Anesthesiologists) physical status classification between 3 and 5. Almost all (97.9%) of patients were of independent functional status. This study's most common patient comorbidity was hypertension (50.1%). When divided into their respective groups based on time

of surgery, there were significantly more patients who lost >10% body weight in the 6 months prior to surgery ($p = 0.056$) in the prepandemic group. Although not significant, there was a trend toward overall differences in the race among patients based on time of surgery ($p = 0.06$). A summary of the population demographics for both groups is presented in ►Table 1.

Surgical Characteristics

A significant difference was evidenced in HN flap reconstruction trends between the cohorts (►Fig. 1). The majority (98.9%) of patients had their surgery performed in the inpatient setting. “Clean” or “clean contaminated” wound classification was present in 95.2%. The length of hospital stay was 6.0 days for those who underwent HN free flap reconstruction prior to the pandemic and 5.6 days for those during the pandemic. When comparing the operative time

between the COVID-19 and prepandemic groups, the COVID-19 group had a shorter operative time of only 7.5 minutes, which was not statistically significant. When stratified by location of reconstruction in the total cohort, 61.1% of patients had reconstruction of the oral cavity, 20.8% had reconstruction of the pharynx, and the remaining 18.1% had reconstruction of the mandible (4.6%), maxilla (3.9%), and surface structures (9.6%). The surgical specialty performing HN free flap reconstruction cases was otolaryngology (75.4%), with plastic surgery performing 22.4%. Surgical characteristics for both groups can be found in ►Table 2.

Postoperative Complications

The most common postoperative complications for the cohort were hematologic complication and unplanned reoperation, with 23.1 and 17.6% during the first 30 PODs, respectively (►Table 3). The next most common

Table 1 Patient demographics based on time of operation

	Historical control N = 462	COVID-19 N = 301	Total cohort N = 763	p-Value
Age mean (SD)	63.6 (11.8)	63.6 (11.0)	63.6 (11.5)	0.9786
BMI mean (SD)	27.0 (6.4)	27.1 (6.7)	27.0 (6.5)	0.7374
Race, n (%)				
White	286 (62.6)	189 (62.8)	475 (62.7)	0.056
African American	22 (4.8)	14 (4.7)	36 (4.8)	
Asian	19 (4.2)	17 (5.7)	36 (4.8)	
American Indian	0 (0)	2 (0.7)	2 (0.3)	
Native Hawaiian or Other Pacific Island	0 (0)	4 (1.3)	4 (0.5)	
Other/unknown	130 (28.5)	75 (24.9)	205 (27.0)	
Functional status, n (%)				
Independent	451 (97.6)	296 (98.3)	747 (97.9)	0.879
Partially dependent	9 (2.0)	4 (1.3)	13 (1.7)	
Totally dependent	1 (0.2)	1 (0.3)	2 (0.3)	
Unknown	1 (0.2)	0 (0)	1 (0.1)	
ASA classification, n (%)				
1–2	95 (20.6)	61 (20.3)	156 (20.5)	1.000
3–5	367 (79.4)	240 (79.7)	607 (79.6)	
Current smoker, n (%)	126 (27.3)	85 (28.2)	211 (27.7)	0.804
Comorbidities, n (%)				
Diabetes mellitus	25 (5.4)	21 (7.0)	46 (6.0)	0.437
COPD	30 (6.5)	30 (10.0)	60 (7.9)	0.098
Hypertension	235 (50.9)	147 (48.8)	382 (50.1)	0.605
Disseminated cancer	30 (6.5)	10 (3.3)	40 (5.2)	0.0670
Preoperative steroids	16 (3.5)	18 (6.0)	34 (4.5)	0.108
Bleeding disorder	7 (1.5)	6 (2.0)	13 (1.7)	0.776
Preoperative weight loss	48 (10.4)	17 (5.7)	65 (8.5)	0.024 ^a

Abbreviations: ASA, American Society of Anesthesiologists; BMI, body mass index; COPD, chronic obstructive pulmonary disease; COVID-19, coronavirus disease 2019; n, frequency; SD, standard deviation.

^aStatistical significance was set up for a p-value less than 0.05.




Fig. 1 Number of head and neck free flap reconstructions by quarter from 2019 to 2020.

Table 2 Surgical characteristics

	Historical control N = 462	COVID-19 N = 301	Total cohort N = 763	p-Value
Surgical team				
Otolaryngology	345 (74.7)	230 (76.4)	575 (75.4)	0.676
Plastic surgery	105 (22.7)	66 (21.9)	171 (22.4)	
Other	12 (2.6)	5 (1.7)	17 (2.2)	
Location of reconstruction, n (%)				
Oral cavity	247 (62.1)	149 (59.6)	396 (61.1)	0.761
Pharynx	79 (19.9)	56 (22.4)	135 (20.8)	
Mandible	21 (5.3)	9 (3.6)	30 (4.6)	
Maxilla	15 (3.8)	10 (4.0)	25 (3.9)	
Surface structures	36 (9.1)	26 (10.4)	61 (9.6)	
Surgery setting, n (%)				
Inpatient	457 (98.9)	297 (98.7)	754 (98.8)	0.745
Outpatient	5 (1.1)	4 (1.3)	9 (1.2)	
Wound classification, n (%)				
Clean	127 (27.5)	88 (29.2)	215 (28.2)	0.685
Clean/contaminated	310 (67.1)	201 (66.8)	511 (67.0)	
Contaminated	13 (2.8)	8 (2.7)	21 (2.8)	
Dirty/infected	12 (2.6)	4 (1.3)	16 (2.1)	
Operation time (min), mean (SD)	561.8 (206.0)	554.3 (196.5)	558.8 (202.2)	0.6178
Length of hospital stay (d), mean (SD)	6.0 (24.1)	5.6 (22.0)	5.9 (23.3)	0.8213

Abbreviations: COVID-19, coronavirus disease 2019; n, frequency; SD, standard deviation.

Table 3 Postoperative complications following head and neck free flap reconstruction (total)

Postoperative complication	Historical control N = 462	COVID-19 N = 301	Total cohort N = 763	p-Value
Wound-related, n (%)				
Superficial skin infection	42 (9.1)	27 (9.0)	69 (9.0)	1.000
Deep incisional infection	15 (3.3)	8 (2.7)	23 (3.0)	0.829
Organ space infection	14 (3.0)	14 (4.7)	28 (3.7)	0.245
Wound dehiscence	25 (5.4)	15 (5.0)	40 (5.2)	0.869
Pulmonary, n (%)				
Pneumonia	28 (6.1)	13 (4.3)	41 (5.4)	0.328
Pulmonary embolism	4 (0.9)	1 (0.3)	5 (0.7)	0.653
Renal/genitourinary, n (%)				
Renal insufficiency	0 (0)	0 (0)	0 (0)	–
Acute renal failure	0 (0)	0 (0)	0 (0)	–
Urinary tract infection	2 (0.4)	7 (2.3)	9 (1.2)	0.033 ^a
Cardiac, n (%)				
Cardiac arrest	8 (1.7)	3 (1.0)	11 (1.4)	0.541
Myocardial infarction	4 (0.9)	4 (1.3)	8 (1.1)	0.719
Hematologic, n (%)				
Bleeding complication	92 (19.9)	66 (21.9)	158 (20.7)	0.523
DVT	13 (2.8)	5 (1.7)	18 (2.4)	0.341
Systemic, n (%)				
Shock/sepsis	22 (4.8)	8 (2.7)	30 (3.9)	0.182
Unplanned reoperation, n (%)	73 (15.8)	61 (20.3)	134 (17.6)	0.120

Abbreviations: COVID-19, coronavirus disease 2019; DVT, deep vein thrombosis; n, frequency.

^aStatistical significance was set up for a p-value less than 0.05.

complication was surgical wound-related complication in 20.9% of the cases. The rates of pulmonary (6.1%), cardiac (2.5%), renal/genitourinary (1.2%), and systemic complications (3.9%) were overall low in this cohort. On univariate analysis, the only significant difference in complications between the groups was the higher rate of urinary tract infections in the COVID-19 group.

Furthermore, there were no significant differences between the two groups on subgroup analysis of those who underwent HN free flap reconstruction during the second quarter of the year (April–June; ► **Table 4**). After stratifying cases by the location of reconstruction, we found that those who had free flap reconstruction of the mandible were significantly more likely to have a medical complication ($p < 0.05$; ► **Table 5**).

Risk Factors and Timing of Unplanned Reoperation

Similar type of surgical procedures for unplanned reoperation were evidenced between the two groups except for debridement procedures, in which pre-COVID-19 group had 23.08% cases versus 5.55% in the during COVID-19 group (► **Supplementary Table S3**). When controlling for sociodemographic and clinical risk factors, no differences were found between the pre-COVID-19 and during COVID-19 groups concerning unplanned reoperation ($p = 0.127$; ► **Table 6**).

Additionally, dependent function status ($p = 0.021$) and postoperative wound infection ($p < 0.001$) was found to have increased odds of undergoing unplanned reoperation after HN flap reconstruction after adjusting other cofactors constant (► **Table 6**, ► **Fig. 2**).

Discussion

The COVID-19 pandemic has impacted patient care significantly due to its unprecedented nature. The health care system had to adapt in a sensible and timely manner to consistently provide the best service conceivable. Patients with HN malignancies were not the exception to the burden generated by the COVID-19 pandemic.^{13–18} This study's findings highlight from a national standpoint that, in general, patients who underwent HN flap reconstruction during the pandemic were not statistically significantly different from patients before the pandemic in regard to clinical outcomes. In other words, the health care system was able to overcome surgical challenges among the HN population, which translated into similar postoperative clinical outcomes between before and during pandemics cohorts. This likely included COVID-19 screening, management of treating COVID-19 positive patients perioperatively, and following these patients after discharge.

Table 4 Postoperative complications following head and neck free flap reconstruction during the second quarter of the year (April-June)

Postoperative complication	Historical Q2 N = 79	COVID-19 Q2 N = 107	Total N = 186	p-Value
Wound-related, n (%)				
Superficial skin infection	9 (11.4)	4 (3.7)	13 (7.0)	0.077
Deep incisional infection	2 (2.5)	4 (2.8)	5 (2.7)	1.000
Organ space infection	4 (5.1)	7 (6.5)	11 (5.9)	0.762
Wound dehiscence	4 (5.1)	5 (4.7)	9 (4.8)	1.000
Medical-related complications				
Pulmonary, n (%)				
Pneumonia	4 (5.1)	5 (4.7)	9 (4.8)	1.000
Pulmonary embolism	0 (0)	0 (0)	0 (0)	–
Renal/genitourinary, n (%)				
Renal insufficiency	0 (0)	0 (0)	0 (0)	–
Acute renal failure	0 (0)	0 (0)	0 (0)	–
Urinary tract infection	0 (0)	3 (2.8)	3 (1.6)	0.263
Cardiac, n (%)				
Cardiac arrest	0 (0)	1 (0.9)	1 (0.5)	1.000
Myocardial infarction	0 (0)	0 (0)	0 (0)	–
Hematologic, n (%)				
Bleeding complication	14 (17.7)	25 (23.4)	39 (21.0)	0.369
DVT	2 (2.5)	1 (0.9)	3 (1.6)	0.575
Systemic, n (%)				
Shock/sepsis	1 (1.3)	0 (0)	1 (0.5)	0.425
Unplanned reoperation, n (%)	14 (17.7)	19 (17.8)	33 (17.7)	1.000

Abbreviations: COVID-19, coronavirus disease 2019; DVT, deep vein thrombosis; n, frequency.

Table 5 Differential postoperative complications following head and neck free flap reconstruction

Location of reconstruction, n (%)	Differential in medical complications ^a	p-Value	Differential in wound complications ^a	p-Value	Differential in reoperation ^a	p-Value
Oral cavity	(+)0.68%	0.903	(+)1.38	0.787	(+)7.3	0.085
Pharynx	(+)0.05%	1.000	(+)0.43	1.000	(+)6.24	0.369
Mandible	(-)41.27%	0.049 ^a	(-)1.59	1.000	(-)17.46	0.393
Maxilla	(-)36.67%	0.111	(+)3.33	1.000	(+)33.33	0.121
Surface structures	(+)5.77	0.774	(+)4.91	0.567	(-)6.2	0.689

^aRefer to at least one complication within those groups were present.

HN surgeons triaged patients based on their disease, urgency, and probability of improved outcome with surgery.¹³ This contingency measure was due to the fact that these type of procedures pose a unique risk compared with other specialties, as they work with the upper airway.¹³ Thus, both patients and health care providers are specifically vulnerable to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) transmission.¹⁹ It has been described that patients with SARS-CoV-2 disease who undergo surgical procedures suffer worse surgical outcomes

than those without the disease.^{20–23} Therefore, much-needed recommendations and new protocols rapidly emerged to decrease virus spread while preventing patients' disease progression and surgical complications.^{14,15,24–27} In fact, patients with indolent neoplastic diseases were shifted to undergo nonsurgical alternatives more than before the pandemic.¹³ Indisputably, long-term studies assessing the true impact of this first-time shift are needed to expose not only changes in clinical outcomes, but also in patient-reported outcomes.

Table 6 Multivariable analysis of risk factors for reoperation

Potential risk factors for reoperation	COVID-19 group ^a	
	OR (95% CI)	p-Value
Reconstruction during COVID-19	1.51 (0.89–2.57)	0.127
Age > 55	0.65 (0.32–1.30)	0.221
BMI \geq 30	0.74 (0.38–1.45)	0.387
African American race	0.80 (0.26–2.48)	0.704
Hispanics	1.24 (0.36–4.29)	0.737
Dependent functional status	6.92 (1.34–35.74)	0.021 ^a
ASA classification > 2	1.04 (0.49–2.20)	0.919
Smoker	1.51 (0.82–2.80)	0.189
Diabetes	0.36 (0.09–1.49)	0.158
Hypertension	1.26 (0.70 – 2.25)	0.441
COPD	0.84 (0.31–2.29)	0.728
Disseminated cancer	1.19 (0.39–3.60)	0.756
Bleeding disorder	2.80 (0.47–16.71)	0.256
Preoperative weight loss > 10% of body weight	0.89 (0.38–2.09)	0.789
Preoperative steroid use	1.70 (0.46–6.22)	0.423
Postoperative wound infection	5.41 (2.98–9.81)	< 0.001 ^b
Wound classification > 2 (contaminated)	0.77 (0.38–1.54)	0.457
Hematocrit < 30	0.67 (0.28–1.62)	0.378
Albumin < 3.5	0.78 (0.45–1.38)	0.395
Operative time in top 25% (> 679 min)	1.55 (0.85–2.84)	0.156
Length of hospital stay in top 25% (>13 d)	4.56 (2.56–8.12)	< 0.001 ^b

Abbreviations: ASA, American Society of Anesthesiologists; BMI, body mass index; CI, confidence interval; COPD, chronic obstructive pulmonary disease; COVID-19, coronavirus disease 2019; OR, odds ratio.

^aStatistical significance was set up for a p-value less than 0.05.

Our study results showed that HN flap reconstruction for malignant diseases could be performed safely during COVID-19. Overall, no major differences in terms of sociodemographic characteristics, clinical profile, operative time, length of hospital stay, and 30-day postoperative outcomes between before and during the COVID-19 pandemic were evidenced in this cohort. Of note, we found statistically significantly higher rates of urinary tract infection in the COVID-19 group. However, we believe that overall complication rates were very low; therefore, this finding is not clinically relevant. Similar findings were demonstrated by Wai et al, where the authors presented in 2020 their COVID-19 pandemic experience among this population.⁹ In 63 operations during COVID-19 and 84 operations during pre-COVID-19, the authors found similar perioperative outcomes with no recorded viral transmission within both health care providers and patients.⁹ This might reflect proper preoperative screening protocols. In fact, studies have shown that with an appropriate screening process and postoperative care, these elective procedures can be performed safely during the current pandemic while minimizing delays in treatment.^{28–31}

Furthermore, it could be then reasonable to infer that due to the massive elective surgeries' cancellations secondary to the COVID-19 pandemic, delays in diagnosis and treatment,

as well as a surge of advanced HN cancer are anticipated.^{32–34} Moreover, Kiong et al, in a retrospective review, evidenced an increased tumor burden in patients with HN malignancies, despite having similar time of diagnosis.³² Not surprisingly, Rygalski et al described that delays in surgical treatment significantly increased the risk of death.³⁵ Even pre-pandemic, the hazard of death increases by 4.6% for every 30-day delay in time-to-surgery among this population.³⁵ However, the long-term impact of the pandemic, in relation to organic and mental burden among HN patients who suffered from delays is yet to be elucidated.³⁶ Until now, lessons learned to maximize outcomes have been described in the current literature. During the pandemic, Han et al proposed a paradigm shift in HN cancer management.²⁶ The authors suggested the use of a multidisciplinary team to define "essential surgery" that require immediate life or function-threatening diseases that necessitate surgeries, as well as to identify those patients that might benefit from nonsurgical options.²⁶ Therefore, a multidisciplinary approach might play an important role in maximizing patient outcomes. If this approach is implemented in the future, this could potentially (1) avoid delays and (2) improve clinical and patient-reported outcomes.²⁶




Fig. 2 Percentage of reoperation over the first postoperative month. COVID-19, coronavirus disease 2019; POD, postoperative day.

Additionally, HN surgeons focused on reconstruction must be aware of the potential risk factors increasing the likelihood of undergoing unplanned reoperations. This study's results demonstrate that dependent functional status and postoperative wound infection were found to be risk factors for unplanned reoperation after conducting an adjusted analysis. Undergoing unplanned reoperations increases the burden placed not only on the patients, but also surgeons and the health care system.³⁷ Sangal et al findings aligns with ours; they found that total operative time, surgical site infection, and wound dehiscence were significantly associated with reoperation in major HN surgeries.³⁷ Moreover, they also identified further risk factors for reoperation not evidenced in our cohort, such as being African American, having disseminated cancer, and being ventilator dependent for more than 48 hours after surgery.³⁷ Therefore, proper control of the modifiable risk factors in the postoperative setting is critical to avoid reoperation among this population.

Limitations of a big database should be taken into consideration when assessing the internal and external validity of this study. This study used a national database with predetermined data collection points that limit the assessment of other variables that might be of interest for patients who underwent HN flap reconstruction, such as type of anesthesia medications, recovery protocols, cancer stage, and history of radiation. Also, long-term complications cannot be evaluated with this database. Lastly, surgical techniques, surgical decisions, and specific patient preoperative details were not able to be ascertained.

Conclusion

HN flap reconstruction can be performed safely during the COVID-19 era. Similar profile status on immediate postoperative outcomes were evidenced between the cohorts. Standardized and rigorous protocols for surgical candidates must be strictly followed to avoid disease progression and optimize surgical outcomes. Lastly, further studies assessing long-term outcomes during the pandemic are of utmost importance to elucidate the true impact of the COVID-19 pandemic on this population.

Ethical Approval

The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. This study was IRB approved (#2021D001052) at the Beth Israel Deaconess Medical Center.

Author Contributions

Conception and design are made by V.P.B., N.E., B.T.L., S.J.L.; administrative support was given by B.T.L., S.J.L.; collection and assembly of data was done by V.P.B., N.E.; data analysis and interpretation were done by V.P.B., N.E., S.J.L.; V.P.B., N.E., A.V. did the manuscript writing; final approval of manuscript was made by V.P.B., N.E., A.V., C.D.C., J.R.K., S.M.M., N.H., B.T.L., S.J.L.

Funding

S.J.L. has Royalties from McGraw-Hill Publishing. The rest of the authors do not have a financial interest in any of the information mentioned in this manuscript.

Conflict of Interest

None declared.

References

- 1 Kang J-Y, Michels A, Lyu F, et al. Rapidly measuring spatial accessibility of COVID-19 healthcare resources: a case study of Illinois, USA. *Int J Health Geogr* 2020;19(01):36
- 2 Moynihan R, Sanders S, Michaleff ZA, et al. Impact of COVID-19 pandemic on utilisation of healthcare services: a systematic review. *BMJ Open* 2021;11(03):e045343
- 3 Chi D, Chen AD, Dorante MI, Lee BT, Sacks JM. Plastic surgery in the time of COVID-19. *J Reconstr Microsurg* 2021;37(02):124–131
- 4 Elmer NA, Bustos VP, Veeramani A, et al. Trends of autologous free-flap breast reconstruction and safety during the coronavirus disease 2019 pandemic. *J Reconstr Microsurg* 2023;39(09):715–726
- 5 Wong S, Payton JJ, Lombana NF, et al. A protocol for safe head and neck reconstructive surgery in the COVID-19 pandemic. *Plast Reconstr Surg Glob Open* 2020;8(11):e3258
- 6 Jain S, Gupta S, Singh TP, et al. Short term outcomes of head and neck oncology surgery during Covid-19 pandemic: experience from a tertiary cancer care centre in North India. *Indian J Otolaryngol Head Neck Surg* 2022;74(Suppl 2):2822–2826
- 7 Thione A, Sánchez-García A, Pérez-García A, García-Vilarinho E, Salmerón-González E, Balaguer-Cambra J. A protocol for performing reconstructive microsurgery on patients with COVID-19. *Plast Surg Nurs* 2021;41(01):36–39
- 8 Thacoor A, Sofos SS, Miranda BH, et al. Outcomes of major head and neck reconstruction during the COVID-19 pandemic: the St. Andrew's centre experience. *J Plast Reconstr Aesthet Surg* 2021;74(09):2133–2140
- 9 Wai KC, Xu MJ, Lee RH, et al. Head and neck surgery during the coronavirus-19 pandemic: the University of California San Francisco experience. *Head Neck* 2021;43(02):622–629
- 10 ACS National Surgical Quality Improvement Program. Accessed June 28, 2021 at: <https://www.facs.org/quality-programs/acs-nsqip>
- 11 Shiloach M, Frencher SK Jr, Steeger JE, et al. Toward robust information: data quality and inter-rater reliability in the American College of Surgeons National Surgical Quality Improvement Program. *J Am Coll Surg* 2010;210(01):6–16
- 12 Henderson WG, Daley J. Design and statistical methodology of the National Surgical Quality Improvement Program: why is it what it is? *Am J Surg* 2009;198(05):S19–S27
- 13 Piccirillo JF. Otolaryngology-head and neck surgery and COVID-19. *JAMA* 2020;324(12):1145–1146
- 14 See A, Go LK, Teo CEH, Teo NWY, Toh ST. Adaptations of a tertiary otorhinolaryngology Head and Neck Surgery Department in Singapore during the COVID-19 outbreak. *Ann Otol Rhinol Laryngol* 2021;130(02):177–181
- 15 Givi B, Schiff BA, Chinn SB, et al. Safety recommendations for evaluation and surgery of the head and neck during the COVID-19 pandemic. *JAMA Otolaryngol Head Neck Surg* 2020;146(06):579–584
- 16 Cho RHW, Yeung ZWC, Ho OYM, et al. Pearls of experience for safe and efficient hospital practices in otorhinolaryngology-head and neck surgery in Hong Kong during the 2019 novel coronavirus disease (COVID-19) pandemic. *J Otolaryngol Head Neck Surg* 2020;49(01):30
- 17 Liu Z, Zhang L. At the center of the COVID-19 pandemic: lessons learned for otolaryngology-head and neck surgery in China. *Int Forum Allergy Rhinol* 2020;10(05):584–586
- 18 Kowalski LP, Sanabria A, Ridge JA, et al. COVID-19 pandemic: effects and evidence-based recommendations for otolaryngology and head and neck surgery practice. *Head Neck* 2020;42(06):1259–1267
- 19 Khariwala SS, Weinreich HM, McCoul ED, et al. Leveraging COVID-19-inspired changes to advance otolaryngology-here to stay. *JAMA Otolaryngol Head Neck Surg* 2020;146(07):605–607
- 20 Colosimo C, Kelly J, Coker J, et al. Unscreened: urgent and emergent surgical outcomes in the early COVID-19 pandemic. *Cureus* 2020;12(12):e11878
- 21 Bustos SS, Bustos VP, Nguyen MT. Elective surgery in asymptomatic patients with undetected SARS-CoV-2 infection: lessons learned from the global operating room. *Br J Surg* 2020;107(10):e399–e400
- 22 Mohanka MR, Mahan LD, Joerns J, et al. Clinical characteristics, management practices, and outcomes among lung transplant patients with COVID-19. *J Heart Lung Transplant* 2021;40(09):936–947
- 23 Osorio J, Madrazo Z, Videla S, et al; COVID-CIR Collaborative Group Members of the COVID-CIR Collaborative Group. Analysis of outcomes of emergency general and gastrointestinal surgery during the COVID-19 pandemic. *Br J Surg* 2021;108(12):1438–1447
- 24 Galloway TJ, Kowalski LP, Matos LL, Junior GC, Ridge JA. Head and neck surgery recommendations during the COVID-19 pandemic. *Lancet Oncol* 2020;21(09):e416
- 25 Krajewska Wojciechowska J, Krajewski W, Zub K, Zatoński TKrajewska (Wojciechowska) J. Review of practical recommendations for otolaryngologists and head and neck surgeons during the COVID-19 pandemic. *Auris Nasus Larynx* 2020;47(04):544–558
- 26 Han AY, Miller JE, Long JL, St John MA. Time for a paradigm shift in head and neck cancer management during the COVID-19 pandemic. *Otolaryngol Head Neck Surg* 2020;163(03):447–454
- 27 Wunsch A, Pitak-Arnnop P. Strategic planning for maxillofacial trauma and head and neck cancers during COVID-19 pandemic-December 2020 updated from Germany. *Am J Otolaryngol* 2021;42(04):102932
- 28 Ferrari M, Paderno A, Giannini L, et al. COVID-19 screening protocols for preoperative assessment of head and neck cancer patients candidate for elective surgery in the midst of the pandemic: a narrative review with comparison between two Italian institutions. *Oral Oncol* 2021;112:105043
- 29 Brar S, Ofo E, Hyde N, et al. Outcomes of elective head and neck confirmed or suspected cancer surgery during the COVID-19 pandemic. *Eur Arch Otorhinolaryngol* 2021;278(04):1277–1282
- 30 Ito A, Kobayashi K, Shiotsuka M, et al. Uniform infection screening allowed safe head and neck surgery during the coronavirus disease 2019 pandemic in Japan. *Jpn J Clin Oncol* 2021;51(03):400–407
- 31 Chen J, Chow A, Lee E, et al. Reintroducing flap reconstruction: one institution's safe return to flap surgery during the COVID-19 pandemic. *J Reconstr Microsurg* 2023;39(01):59–69
- 32 Kiong KL, Diaz EM, Gross ND, Diaz EM Jr, Hanna EY. The impact of COVID-19 on head and neck cancer diagnosis and disease extent. *Head Neck* 2021;43(06):1890–1897
- 33 Tevetoglu F, Kara S, Aliyeva C, Yildirim R, Yener HM. Delayed presentation of head and neck cancer patients during COVID-19 pandemic. *Eur Arch Otorhinolaryngol* 2021;278(12):5081–5085
- 34 De Luca P, Bisogno A, Colacurcio V, et al. Diagnosis and treatment delay of head and neck cancers during COVID-19 era in a tertiary care academic hospital: what should we expect? *Eur Arch Otorhinolaryngol* 2022;279(02):961–965
- 35 Rygalski CJ, Zhao S, Eskander A, et al. Time to surgery and survival in head and neck cancer. *Ann Surg Oncol* 2021;28(02):877–885
- 36 Dermody SM, Shuman AG. Psychosocial implications of COVID-19 on head and neck cancer. *Curr Oncol* 2022;29(02):1062–1068
- 37 Sangal NR, Nishimori K, Zhao E, Siddiqui SH, Baredes S, Chan Woo Park R. Understanding risk factors associated with unplanned reoperation in major head and neck surgery. *JAMA Otolaryngol Head Neck Surg* 2018;144(11):1044–1051