

Endoscopy International Open

Plan-do-study-act (PDSA) interventions to improve real-world endoscopy unit productivity

Christopher Bradley, Matt Sumethasorn, Jin S Kim, Sarah Wang, Leslie Martinez, Megan Chang, Lillian Lemus, Behrod Katebian, Dara Bruce, Andy Lee, Rachel Baden, Hal Yee, James Buxbaum.

Affiliations below.

DOI: 10.1055/a-2290-0263

Please cite this article as: Bradley C, Sumethasorn M, Kim J S et al. Plan-do-study-act (PDSA) interventions to improve real-world endoscopy unit productivity. Endoscopy International Open 2024. doi: 10.1055/a-2290-0263

Conflict of Interest: The authors declare that they have no conflict of interest.

Abstract:

ABSTRACT:

Introduction: The Plan-Do-Study Act (PDSA) ramp is a framework that uses initial small changes to build consensus and momentum for subsequent, iterative process improvement. Our aim was to study its impact on endoscopy unit efficiency and throughput.

Methods: Following a granular time-and-motion analysis to evaluate baseline performance (phase 1) we instituted successive interventions and measured their impact on core efficiency metrics including procedure volume and turnover time (phases 2-3).

Results: We identified that inefficiency in turnover of anesthesia-supported endoscopy was the most crucial issue. Implementation of pre-procedure anesthesia visit in phase 2 reduced turnover time by 15.5 (95% CI 3.9-27.1 minutes). Subsequent changes (phase 3) including front-loaded procedure scheduling and parallel in-room preparation resulted in an 18% increase in procedure volume.

Conclusion:

The PDSA ramp model is an effective means of assessing operational processes, developing novel interventions, and building consensus to improve the real-world productivity in a resource-conscious manner.

MESH TERMS: Time and Motion Studies; Endoscopy, Gastrointestinal; Workflow; Efficiency, Organizational; Operative Time; Recovery Room; Cost savings; Hospital Units; Patient Care Team; Benchmarking; COVID-19

Corresponding Author:

Dr. James Buxbaum, University of Southern California Keck School of Medicine, Department of Internal Medicine, Division of Gastroenterology, Los Angeles, United States, James.Buxbaum@med.usc.edu

Affiliations:

Christopher Bradley, University of Southern California Keck School of Medicine, Department of Internal Medicine, Division of Gastroenterology, Los Angeles, United States

Matt Sumethasorn, University of Southern California Keck School of Medicine, Department of Internal Medicine, Division of Gastroenterology, Los Angeles, United States

Jin S Kim, University of Southern California, Division of Gastrointestinal Liver Disease, Los Angeles, United States

[...]

James Buxbaum, University of Southern California Keck School of Medicine, Department of Internal Medicine, Division of Gastroenterology, Los Angeles, United States



Plan-do-study-act (PDSA) interventions to improve real-world endoscopy unit productivity

ABSTRACT:

Introduction: The COVID-19 pandemic and its aftermath have exacerbated the mismatch between the high demand for endoscopic services and limited staff and resources. The Plan-Do-Study Act (PDSA) ramp is a framework that uses initial small changes to build consensus and momentum for subsequent change and continual improvement of a process. Our aim was to study its impact on endoscopy unit efficiency and throughput.

Methods: During phase 1, we conducted a granular time-and-motion analysis to evaluate baseline performance. A multidisciplinary team (MDT) comprised of study and quality improvement staff, physicians, nurses, and anesthesiologists analyzed our performance in comparison to benchmarks and developed interventions which were implemented in phases 2 and 3. Productivity outcomes included monthly and daily procedure volume for both anesthesia-supported-monitored anesthesia care (MAC) and moderate sedation procedures. Secondary efficiency outcomes included room turnover and first-case start time.

Results: We identified multiple operational inefficiencies in phase 1 and addressed them with sequential operational changes using the PDSA ramp model in phases 2

and 3. Utilization of a pre-procedure anesthesia clinic (PAC) for patients requiring MAC was initiated in phase 2. The turnover time decreased by a mean of 15.5 (95% CI 3.9-27.1 minutes) and mean daily MAC room throughput increased from 5.6 to 8.3 (mean difference 2.8 (95% CI 0.9-4.7)). During phase 3, we implemented front-loaded procedure scheduling, first procedure preparation in the procedure room, parallel MD/RN evaluation, as well as streamlined documentation processes. Data from phase 3 showed an increase in monthly procedure volume from 495.8 (± 41) to 583 (± 73); mean difference 87.2 (95% CI 0.1-174) and improvement in multiple efficiency metrics, including first case start time in both MAC (OR 28 [95% CI 2.6-297.6]) and moderate sedation (OR 3.6 [95% CI 1.7-7.8]) rooms.

Conclusion:

The PDSA ramp model is an effective means of assessing operational processes, developing novel and effective interventions, and building consensus to improve the real-world efficiency and productivity of a busy safety-net endoscopy unit in a resource-neutral manner.

ACKNOWLEDGEMENT:

MESH TERMS: Time and Motion Studies; Endoscopy, Gastrointestinal; Workflow; Efficiency, Organizational; Operative Time; Recovery Room; Cost savings; Hospital Units; Patient Care Team; Benchmarking; COVID-19

INTRODUCTION:

During and after the COVID-19 pandemic, staff shortages and mounting procedure backlogs exacerbated a pre-existing shortage of endoscopy resources. In the previous two decades, widespread screening for GI malignancies, an aging population, and reduced endoscopist training set the context for this shortfall.[1-3]

The post-pandemic shortage of nurses and technicians in the worldwide remains problematic, particularly in safety net (under-served) settings with limited resources available to support operations

Operations analysis, including time and flow studies, emerged in the early 20th century to optimize manufacturing. [5] Despite widespread adoption in anesthesia and surgical settings, these strategies have only recently been introduced to endoscopy units (EU).[6] The Plan-Do-Study-Act (PDSA) ramp cycle is a framework used to systematically introduce and implement changes for quality improvement and involves a stepwise process. The four steps are development of a proposed intervention (plan), implementation of this intervention (do), analysis of how the intervention affects outcomes (study) and then adaptation of the findings into practice (act). This cycle is then repeated with modifications to the

intervention as needed. Interventions start small to prevent disruptions in opinion and workflow and are subsequently scaled or “ramped up” over time as support is gathered through proof of concepts. Initial results encourage organizational consensus and stakeholder buy-in to the quality improvement (QI) initiative and support larger changes and operational improvements downstream (Figure 1). We report the implementation of the PDSA ramp model to evaluate and improve endoscopy unit efficiency and productivity.

METHODS:

Setting:

The study was conducted in the Los Angeles County University of Southern California (LAC+USC) Medical Center Endoscopy Unit (EU). Prior to initiation, it was approved by the University of Southern California Health Sciences Institutional Review Board. The EU has five endoscopy rooms and 90% of procedures are performed using moderate sedation. One day each week, one room is dedicated to monitored anesthesia care (MAC) procedures supported by a certified registered nurse anesthetist (CRNA). All endoscopies are performed by fellows under attending supervision. The EU is supported by an adjacent 12 bed pre-procedure/recovery unit (PPU).

Process-flow map and time-flow study

An initial EU process flow map was created via observation of workflow and interviews with unit staff (**Figure 2**). To record the patient journey through the EU, the process flow was transferred to a timesheet and attached to the patient chart. Unit or study staff manually recorded the time when each step was initiated and concluded. (**Supplementary Figure 1**). A separate flow sheet was used for MAC cases to account for additional steps involved. Data were collected weekly for all procedures in each standard procedure room. For MAC procedure days, data were collected on the same day each week. Time points were cross-checked against mandatory “scope tracking logs,” including time of patient entry to procedure room, procedure start time, scope exit time, and time of patient exit from procedure room (**Supplementary Figure 1**). Overall procedure volume was aggregated from the electronic medical records (EMR) by EU administrative staff and included both MAC and non-MAC days for all rooms.

Phases of the PDSA Cycle

Three phases were designated. Phase 1 was observational, aimed at establishing

baseline performance and identifying areas for intervention (**Figure 1**). A multi-disciplinary team (MDT) of study staff, endoscopy nurses, QI staff, physicians, and anesthesiologists reviewed data against established benchmarks (7,8) and proposed interventions aimed at improving efficiency and throughput. The aim was to include all key stakeholders (nurses, endoscopists, anesthesiologists, and administrators) in decision making to obtain the perspective and buy-in of these groups. The entire medical team of the endoscopy unit was provided with a survey on perceived operational inefficiencies to better help identify areas of intervention. Although some issues such as additional staff or equipment could not be addressed due to cost containment constraints, other intervenable domains such as room turnover time and late first case start time were identified. Generally, perceptions of operational inefficiencies correlated well with deficiencies seen in comparison to established benchmarks. The combination of objective and subjective data allowed us to quickly implement interventions using the PDSA ramp model (**Figure 1**).

Phase 2 involved implementing the first targeted intervention, the utilization of a pre-anesthesia clinic (PAC) visit, to address two major deficiencies identified in Phase 1, excessive turnover time and low throughput through the MAC room. This required collaboration with our anesthesia colleagues to help set up a new

workflow to include the PAC prior to endoscopy. To help acclimate staff to these new changes, a process flow map was provided to each staff member to graphically describe the new pathway. At the end of phase 2, we studied or reviewed the impacts of our primary intervention and found significant improvement on MAC room turnover time and throughput. It was initially difficult to motivate staff to engage in the proposed changes. Once we were able to demonstrate proven and measurable outcomes, we were able to gain stakeholder and staff support to enact larger and broader changes. In phase 3, we enacted multiple simultaneous interventions to further improve the efficiency of the endoscopy units (see results section). At the end of this cycle, impacts of all primary and secondary interventions were measured to determine their impact.

Main (Productivity) Outcomes

The primary outcome of interest was monthly total (anesthesia supported MAC and non-MAC) procedure volume performed in the EU pre- and post-phase 2 intervention. A co-primary productivity outcome was the daily volume of procedures in the MAC room following phases 2 and 3.

Operational Metrics/Outcomes

Outcomes included 'Turnover time' which was defined as the interval between a patient exiting the procedure room and entry of the next patient, 'Procedure end-to-

room exit time,' the interval between endoscope removal and patient egress from the procedure room and 'In-room-to-procedure start time,' the interval between patient entry to procedure room and endoscope insertion. 'First case on-time start' was defined as the proportion of cases in which the endoscope had been inserted by 08:00.

Statistical Analysis:

We calculated means (standard deviations/confidence intervals) for continuous variables with normal distribution and medians (interquartile ranges) for nonparametric distributions. To compare baseline (phase 1) metrics versus benchmarks we used one-sample t-tests for continuous (room turnover time, in room to procedure start time, procedure time, procedure end to room exit time) and one sample test of proportions for categorical variables (first case on-time start).

To compare the continuous productivity outcomes (main outcomes of the study) of total monthly procedure volume and daily MAC volume we used linear regression. Sensitivity analysis of total volume excluding March (as interventions were implemented mid-month) was performed. For the operational outcomes of room turnover time, procedure end-to-room-exit time, in room to procedure start time)we used linear regression. For the the categorial operational outcome of first

case on-time start we used logistical regression. Statistical analysis was performed using STATA 14.0 (College Station, TX).

RESULTS:

Patients:

Prospective data was captured for 673 patients undergoing endoscopic procedures during the study period of 9 months divided between the phases. In phase 1, 265 patients were evaluated to establish baseline metrics and derive primary interventions. During phase 2, primary interventions were assessed for 167 patients. Phases 1 and 2 results were used to develop secondary interventions and assessed during phase 3, involving 241 additional patients.

Population Metrics versus Benchmarks

Baseline analysis revealed mean turnover time of 45.3 minutes (95% CI 31.8-58.8) for MAC procedures exceeding the benchmark of 26.6 minutes. First case on-time start for both the MAC (20%) and moderate sedation rooms (29.3%) were below the benchmark of 64.5% (**Supplementary Table 1**). Other time measures for MAC procedures met benchmarks. For moderate sedation procedures, the time

from procedure end to room exit was 16.5 minutes (95% CI 15.3-17.6) vs the benchmark of 9.4 minutes. According to the stakeholder survey, room turnover time was the greatest perceived inefficiency in the EU (**Figure 3**).

Identification and Impact of First Targeted Intervention

Suboptimal throughput in the MAC room was designated as the first target for intervention. The MDT identified that delayed turnover was the rate limiting factor to productivity which was primarily a consequence of the need for patient evaluation on the day of procedure by the anesthetist. To address this concern, the team recommended that all endoscopy patients scheduled for MAC procedures be referred to a pre-anesthesia clinic visit (PAC) which eliminated any inter-procedure anesthesia evaluations the day of the procedure by moving them to the day of endoscopy teaching (**Table 1**). Standing appointments were created, allowing patients to visit the PAC immediately after their endoscopy teaching visit. In the PAC, history and physical assessment were performed and, if necessary, cardiac/biochemical tests ordered so that results were available in advance, rather than on procedure day; and was thus a resource neutral change. This was done at the same time as the required endoscopy teachings to further streamline efficiency. Following implementation, turnover time for the MAC room decreased by a mean

of 15.5 minutes (95% CI of 3.8 to 27.1) (**Table 2**) and now aligned with established benchmarks. The daily anesthesia room volume from phase 1 to 2 correspondingly increased from 5.6 (± 2.1) to 8.3 (± 2.1); mean difference of 2.75 (95% 0.8-4.7) (**Table 2**).

Definition of Secondary Interventions

Following phase 2 the team devised additional interventions for implementation during phase 3 (**Figure 1, Table 1**). The first MAC and moderate sedation patients of the day were prepared in the procedure room to directly bypass the pre-procedure unit and minimize transfer time. Daily procedure start times were recorded in a report card. Additionally, a greater proportion of procedures were scheduled earlier in the day to mitigate the impact of patient no-shows, or delays due to unplanned (or excessively long) procedures or adverse events (**Table 1**).

. To address late first procedure start and turnover we instructed our team to prepare the first patients of the day in the procedure room to minimize transfer time and perform concurrent tasks for patient preparation throughout the day. Specifically, we asked our nurses to place intravenous lines and monitoring equipment (i.e. electrocardiogram leads, blood pressure cuffs, oxygen saturation

monitors) while the endoscopists documented clinical history and obtained informed consent. A policy requiring procedure report completion before patient egress from the endoscopy room was identified as delaying procedure completion to room exit time. During phase 3, this policy was modified such that only a very brief note and instruction by the endoscopists to guide care in the recovery room were required before egress from the procedure room.

Primary Outcome: Endoscopy Unit Productivity

Post-interventions, the primary outcome of overall procedure volume per month increased from 495.8 (± 40.7) to 583 (± 73); mean difference of 87.2 (95% CI 0.1-174) (**Table 2**). In sensitivity analysis excluding March 2022 (changes implemented mid-month), this increased to 623.5 (± 29); mean difference 127.7 (95% CI, 49.8-205.6). Additionally, the other main productivity outcome, mean number of daily MAC procedures, increased from 5.6 (± 2.1) to 9.6 (± 2.7) following phase 2 and phase 3 interventions: mean difference 4.0 (95% CI, 2.1-5.9).

Operations Outcomes

The reduction in MAC turnover time was sustained at 30 (± 24 minutes) in phase 3, equal to the benchmark (**Table 2**). Additionally, there was an improvement in the

first case on-time start percentage for the MAC room to 87.5% (OR 28 (95% CI 2.6-297.9) and the moderate sedation room to 60% (OR 3.6 (95% CI 1.7-7.8). Procedure end-to-room-exit time decreased by 2.1 (95% CI 0.8-3.3) minutes. There was a non-significant reduction in turnover time for moderate sedation procedures from 24.5 (± 36.7) to 22 (± 32.1) minutes.

DISCUSSION:

Time-flow and operations management approaches were introduced in healthcare to optimize operating room function. [5, 7][6, 15, 16] Akin to surgery, innovative endoscopic technology has increased costs while reimbursements have been reduced creating the need to improve efficiency.[6] With more than 20 million procedures performed annually, GI endoscopy represents the highest volume procedure performed in ambulatory care centers in the United States and improved efficiency has the potential to provide substantial benefit.[1, 2]. This study demonstrates that the PDSA ramp model is effective at maximizing efficiency and productivity in a resource neutral manner.

Prior work to use data to improve EU operations has focused on single interventions and, in most cases, utilized simulated results over real world data.

Day et al used a time-motion analysis and discrete event simulation to propose changes that might improve efficiency including shorter procedure times (60 to 45 minutes), modified scheduling, and expanded human resources.[9] Nevertheless, simulation was the primary method of validation as testing interventions in a large patient population was considered unfeasible. Kaushal et al performed a prospective flow analysis to identify a small pre-procedure/recovery unit as the bottleneck in their EU and responded to this by utilizing procedure rooms to prepare patients whenever available. [8] They demonstrated that this single change improved on-time procedure start by 51% and reduced overhead costs in a large subsequent cohort. Others have collected critical data about efficiency metrics to identify problems and used modeling and other systems approaches to propose solutions. [10-14].

The success of the present study in implementing multiple interventions is credited to the PDSA ramp model, using carefully chosen, limited preliminary actions to gain stakeholder buy-in prior to enacting extensive changes as well as the use of a multidisciplinary team approach. Our study identified the importance of increasing anesthesia room productivity with time-flow analysis delineating turnover time as the primary problem. The MDT recognized that testing and evaluation by the anesthetist on the day of the procedure delayed patient flow and recommended

evaluating patients in the PAC before their procedures. This intervention was chosen to generate agency in the project for EU personnel and correspondingly addressed the most reported cause of delays from the stakeholder survey. The success of this first intervention in the “ramp” empowered the intervention team and prompted support from EU staff to carry out more extensive interventions in later phases. Following this intervention, the mean number of procedures doubled during phase 2, and was sustained during phase 3. Other interventions such as strategies to prepare the first patient of the day in the procedure room, parallel processing, and booking cases earlier in the day were simultaneously implemented. Consequently, in phase 3 we confirmed a significant improvement in the operational outcomes of first case on-time start and turnover times. Of even greater importance, the PDSA process as whole resulted in a significant improvement in the main outcome of interest which was endoscopy unit productivity. This was measured by the overall number of monthly procedure and the daily number of procedures in the resource intensive room requiring anesthesia support.

In the current study, the multidisciplinary team (MDT) reviewed the data, crafted proposed interventions, and paired our acquisition of operational metrics with a stakeholder survey to define barriers to productivity. There are valid concerns that using quantitative information to guide operations may dehumanize clinical

practice[17]. In contrast to top-down interventions, the use of a multidisciplinary team cognizant of local factors in the PSDA ramp model enables a nuanced approach that accounts for cultural and structural factors of the environment. [8, 11] Including nurses, endoscopists, anesthesiologists, and administrators not only provides broad perspective for the group but facilitates dissemination and improves the willingness of the entire medical team to carry out the recommendations of the MDT.

Our study has several limitations. Each institution has its own unique qualities and resources that presents as logistical barriers specific to that site. The results of the present study were initiated in a safety-net health center with the challenges of limited resources exacerbated by operating in the post-pandemic setting. Safety net hospitals serve a disproportionately higher number of vulnerable patients such as the uninsured, unhoused, and ethnic minorities regardless of their ability to pay. They often operate with limited financial resources and staffing. Though generalizability of our exact interventions may be limited to medical centers with similar patient populations and resources, this strategy is transferable to multiple settings. While we highlight specific low-cost strategies that may be implemented in institutions with similar challenges, a greater aim of this study is to demonstrate that the general methodology of the PSDA cycle offers a systematic way of implementing

and scaling multiple interventions to maximize efficiency and productivity. This would provide an added benefit in settings with incentives for improved productivity and efficiency

Additionally, though relatively fewer procedures were supported by the anesthesia services, optimization of this resource was the target of our primary intervention. Another challenge is that prolonged procedures or technical complications may adversely impact endoscopy unit function. To systematically adjust for no-shows, lengthy procedures, and adverse events, as part of our secondary interventions we scheduled a greater proportion of procedures in the morning. This affords greater flexibility for subsequent procedure flow throughout the day. Although this does not completely eliminate delays due to emergent cases or lengthy procedures, overall EU productivity ultimately improved in part due to this buffer for unforeseen events. Finally, observational bias due to the Hawthorne effect may have led us to overestimate improvement, which may wane when the period of observation concludes. We aim to continue monitoring operations by programming key steps of EU processes into the EMR and meeting regularly to follow up on and recommend additional interventions.

This study also underscores several needs in the field of endoscopy operations research. In phase one, we determined baseline performance metrics, comparing them to values from prior studies. [7, 8] Nevertheless, terms such as “turnover time” are defined heterogeneously. Future studies will benefit from standardized metrics (quantifiable variables) of endoscopy operations according with Agency for Health Care Research and Quality (AHRQ) guidance and reference values for defined scenarios (benchmarks). [7, 18] Future systematic work using a scientific approach to operations may be used to develop practical endoscopy operations guidelines. For example, Kaushal et al found that a ratio of 1.67 endoscopy rooms to pre/post recovery unit (PPU) beds was inadequate and proposed a ratio of 2 to 2.5. [8] A number of studies have demonstrated that providing more than 1 procedure room per physician increases efficiency if the “efficiency quotient” (actual procedure time : total time in EU) is less than 0.5. [11, 19, 20]

In this study, we demonstrate the use of the PDSA ramp model to improve core efficiency metrics in the endoscopy unit, employing an multidisciplinary team cognizant of local factors to guide successful intervention. The ramp model employed quantitative metrics to drive policy in a patient care setting and engendered cooperation amongst personnel to optimize the implementation of interventions. Prospective analysis confirms that this approach may additionally

improve room throughput, overall productivity, and core efficiency metrics (i.e. first case on-time start) in an endoscopy unit.

Figure Legend

Figure 1. The Plan-Do-Study-Act (PDSA) ramp model process began in phase 1 with a baseline analysis of the endoscopy unit to identify inefficiencies in the workflow. Targeted interventions were developed between phases 2 and 3 and their effect analyzed between phases. Promotion of consensus and buy-in during the process allows for progressively greater operational interventions to be achieved over time.

Figure 2. A diagram of the patient flow through the endoscopy unit (EU) from check-in to discharge.

Figure 3. Results from an open-ended stakeholder survey sent to all EU staff, fellows, and faculty, that asked respondents to identify the most significant contributor to inefficiency in the EU.

Supplementary Figure 1: Time flow data sheet used for monitored anesthesia care (MAC) procedures. The CRNA (certified nurse anesthetist) assessment was not included on the data sheet used for moderate sedation procedures. Procedure room

times were cross checked with Scope Tracking Logs that our institution uses for all endoscopy procedures in the EU.

REFERENCES:

1. Lin MY, Mishra G, Ellison J, et al. Differences in patient outcomes after outpatient GI endoscopy across settings: a statewide matched cohort study. *Gastrointest Endosc.* 2022;95(6):1088-97 e17.
2. Peery AF, Crockett SD, Murphy CC, et al. Burden and Cost of Gastrointestinal, Liver, and Pancreatic Diseases in the United States: Update 2021. *Gastroenterology.* 2022;162(2):621-44.
3. Meyer GS, Jacoby I, Krakauer H, et al. Gastroenterology workforce modeling. *JAMA.* 1996;276(9):689-94.
4. Buxbaum JL, Bradley CP, Lee WA, et al. Addressing postpandemic disruption in a safety net EU. *Gastrointest Endosc.* 2022;96(1):135-8 e1.
5. Lopetegui M, Yen PY, Lai A, et al. Time motion studies in healthcare: what are we talking about? *J Biomed Inform.* 2014;49:292-9.
6. Friedman DM, Sokal SM, Chang Y, Berger DL. Increasing operating room efficiency through parallel processing. *Ann Surg.* 2006;243(1):10-4.
7. Day LW, Belson D. Studying and Incorporating Efficiency into Gastrointestinal Endoscopy Centers. *Gastroenterol Res Pract.* 2015;2015:764153.
8. Kaushal NK, Chang K, Lee JG, Muthusamy VR. Using efficiency analysis and targeted intervention to improve operational performance and achieve cost savings in the endoscopy center. *Gastrointest Endosc.* 2014;79(4):637-45.
9. Day LW, Belson D, Dessouky M, et al. Optimizing efficiency and operations at a California safety-net endoscopy center: a modeling and simulation approach. *Gastrointest Endosc.* 2014;80(5):762-73.
10. Yong E, Zenkova O, Saibil F, et al. Efficiency of an endoscopy suite in a teaching hospital: delays, prolonged procedures, and hospital waiting times. *Gastrointest Endosc.* 2006;64(5):760-4.
11. Zamir S, Rex DK. An initial investigation of efficiency in endoscopy delivery. *Am J Gastroenterol.* 2002;97(8):1968-72.
12. Berg B, Denton B, Nelson H, et al. A discrete event simulation model to evaluate operational performance of a colonoscopy suite. *Med Decis Making.* 2010;30(3):380-7.
13. Yang D, Summerlee R, Suarez AL, et al. Evaluation of interventional EU efficiency metrics at a tertiary academic medical center. *Endosc Int Open.* 2016;4(2):E143-8.
14. Almeida R, Paterson WG, Craig N, Hookey L. A Patient Flow Analysis: Identification of Process Inefficiencies and Workflow Metrics at an Ambulatory EU. *Can J Gastroenterol Hepatol.* 2016;2016:2574076.
15. Weinbroum AA, Ekstein P, Ezri T. Efficiency of the operating room suite. *Am J Surg.* 2003;185(3):244-50.
16. Overdyk FJ, Harvey SC, Fishman RL, Shippey F. Successful strategies for improving operating room efficiency at academic institutions. *Anesth Analg.* 1998;86(4):896-906.
17. Bjorkman DJ. Efficiency in endoscopy: imperatives and pitfalls. *Gastrointest Endosc.* 2008;68(6):1054-5.

18. Gellad ZF, Thompson CP, Taheri J. EU efficiency: quality redefined. *Clin Gastroenterol Hepatol*. 2013;11(9):1046-9 e1.
19. Rex DK, Lahue BJ, Dronzek RW. Impact of two procedure rooms per physician on productivity and computer simulation examines the impact of process change in the hospital gastroenterology department. *Gastrointest Endosc*. 2005;61:AB154.
20. Harewood GC, Chrysostomou K, Himy N, Leong WL. A "time-and-motion" study of endoscopic practice: strategies to enhance efficiency. *Gastrointest Endosc*. 2008;68(6):1043-50.



Table 1: Comparison of baseline times at LAC+USC to benchmarks with possible causes and interventions

	Monitored	anesthesia care	p-value		Moderate sedation	p-value		
Metric	LAC+USC	Benchmark*		LAC+USC	Benchmark*		Explanation	Interventions
Room turnover time	45.3 (31.8-58.8)	26.6 ¹	<0.01*	25.2 (19.7-30.7)	26.6 ¹	0.6	Anesthesia evaluations by CRNA performed between procedures	PAT clinic, Streamline inter-procedure processes
First case on-time start	9 (0-26)%	64.5% ¹	<0.01*	33 (20-47)%	64.5% ¹	<0.01*	Patient prep outside room Same-day anesthesia evaluation	Pre-op 1 st patient in room Double schedule early cases First case delay log
In room to procedure start time	16.0 (12.5-19.4)	20.8 ²	<0.01**	25.6 (23.2-27.9)	33.7 ²	<0.01*	Benchmark met	None
Procedure time	31.6 (26.7-36.4)	38.4 ²	<0.01**	36.6 (33.4-39.9)	31.1 ²	<0.01*	Benchmark met	None
Procedure end to room exit time	12.4 (10.6-14.2)	13 ²	0.5	16.5 (15.3-17.6)	9.4 ²	<0.01**	Detailed report required before patient leaves procedure room	Brief procedure note Re-time procedure note

*tested for equality with benchmark mean or proportion at 95% confidence interval, significant p value signifies significant difference with benchmark, did not meet metric ***tested for equality with benchmark mean or proportion at 95% confidence interval, significant p value signifies significant difference with benchmark, performance exceeded metric

- 1) Day L, Belson D, Gastroenterology Research and Practice 2015; 764153
- 2) Kaushal, K, Chang K, Lee J, Muthusamy V. Gastrointestinal Endoscopy 2014; 79(4): 637-645

Table 2: Core productivity and efficiency parameters before and after PDSA guided intervention

	Pre-intervention	Post-intervention	Major Changes Implemented*	Statistical Significance
	Mean (+SD)	Mean (+SD)		Mean difference (95% CI)
Total Volume (Per Month)	495.8 (+40.7)	583.0 (+73)	All	87.2 (0.1 to 174)
Anesthesia Room Volume (Per Day)	5.6 (2.1)	8.3 (2.1)	Pre-anesthesia clinic visit (PAT)*	2.8 (0.9-4.7)
Room turnover time (MAC)(Minutes)	45.3 (±41)	29.8 (±23.5)		-15.5 (-3.9 to -27.1)
Room turnover time (Moderate Sedation)(Minutes)	24.5 (+36.7)	22.0 (+32.1)	Parallel patient preparation by RN and MD**, PAT*	-2.4(-8.7 to 3.8)
Scope out to out of room (Moderate Sedation)	16.5 (8.1)	14.4 (.6.8)	Brief procedure note**	-2.1 (-3.3 to -0.8)
	N(%)	N(%)	Phase	OR
First-case on time start % (anesthesia supported)	20%	87.5%	Pre-op 1 st patient in procedure room**	28 (2.6-297.9)
First-case on time start % (moderate Sedation)	29.3%	60.0%	Double schedule early cases to mitigate impact of no shows**	3.6 (1.7-7.8)

*Interventions implemented in phase 2. **Interventions implemented in phase 3

Figure 1) PDSA Ramp




Figure 2A) Process Flow of Endoscopy Unit





Figure 3) Stakeholder Survey



Time Flow Data Sheet

Date: _____

Check in with Clerk

Time at Check in: _____

GI Nurse Assessment

Time: _____

MAC? ☐ Y ☐ N

CRNA Nurse Assessment

Time: _____

Procedure Room

In Room	Room # _____
Scope In	Patient # _____
End Time	<input type="checkbox"/> EGD
Out of Room	<input type="checkbox"/> Colonoscopy
	<input type="checkbox"/> Other _____

Recovery

Time: _____

Discharge

Time: _____

Patient MRN/Sticker

Scope Tracking Log

RN: _____ Room: _____ Date: _____

Patient Name/MRN Sticker	TIME
<input type="checkbox"/> EGD <input type="checkbox"/> Colonoscopy <input type="checkbox"/> Flex sigmoid <input type="checkbox"/> Other _____ Scope Serial No: _____ Physician's Name: _____ RN II name: _____	TIME IN ROOM _____ SCOPE-IN _____ END TIME _____ OUT OF ROOM _____
<input type="checkbox"/> EGD <input type="checkbox"/> Colonoscopy <input type="checkbox"/> Flex sigmoid <input type="checkbox"/> Other _____ Scope Serial No: _____ Physician's Name: _____ RN II name: _____	TIME IN ROOM _____ SCOPE-IN _____ END TIME _____ OUT OF ROOM _____
<div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 0 auto;"> <p>TIME</p> <p>TIME IN ROOM</p> <p>SCOPE-IN</p> <p>END TIME</p> <p>OUT OF ROOM</p> </div>	
<input type="checkbox"/> EGD <input type="checkbox"/> Colonoscopy <input type="checkbox"/> Flex sigmoid <input type="checkbox"/> Other _____ Scope Serial No: _____ Physician's Name: _____ RN II name: _____	TIME IN ROOM _____ SCOPE-IN _____ END TIME _____ OUT OF ROOM _____