

Journal of Neurological Surgery Reports

Nuanced Management of a Skull Base Tumor in the Setting of Relapsed Acute Lymphoblastic Leukemia

Geena Jung, Emery Buckner-Wolfson, Adit Tal, Ryan Fatemi, Timothy Kim, Genesis Liriano, Andrew Kobets.

Affiliations below.

DOI: 10.1055/a-2297-4265

Please cite this article as: Jung G, Buckner-Wolfson E, Tal A et al. Nuanced Management of a Skull Base Tumor in the Setting of Relapsed Acute Lymphoblastic Leukemia. Journal of Neurological Surgery Reports 2024. doi: 10.1055/a-2297-4265

Conflict of Interest: The authors declare that they have no conflict of interest.

Abstract:

Introduction: Relapsed acute lymphoblastic leukemia involving the central nervous system (CNS) is a significant issue that contributes to both morbidity and mortality. Given the poor outcomes in patients with CNS relapse, understanding how ALL involving intracranial relapse presents and is treated is critical. Here, we present a complex case of relapsed recurrent ALL in a pediatric patient.

Case Report: An eleven-year-old patient presented with double relapse of ALL in the form of an extensive skull base lesion and again with leptomeningeal disease. For the skull base lesion, she was treated non-surgically with chemotherapy and radiation, which led to a remarkable reduction in the size of the lesion. However, she was found to have early recurrence with leptomeningeal enhancement resulting in hydrocephalus five months after completing therapy. A shunt was placed successfully. Currently, she is being managed with monthly intrathecal chemotherapy with CSF sampling and bone marrow biopsies every two months.

Discussion: We report the significant effect of chemotherapy and radiotherapy in reducing the size of the extensive skull base lesion, saving the patient from the risks associated with surgery. This patient's initial relapse, with a large skull base lesion that had intracranial involvement, is an unusual presentation of relapsed ALL. The additional early recurrence of leptomeningeal disease further makes this case unique and the management even more nuanced. Here, we demonstrate a multidisciplinary approach for the successful treatment of our patient, which can help guide the management of similar patients in the future.

Corresponding Author:

Geena Jung, Albert Einstein College of Medicine, Neurological Surgery, Bronx, United States, geena.jung@einsteinmed.edu

Affiliations:


Geena Jung, Albert Einstein College of Medicine, Neurological Surgery, Bronx, United States


Emery Buckner-Wolfson, Albert Einstein College of Medicine, Neurological Surgery, Bronx, United States


Adit Tal, Montefiore Medical Center, Pediatrics, Bronx, United States

[...]

Andrew Kobets, Montefiore Medical Center, Neurological Surgery, Bronx, United States







Nuanced Management of a Skull Base Tumor in the Setting of Relapsed Acute Lymphoblastic Leukemia

Geena Jung, BA,¹ Emery Buckner-Wolfson, BS,¹ Adit Tal, MD², Ryan Fatemi, BS,¹ Timothy Kim, BS,¹ Genesis Liriano, MD,¹ Andrew Kobets, MD¹

¹Departments of Neurosurgery, Montefiore Medical Center and the Albert Einstein College of Medicine, Bronx, NY, USA

²Department of Pediatrics, Montefiore Medical Center and the Albert Einstein College of Medicine, Bronx, NY, USA

Corresponding Author

Abstract

Introduction: Relapsed acute lymphoblastic leukemia involving the central nervous system (CNS) is a significant issue that contributes to both morbidity and mortality. Given the poor outcomes in patients with CNS relapse, understanding how ALL involving intracranial relapse presents and is treated is critical. Here, we present a complex case of relapsed recurrent ALL in a pediatric patient.

Case Report: An eleven-year-old patient presented with double relapse of ALL in the form of an extensive skull base lesion and again with leptomeningeal disease. For the skull base lesion, she was treated non-surgically with chemotherapy and radiation, which led to a remarkable reduction in the size of the lesion. However, she was found to have early recurrence with leptomeningeal enhancement resulting in hydrocephalus five months after completing therapy. A shunt was placed successfully. Currently, she is being managed with monthly intrathecal chemotherapy with CSF sampling and bone marrow biopsies every two months.

Discussion: We report the significant effect of chemotherapy and radiotherapy in reducing the size of the extensive skull base lesion, saving the patient from the risks associated with surgery. This patient's initial relapse, with a large skull base lesion that had intracranial involvement, is an unusual presentation of relapsed ALL. The additional early recurrence of leptomeningeal disease further makes this case unique and the management even more nuanced. Here, we demonstrate a multidisciplinary approach for the successful treatment of our patient, which can help guide the management of similar patients in the future.

Keywords

Skull base tumor, Acute lymphocytic leukemia, central nervous system involvement, clinical management

Introduction:

Acute lymphocytic leukemia (ALL) is a progressive cancer characterized by the proliferation of immature lymphoid cells in the bone marrow, blood, and extramedullary sites. Typically, most cases of ALL are diagnosed in the pediatric population. In fact, ALL is the most common form of childhood cancer, accounting for up to 25% of cancers in children under the age of 15 years¹. Although childhood ALL is highly curable, relapsed disease is a significant problem that contributes to morbidity and mortality^{2,3}. For patients with relapse, the 5-year survival rate is estimated to be between 25-50%⁴⁻⁸.

The CNS is one of the most common sites of extramedullary ALL relapse and can occur as isolated recurrence or alongside bone marrow involvement, both of which have devastating consequences^{9,10}. Given the poor outcomes in patients with CNS relapse, understanding how ALL recurrence involving intracranial infiltration presents and can be managed is essential.

Here we describe the case of a patient with late isolated CNS relapse of ALL in the form of an extensive skull base lesion, who was effectively treated non-surgically with chemotherapy and radiation, which led to a drastic decrease in the size and intensity of the lesion.

Case Presentation:

An 11-year-old girl with a history of standard-risk B-cell ALL 7 years prior presented with left facial droop, recurrent mastoiditis/otitis, and right elbow swelling. She was first diagnosed with SR-ALL at 18 months of age and was enrolled on study DFCI 11-001. She went into remission and completed 2 years of therapy with minimal complications.

When she presented presently, physical examination was limited due to agitation. She had a notable left facial droop and swelling of her left ear and mastoid with bony prominence, though the area was not tender or erythematous. A peripheral CNVII palsy was observed, without evidence of other cranial nerve dysfunction. She was able to move all extremities and demonstrated normal strength and sensation. The remainder of her exam was noncontributory.

MRI of the brain revealed a large skull base lesion centered on the sphenoid bones and clivus with extensive intracranial, nasopharyngeal, and left neck extension (Figure 1). The mass encapsulated the left VII and VIII cranial nerves, dorsally displacing the pons and medulla, and filling the left cavernous sinus. The lesion was biopsied, and CSF was obtained for cytologic examination. Pathologic evaluation revealed the tumor was consistent with relapsed pre-B ALL

rather than a new second malignancy. Moreover, CSF from the lumbar puncture (LP) showed 15% blast-like cells, with tissue flow from the tumor further being consistent with relapsed pre-B ALL. Notably, the blasts were negative for the B-cell biomarker CD19. The bone marrow biopsy was negative, suggesting late isolated CNS relapse.

She was started on re-induction chemotherapy as per AALL1331, with systemic mitoxantrone, dexamethasone, vincristine, and pegaspargase and weekly intrathecal methotrexate, cytarabine, and hydrocortisone, which she tolerated well (ClinicalTrials.gov identifier: NCT02101853). End of induction lumbar puncture and bone marrow biopsy showed no residual disease. End of induction MRI brain scans, however, showed that while the intracranial lesion decreased significantly, there remained a residual mass centered on the left skull base, with mass effect upon the brainstem. Prolonged systemic chemotherapy was continued as per AALL1331. Surgery was not considered due to the extensive intracranial component of the lesion and the high risk of the procedure. Given the absence of CD19 positivity on the leukemic cells, the patient was also ineligible for CAR-T cell therapy targeting CD19, and no open clinical trials were available. While allogeneic hematopoietic stem cell transplantation was considered, the patient's parents decided to forego the treatment in favor of a targeted radiotherapy and chemotherapy approach.

The patient received chemotherapy for approximately two years as per AALL1331 and targeted radiation (24Gy) to the lesion. Her treatment was complicated by fungal pneumonia. MRI scans of the brain at the end of therapy showed a remarkable decrease in the size and intensity of the skull base lesion as compared to the MRI scan taken two years prior, although her facial droop persisted (Figure 2). Given the duration of treatment for the chemotherapy was exhausted, a discussion at tumor board yielded a decision to monitor the residual lesion and consider biopsy and excision if the mass started to re-grow.

Unfortunately, five months after completing therapy, the patient presented to the hospital with worsening facial droop and neck pain. MRI scans showed the skull base lesion was stable, but there was a new finding of diffuse leptomeningeal enhancement (Figure 3). She subsequently had an initial LP, which showed no blasts but elevated intracranial pressure (ICP) to 41 and a white blood cell count of 35,000/mm³. Repeat LP 1 week later revealed 20% blast cells confirming CNS relapse. Bone marrow biopsy was negative for leukemic infiltration and CT scan further revealed communicating hydrocephalus and possible obstruction from crowding cerebellar tonsils at the level of the foramen magnum. The patient had emergent external ventricular drainage placement for increased ICP and eventually underwent ventriculoperitoneal shunt placement. She also received biweekly intrathecal therapy until the CSF was cleared of blasts.

Currently, she is receiving monthly intrathecal therapy consisting of methotrexate, hydrocortisone, and cytarabine, and a bone marrow biopsy every two months to evaluate for

disease. Given her extensive ALL history and CNS recurrence, bone marrow transplantation was once again recommended. The risks and benefits of transplantation in alignment with the family's wishes has been an ongoing discussion. The patient has been tolerating chemotherapy well and has not had any headaches or emesis. CSF remains negative for leukemia.

Discussion:

Currently, there is no unified treatment plan for CNS relapse in ALL and evaluating the best course of treatment for these patients can be challenging. Depending on the degree of intracranial invasion, strategies typically involve a combination of the following: cranial radiation, conventional chemotherapy, hematopoietic stem cell transplantation, and, more recently, immunotherapies^{11,12}. Early isolated CNS relapses, defined by past literature as relapse <18 months from diagnosis, are often treated with intensive systemic chemotherapy followed by HSCT if patients are eligible^{13,14}. For late isolated CNS relapses, there has been relative success with clinical trials using high dose chemotherapy and cranial radiation¹¹. While the utilization of radiation has improved survival, it is known to have several long-term side effects including cognitive and growth impairments^{15,16}.

Here, we report a case of a young patient with relapsed recurrent ALL involving the CNS who was treated with a non-surgical multidisciplinary approach which drastically reduced the tumor size. The patient's initial relapse with a large skull base lesion and intracranial involvement, is an interesting and uncommon presentation of relapsed CNS-ALL. The patient's early recurrence of leptomeningeal disease is another uncommon presentation of relapsed ALL. Lastly, and most importantly, the patient's radiographic response to standard chemotherapy and radiation was unexpected and profound, highlighting how a patient with relapsed CNS-ALL can be managed non-surgically. Many available treatment options for CNS relapse involve a high risk of toxicity, making such interventions suboptimal in certain patients. Here, however, we demonstrate the profound effect of chemotherapy and radiotherapy in reducing the size of an extensive skull base lesion, saving the patient from the risks associated with surgery.

While our therapeutic approach had a significant impact on the growth of the lesion, complications with this strategy did arise. The early recurrence of the patient's CNS disease, in the form of leptomeningeal enhancement and hydrocephalus, despite intensive chemotherapy, was unfortunate and unexpected. While we believe our course of therapeutic action was the most appropriate given the patient's history of disease and family wishes, it is clear that more effective forms of therapy for CNS-ALL are needed. In order to truly develop an optimal form of therapy, it is crucial to first identify more diagnostic and prognostic biomarkers for relapsed CNS-ALL. Such an endeavor would allow for the identification of patients who are at high risk for CNS relapse as well as the development of more targeted therapies to better manage these patients in the future.

In summary, we report a unique case of an eleven-year-old girl with a history of SR-ALL who presented with relapsed CNS-ALL in the form of a large skull base lesion with intracranial involvement. The patient's initial relapse, additional early recurrence of leptomeningeal disease, and response to standard chemotherapy and radiation makes our case particularly unusual. We demonstrate the intricate decision-making process behind optimizing a treatment plan for such a nuanced case of CNS-ALL and show how similar patients can be managed without the use of surgical interventions. Moving forward, we believe it is important to be able to identify patients

at risk of relapsed CNS-ALL, as this would allow for swift clinical action to be taken before progression of disease. Through early identification as well as the development of more targeted therapies for intracranial disease, we will be able to better manage and treat patients with relapsed ALL involving the CNS.

Abbreviations

ALL - Acute lymphocytic leukemia, B-ALL - B-cell acute lymphocytic leukemia, SR-ALL – Standard risk acute lymphocytic leukemia, LP – lumbar puncture, EVD - external ventricular drainage, VP – ventriculoperitoneal, IT – intrathecal, HSCT – hematopoietic stem cell transplant

Declarations

Ethics approval and consent to participate: Approval for the publication of this work was granted by the Albert Einstein IRB and consent to publish this report was obtained from the patient's family.

Consent for publication: Consent for publication was obtained from the patient's family.

Availability of data and materials: Not applicable

Author's contributions: GJ, EBW, RF, TK were involved in the composition of this manuscript and AT, GL, MB, AK were involved in revision and review of this manuscript. GJ and AK developed the concept for this report.

Author's information: as outlined above

Acknowledgement

None

References

1. Kakaje A, Alhalabi MM, Ghareeb A, Karam B, Mansour B, Zahra B, Hamdan O. Rates and trends of childhood acute lymphoblastic leukaemia: an epidemiology study. *Scientific reports*. 2020 Apr 21;10(1):1-2.
2. Place AE, Stevenson KE, Vrooman LM, Harris MH, Hunt SK, O'Brien JE, Supko JG, Asselin BL, Athale UH, Clavell LA, Cole PD. Intravenous pegylated asparaginase versus intramuscular native Escherichia coli L-asparaginase in newly diagnosed childhood acute lymphoblastic leukaemia (DFCI 05-001): a randomised, open-label phase 3 trial. *The lancet oncology*. 2015 Dec 1;16(16):1677-90.
3. Vora A, Goulden N, Wade R, Mitchell C, Hancock J, Hough R, Rowntree C, Richards S. Treatment reduction for children and young adults with low-risk acute lymphoblastic leukaemia defined by minimal residual disease (UKALL 2003): a randomised controlled trial. *The lancet oncology*. 2013 Mar 1;14(3):199-209.
4. Hunger SP, Lu X, Devidas M, Camitta BM, Gaynon PS, Winick NJ, Reaman GH, Carroll WL. Improved survival for children and adolescents with acute lymphoblastic leukemia between 1990 and 2005: a report from the children's oncology group. *Journal of clinical oncology*. 2012 May 5;30(14):1663.
5. Pui CH, Yang JJ, Hunger SP, Pieters R, Schrappe M, Biondi A, Vora A, Baruchel A, Silverman LB, Schmiegelow K, Escherich G. Childhood acute lymphoblastic

leukemia: progress through collaboration. *Journal of Clinical Oncology*. 2015 Sep 9;33(27):2938.

6. Hunger SP, Raetz EA. How I treat relapsed acute lymphoblastic leukemia in the pediatric population. *Blood*. 2020 Oct 15;136(16):1803-12.
7. Nguyen K, Devidas M, Cheng SC, La M, Raetz EA, Carroll WL, Winick NJ, Hunger SP, Gaynon PS, Loh ML. Factors influencing survival after relapse from acute lymphoblastic leukemia: a Children's Oncology Group study. *Leukemia*. 2008 Dec;22(12):2142-50.
8. Brown PA, Ji L, Xu X, Devidas M, Hogan LE, Borowitz MJ, Raetz EA, Zugmaier G, Sharon E, Bernhardt MB, Terezakis SA. Effect of postreinduction therapy consolidation with blinatumomab vs chemotherapy on disease-free survival in children, adolescents, and young adults with first relapse of B-cell acute lymphoblastic leukemia: a randomized clinical trial. *Jama*. 2021 Mar 2;325(9):833-42.
9. Aldoss I, Al Malki MM, Stiller T, Cao T, Sanchez JF, Palmer J, Forman SJ, Pullarkat V. Implications and management of central nervous system involvement before allogeneic hematopoietic cell transplantation in acute lymphoblastic leukemia. *Biology of Blood and Marrow Transplantation*. 2016 Mar 1;22(3):575-8.
10. Larson RA. Managing CNS disease in adults with acute lymphoblastic leukemia. *Leukemia & Lymphoma*. 2018 Jan 2;59(1):3-13.
11. Thastrup, M., Duguid, A., Mirian, C., Schmiegelow, K., & Halsey, C. (2022). Central nervous system involvement in childhood acute lymphoblastic leukemia: Challenges and solutions. *Leukemia*, 36(12), 2751-2768.
12. Paul, S., & Short, N. J. (2022). Central nervous system involvement in adults with acute leukemia: Diagnosis, prevention, and management. *Current oncology reports*, 24(4), 427-436.
13. Masurekar AN, Parker CA, Shanyinde M, Moorman AV, Hancock JP, Sutton R, Ancliff PJ, Morgan M, Goulden NJ, Fraser C, Hoogerbrugge PM. Outcome of central nervous system relapses in childhood acute lymphoblastic leukaemia—prospective open cohort analyses of the ALLR3 trial. *PLoS One*. 2014 Oct 3;9(10):e108107.
14. Locatelli F, Schrappe M, Bernardo ME, Rutella S. How I treat relapsed childhood acute lymphoblastic leukemia. *Blood, The Journal of the American Society of Hematology*. 2012 Oct 4;120(14):2807-16.
15. Hastings C, Chen Y, Devidas M, Ritchey AK, Winick NJ, Carroll WL, Hunger SP, Wood BL, Marcus RB, Barredo JC. Late isolated central nervous system relapse in childhood B-cell acute lymphoblastic leukemia treated with intensified systemic therapy and delayed reduced dose cranial radiation: A report from the Children's Oncology Group study AALL02P2. *Pediatric blood & cancer*. 2021 Dec;68(12):e29256.
16. Halsey, C., Buck, G., Richards, S., Vargha-Khadem, F., Hill, F., & Gibson, B. (2011). The impact of therapy for childhood acute lymphoblastic leukaemia on intelligence

quotients; results of the risk-stratified randomized central nervous system treatment trial MRC UKALL XI. *Journal of hematology & oncology*, 4(1), 1-12.

Figure 1: Sagittal (A) and coronal (B) T1 weighted MRI from 2020 of the patient's initial relapse prior to therapy demonstrating a large skull base lesion centered on the sphenoid bones and clivus.

Figure 2: Sagittal (A) and coronal (B) T1 weighted MRI from 2022 following the completion of chemotherapy and radiotherapy demonstrating the reduction in the size of the skull base lesion.

Figure 3: Coronal MRI (A) and axial CT (B) scans five months after completing therapy depicting a stable skull base lesion and new leptomeningeal enhancement.