

Intramural hematoma: a rare complication of endoscopic injection therapy for bleeding peptic ulcers

Intramural hematomas of the gastrointestinal tract are a rare entity. We report two cases of intramural hematoma that developed following endoscopic therapy for bleeding ulcers.

Case 1: A 67-year-old man with diabetes was admitted to the intensive care unit for acute respiratory distress syndrome after a bout of pneumonia. He was being treated with prednisolone, ranitidine, and enoxaparin. Fourteen days later, his hemoglobin decreased from 9.6 g/dL to 6.7 g/dL and he was transfused with 3 units of packed red blood cells (with the hemoglobin rising to 10.2 g/dL). Endoscopy revealed oozing bleeding from an ulcer on the anterior wall of the duodenal bulb. Hemostasis was achieved by injecting 5 mL of diluted epinephrine (1 : 10000) and 1 mL of absolute alcohol. However, after 3 days, the patient's hemoglobin decreased again (from 9.1 g/dL to 7.8 g/dL). A second-look endoscopy showed a conspicuous, violet-colored bulge in the duodenum, which seemed to be an intramural hematoma (• [Video 1](#)).

This was confirmed by computed tomography (• [Fig. 1](#)).

Conservative treatment was instituted, but 20 hours later, the patient developed acute abdomen. An emergency laparotomy revealed a large duodenal hematoma extending into the retroperitoneum, with necrosis of the posterior wall (• [Fig. 2](#)).

The hematoma was drained and there were no surgical complications. However, the patient died 30 days later from respiratory failure.

Case 2: A 76-year-old man was admitted with angina and melena since 2 days. He had a history of myocardial infarction, which had been treated with clopidogrel and acetylsalicylic acid. His hemoglobin was 11.8 g/dL. Endoscopy revealed an



Fig. 1 Computed tomography scan showing an inhomogeneous, hyperdense mass, 10 × 7.5 cm in size, in the duodenum wall.




Fig. 2 At surgery, a large duodenal hematoma was visualized, with necrosis of the posterior wall and extending into retroperitoneum.



Fig. 3 Second-look endoscopy – 24 hours after hemostasis – showing three violet-colored bulges in the posterior wall of the antrum, corresponding to intramural hematomas.



Fig. 4 Endoscopy, 6 days later, showing a longitudinal ulcer and complete resolution of the hematomas.

Video 1

Endoscopy showing oozing bleeding from an ulcer on the anterior wall of the duodenal bulb; hemostasis was achieved with 5 mL of diluted epinephrine (1 : 10000) along with 1 mL of absolute alcohol. Three days later, endoscopy showed a prominent violet-colored bulge, corresponding to an intramural hematoma.

oval ulcer on the posterior wall of the gastric antrum, with oozing bleeding. Hemostasis was achieved by injecting 4 mL of diluted epinephrine (1:10000) and 0.5 mL of absolute alcohol. A second-look endoscopy, 24 hours later, revealed three intramural hematomas (● Fig. 3).

Another endoscopy 6 days later showed complete resolution of the hematomas (● Fig. 4).

The cause of most intramural hematomas is blunt abdominal trauma; however, they have also been reported to occur as a complication of anticoagulant therapy [1] and blood dyscrasias [2], and after endoscopic biopsy [3] or therapy [4,5]. In this latter case, the use of large amounts of injected substances and antiplatelet/anticoagulation therapy may favour their development [4]. We believe that in our cases, the presence of comorbidities in the first patient and the excessive anti-aggregation therapy in the second patient were implicated in the development of the intramural hematomas.

Endoscopy_UCTN_Code_CPL_1AH_2AC

Competing interests: None

**A. Sadio¹, P. Peixoto¹, E. Cancela¹,
A. Castanheira¹, V. Marques²,
P. Ministro¹, A. Silva¹, A. Caldas¹**

¹ Department of Gastroenterology,
São Teotónio Hospital, Viseu, Portugal

² Department of Surgery 1, São Teotónio
Hospital, Viseu, Portugal

References

- 1 Chaiteerakij R, Treeprasertsuk S, Mahachai V et al. Anticoagulant-induced intramural intestinal hematoma: report of three cases and literature review. *J Med Assoc Thai* 2008; 91: 1285–1290
- 2 Nogues A, Eizaguirre I, Sunol M et al. Giant spontaneous duodenal hematoma in hemophilia A. *J Pediatr Surg* 1989; 24: 406–408
- 3 Ghisban FK, Werner M, Vieira P et al. Intramural duodenal hematoma: an unusual complication of endoscopic small bowel biopsy. *Am J Gastroenterol* 1987; 82: 368–370
- 4 Rohrer B, Schreiner J, Lehnert P et al. Gastrointestinal intramural hematoma, a complication of endoscopic injection methods for bleeding peptic ulcers: a case series. *Endoscopy* 1994; 26: 617–621
- 5 Sugai K, Kajimara E, Mochizuki Y et al. Intramural duodenal hematoma after endoscopic therapy for a bleeding duodenal ulcer in a patient with liver cirrhosis. *Intern Med* 2005; 44: 954–957

Bibliography

DOI 10.1055/s-0029-1214927

Endoscopy 2011; 43: E141–E142

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

A. A. D. Sadio

Department of Gastroenterology

São Teotónio Hospital

Av. Rei D. Duarte 3504 509

Viseu

Portugal

Fax: +351-232-420591

anasadio@iol.pt