

Lymphoepithelioma-like gastric carcinoma resected by endoscopic submucosal dissection (ESD)




Fig. 1 Endoscopy showed an erythematous, flat depressed lesion in the posterior wall of the upper body of the stomach.

We report here a rare case of lymphoepithelioma-like gastric carcinoma that presented as a flat depressed lesion and was treated by complete en-bloc resection using endoscopic submucosal dissection (ESD).

A 73-year-old man was referred to the department of internal medicine because of an incidental finding suspicious of early gastric cancer. The patient's medical history and laboratory test results were unremarkable. Gastroscopy revealed an erythematous, flat depressed lesion in the posterior wall of the upper part of the gastric body (● Fig. 1).

No abnormalities were detected on abdominal computed tomography. The tumor was removed completely by ESD, using an insulation-tipped knife (KD-610L; Olympus, Tokyo, Japan) (● Fig. 2). Histopathological findings of the resected specimen were compatible with the diagnosis of lymphoepithelioma-like gastric carcinoma (● Fig. 3).

Lymphoepithelioma-like gastric carcinoma is a rare type of gastric carcinomas [1]. It is characterized by the presence of a lymphoid stroma and small nests of cancer cells that are uniformly distributed throughout the lymphoid stroma. There is clear demarcation between the tumor nests and the nondescript lymphocyte-rich stroma, as is the case with lymphoid tissue. Lymphoepithelioma-like gastric carcinoma has a good prognosis and is closely associated with the presence of the Epstein-Barr virus and microsatellite instability [2]. The recently introduced technique of ESD




Fig. 2 **a** The lesion was resected by endoscopic submucosal dissection (ESD) using an insulation-tipped knife. **b** The resected tumor was 19 × 14 mm in size.

can be useful in the diagnosis and treatment of early gastric cancer. In particular, this method enables en-bloc resection of the tumor, regardless of tumor size and location.

In our case, endoscopic examination of the flat depressed lesion revealed an ap-




Fig. 3 **a** Histological examination of the resected specimen revealed that the mass was located in the subepithelial area (hematoxylin and eosin [H&E], magnification × 20). **b** Nests of tumor cells separated by dense lymphoplasmacytic infiltration (H&E, magnification × 200). Immunohistochemical staining was positive for: **(c)** cytokeratin; and **(d)** leucocyte common antigen (LCA; CD 45).

pearance typical of early gastric adenocarcinoma. However, the histological examination revealed that the tumor was a lymphoepithelioma-like gastric carcinoma.

Endoscopy_UCTN_Code_TTT_1AO_2AG

H. S. Moon¹, S. H. Kang¹, J. K. Seong¹,
H. Y. Jeong¹, K. S. Song²

¹ Department of Internal Medicine, Chung Nam University College of Medicine, Daejeon, South Korea

² Department of Pathology, Chung Nam University College of Medicine, Daejeon, South Korea

References

- 1 Herath CH, Chetty R. Epstein-barr virus-associated lymphoepithelioma like gastric carcinoma. Arch Pathol Lab Med 2008; 132: 706–709
- 2 Grogg KL, Lohse CM, Pankratz VS et al. Lymphocyte-rich gastric cancer: association with Epstein-Barr virus, microsatellite instability, histology and survival. Mod Pathol 2003; 16: 641–651

Bibliography

DOI 10.1055/s-0029-1215199

Endoscopy 2010; 42: E73–E74

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

Dr. H. S. Moon

Department of Internal Medicine
Chung Nam University College of Medicine
Dae Sa Dong, 640
Daejeon
South Korea
Fax: +82-42-2544553
mhs1357@cnuh.co.kr