

Appendiceal mucocoele in ulcerative colitis

A 55-year-old woman with ulcerative colitis underwent colonoscopy with a view to assess the activity and extent of disease. During the procedure, a 2.5-cm, smooth, rounded lesion (● Fig. 1) was found in the cecum, distal and separate from the ileocecal valve; the overlying mucosa had a normal appearance.

A contrast computed tomography (CT) scan of the abdomen showed that the appendix was dilated with low-attenuation fluid to 1.5 cm (● Fig. 2; arrow) and showed calcification at its origin and tip. Based on these findings a diagnosis of mucocoele of the appendix was suspected. The patient was referred for laparoscopic resection and a right-sided hemicolectomy was carried out. The surgical specimen included a dilated appendix, 2.8 cm in diameter (● Fig. 3; arrow); sectioning revealed a 1.3-cm spherical tumour. Histological examination showed this to be a mucinous cyst adenoma with low grade dysplasia (● Fig. 4).

Mucocoeles of the appendix are uncommon lesions caused by distension of the appendix with mucinous material. There are four main types: simple or retention mucocoeles, hyperplastic mucocoeles, mucinous cyst adenomas, and mucinous cyst-adenocarcinomas [1]. Mucocoeles of the appendix can be asymptomatic or present with abdominal pain or bleeding, or as a palpable abdominal mass. Rarely they cause intestinal obstruction, intussusception, ureteric obstruction, volvulus, or cutaneous fistulae [2]. The most feared complication is pseudomyxoma peritonei, because of spontaneous or iatrogenic rupture that results in seeding of neoplastic cells in the peritoneal cavity. Colonoscopy, CT, and ultrasound can all help in diagnosis [3]. As it is difficult to differentiate between benign and malignant lesions causing mucocoele, resection is recommended. There have been case reports of appendiceal mucocoele in patients with ulcerative colitis. It has been suggested that appendicular involvement with inflammation, or an associated appendicular neoplasm, in such patients leads to blockage and formation of a mucocoele [4].

Endoscopy_UCTN_Code_CCL_1AD_2AD
Endoscopy_UCTN_Code_CCL_1AD_2AJ




Fig. 1 Colonoscopic view of the lesion.




Fig. 2 Computed tomography (CT) scan of the lesion showing a fluid-filled, dilated appendix (arrow).



Fig. 3 Resected specimen with appendix (arrow).

T. Ghosh, A. Chalmers, C. Verbeke, R. Saunders, S. M. Everett
Leeds General Infirmary, Great George Street, Leeds, UK

References

- 1 Rampone B, Roviello F, Marrelli D, Pinto E. Giant appendiceal mucocoele: report of a case and brief review. *World J Gastroenterol* 2005; 11 (Suppl 30): 4761–4763
- 2 Karakaya K, Barut F, Emre AU et al. Appendiceal mucocoele: case reports and review of current literature. *World J Gastroenterol* 2008; 14 (Suppl 14): 2280–2283
- 3 Madwed D, Mindelzun R, Jeffrey RB Jr. Mucocoele of the appendix: imaging findings. *AJR Am J Roentgenol* 1992; 159 (Suppl 1): 69–72
- 4 Lakatos PL, Gyori G, Halasz J et al. Mucocoele of the appendix: an unusual cause of lower abdominal pain in a patient with ulcerative colitis. A case report and review of literature. *World J Gastroenterol* 2005; 11 (Suppl 3): 457–459



Fig. 4 Histological section of the resected tumour.

Bibliography

DOI 10.1055/s-0029-1243913

Endoscopy 2010; 42: E105 – E106

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

T. Ghosh, MBBS, MRCP

Leeds General Infirmary

Great George Street

Leeds LS1 3EX

UK

ghosh_tilak@yahoo.com