

Ulcer in Barrett's esophagus probably related to severe kyphosis




Fig. 1 A long segment (20 cm) of Barrett's mucosa was found up to 17 cm from the incisor line. A deep longitudinal Barrett's ulcer was observed at 30–34 cm from the incisors.




Fig. 3 Barrett's ulcer healed with partial squamous re-epithelialization after 2 months of proton-pump inhibitor administration.

An 83-year-old man with severe kyphosis and parkinsonism caused by cerebral infarction was admitted to our hospital with a history of episodes of chest discomfort. Endoscopic study revealed a long segment of Barrett's mucosa with large hiatal hernia and a deep ulcer in the Barrett's mucosa (► Fig. 1). No malignancy was proven by endoscopic biopsy of the ulcer. The patient was not receiving any drugs or chemicals. Thus, we diagnosed the ulcer as a "Barrett's ulcer" [1]. Computed tomography showed that the thoracic esophagus was dorsally displaced and the stomach ventrally displaced by kyphosis of the spine. Despite overnight fasting, fluid was retained in the mid-esophagus, which corresponded to the location of the Barrett's ulcer (► Fig. 2a). A barium examination revealed reflux from the stomach to the mid-esophagus around the location of the



Fig. 2 **a** Sagittal CT shows the thoracic esophagus displaced dorsally and the stomach displaced ventrally by kyphosis. Refluxed fluid from the stomach is seen in the region of the ulcer (white arrows) at the lowest part of the esophagus with the patient in the supine position. **b** Barium meal study of the esophagus with the patient semi-upright. A large longitudinal ulcer was found in the mid-esophagus. **c** In the supine position, obvious reflux of barium toward the proximal esophagus and subsequent retention of contrast was observed.



ulcer when the patient was semi-upright and supine (► Fig. 2b,c). The patient had to sleep in the supine position during the night because his ability to turn over in bed was restricted by the parkinsonism. No evidence of *Helicobacter pylori* infection was found in the serological test. Evidently, highly acidic reflux fluid might be retained for a long time in the mid-esophagus, where the ulcer developed.

The ulcer healed completely after treatment with omeprazole 20 mg/day for 2 months (► Fig. 3). The onset mechanism of Barrett's ulcer is thought to be peptic ulceration in Barrett's esophagus due to gastroesophageal reflux [2]. It is suggested that kyphosis increases the severity of hiatal hernia and subsequently induces gastroesophageal reflux [3]. In our hospital all three patients

with a nonmalignant ulcer in a Barrett's segment had either kyphosis or scoliosis. This suggests that nonmalignant Barrett's ulcers may be associated with vertebral deformities including kyphosis in Japanese patients.

Competing interests: None

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AZ

A. Oka¹, Y. Amano², T. Yuki²,
R. Kusunoki¹, N. Ishimura¹, S. Ishihara¹,
Y. Kinoshita¹

¹ Second Department of Internal Medicine, Shimane University School of Medicine, Izumo, Shimane, Japan
² Division of Endoscopy, Shimane University Hospital, Izumo, Shimane, Japan

References

- 1 Komorowski RA, Hogan WJ, Chausow DD. Barrett's ulcer: the clinical significance today. *Am J Gastroenterol* 1996; 91: 2310–2313
- 2 Bremner CG. The columnar-lined (Barrett's) esophagus. *Surg Annu* 1977; 9: 103–123
- 3 Kusano M, Hashizume K, Ehara Y et al. Size of hiatus hernia correlates with severity of kyphosis, not with obesity, in elderly Japanese women. *J Clin Gastroenterol* 2008; 42: 345–350

Bibliography

- DOI** 10.1055/s-0030-1255596
Endoscopy 2010; 42: E286–E287
© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

Y. Amano MD
Division of Endoscopy
Shimane University Hospital
89-1 Enya-cho, Izumo-shi
Shimane 693-8501
Japan
Fax: +81-853-202187
amano@med.shimane-u.ac.jp