

Analysis of cyst fluid obtained by endoscopic ultrasound-guided fine-needle aspiration supporting the diagnosis of a pancreatic neuroendocrine neoplasm

A 28-year-old man presented at the emergency ward for bronchopneumonia with hemoptysis. A computed tomography (CT) scan disclosed an incidental 18-mm-wide lesion in the pancreatic tail that appeared cystic with magnetic resonance imaging, with a thick wall and a solid projection, both contrast-enhanced. Serological tumor markers were in the normal range.

Endoscopic ultrasound (EUS) evaluation showed an oval, protruding mass with a mixed solid and cystic echo structure (● Fig. 1).

Fine-needle aspiration (FNA) produced cystic fluid; two slides were smeared and one was stained with hematoxylin and eosin for rapid on-site evaluation. Part of the fluid was sent to the laboratory for tumor marker analysis, while the remainder was preserved in 95% ethanol for cell block preparation. The observation by the on-site cytopathologist of a small group of cells suspected of being a pancreatic endocrine neoplasm (PEN) (● Fig. 2) prompted the request for analysis of chromogranin A in the cystic fluid.

Cell-block sections showed discohesive epithelial cells with a plasmocytoid appearance, regular nuclear membrane, and finely granular chromatin; immunocytochemistry (ICC) results (positivity for chromogranin A and synaptophysin) confirmed the endocrine differentiation. The proliferation index with Ki-67 was positive in < 1% of neoplastic cells (● Fig. 3).

The final cytological diagnosis of a neuroendocrine tumor was supported by the cyst fluid analysis, showing high levels of chromogranin A (138 ng/mL, normal range 20–100 ng/mL), while amylase and carcinoembryonic antigen were low.

Pancreatic endocrine neoplasms are occasionally manifested as cystic lesions [1–4]. Differential diagnosis of pancreatic cystic neoplasms is significantly enhanced by cyst fluid analysis [5].

To our knowledge, this is the first report that demonstrates a high chromogranin A level in the fluid of a cystic pancreatic neuroendocrine tumor sampled during EUS-guided FNA. This can be a useful diagnostic tool confirming a preoperative diagnosis of PEN, especially in those cases




Fig. 1 Endoscopic ultrasound (EUS) image demonstrating the pancreatic lesion, which appeared oval with a thin and hypoechoic rim and a central, anechoic part; the lesion was 22 × 18 mm.




Fig. 2 A direct smear with Giemsa staining showed few epithelial cells with the cytomorphological features of neuroendocrine tumors: smooth, round nuclei, finely stippled chromatin, and delicate cytoplasm. These bland tumor cells seem to be arranged in rosette-like structures.

where FNA gives little material for traditional cytological and ICC investigations.

Endoscopy_UCTN_Code_TTT_1AS_2AD
Endoscopy_UCTN_Code_CCL_1AF_2AZ_3AB

Competing interests: None

F. Maletta¹, D. Pacchioni¹, P. Carucci², G. Accinelli¹, M. Bruno², F. Brizzi², P. Allegranza², M. Rizzetto², G. Bussoletti¹, C. De Angelis²

¹ Department of Biomedical Science and Oncology, University of Turin, Italy

² Department of Gastrohepatology, University of Turin, Italy

References

- 1 Baker MS, Knuth JL, DeWitt J et al. Pancreatic cystic neuroendocrine tumors: preoperative diagnosis with endoscopic ultrasound and fine-needle immunocytology. *J Gastrointest Surg* 2008; 12: 450–456
- 2 Bordeianou L, Vagefi PA, Sahani D et al. Cystic pancreatic endocrine neoplasms: a distinct tumor type? *J Am Coll Surg* 2008; 206: 1154–1158
- 3 Goh BK, Ooi LL, Tan YM et al. Clinico-pathological features of cystic pancreatic endocrine neoplasms and a comparison with




Fig. 3 Cell block preparation showing numerous cells with a plasmacytoid appearance, regular nuclear membrane, and finely granular chromatin pattern. **a** Hematoxylin and eosin stain. The neuroendocrine differentiation was confirmed by strongly positive immunostaining for **b** chromogranin A and **c** synaptophysin. **d** The proliferative index (Ki-67) was low (< 1%).

their solid counterparts. *Eur J Surg Oncol* 2006; 32: 553–556

- 4 Charfi S, Marcy M, Bories E *et al*. Cystic pancreatic endocrine tumors: an endoscopic ultrasound-guided fine-needle aspiration biopsy study with histologic correlation. *Cancer Cytopathol* 2009; 117: 203–210
- 5 Linder JD, Geenen JE, Catalano MF. Cyst fluid analysis obtained by EUS-guided FNA in the evaluation of discrete cystic neoplasms of the pancreas: a prospective single-center experience. *Gastrointest Endosc* 2006; 64: 697–702

Bibliography

DOI 10.1055/s-0030-1256036

Endoscopy 2011; 43: E34–E35

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

C. De Angelis, MD

S.C. Gastro-Epatologia D.U.

Ospedale Molinette

Corso Bramante 88

10126 Torino

Italy

Fax: +39-011-6335927

eusdeang@hotmail.com