

Endosonographic and elastographic features of a rare epidermoid cyst of an intrapancreatic accessory spleen

Accessory spleens are found in approximately 10% of the general population, of which 16% are intrapancreatic [1]. A previously healthy 49-year-old patient was referred to our tertiary center for further evaluation of a pancreatic mass. She had initially presented to another hospital with nonspecific abdominal pain. An abdominal computed tomography (CT) scan had revealed a solid mass in the tail of the pancreas. Endoscopic ultrasound-guided fine needle aspiration (EUS-FNA) was suggestive of a pancreatic neuroendocrine tumor (PNET).

At our institution, the patient had a normal physical examination with no lymphadenopathy, organomegaly, or a palpable abdominal mass. The patient was referred for EUS-guided tattooing to guide laparoscopic distal pancreatectomy. The EUS revealed a hypoechoic, 2.3-cm mass in the tail of the pancreas. The mass was round and homogeneous, with well-demarcated and sharp borders, and a small cystic component (● Fig. 1).

EUS-elastography showed inhomogeneous hardness as compared with the surrounding tissue (● Fig. 2).

EUS-guided tattooing was carried out by injecting 2.5 mL of sterile, purified carbon particle just proximal to the lesion as the needle was withdrawn to the surface of the pancreas (● Fig. 3).

The patient subsequently underwent laparoscopic spleen-preserving distal pancreatectomy without complications. The tattoo was readily identified and demarcated a precise line of resection. Pathologic examination of the surgical specimen demonstrated a cystic mass within the pancreas. The mass had a well-defined capsule within which was splenic parenchyma and a small cyst lined by a layer of benign squamous epithelium. Pathologic diagnosis was consistent with an epidermoid cyst in an intrapancreatic accessory spleen (IPAS) (● Fig. 4).

None of the 16 reported IPAS cases were diagnosed preoperatively as they are known to be difficult to clinically distinguish from other tumors. The value of EUS-FNA for their diagnosis needs further study. FNA was not done in the current case because the lesion was presumed to




Fig. 1 Endoscopic ultrasound showing a hypoechoic, round, and well-circumscribed mass in the tail of the pancreas.




Fig. 2 Endoscopic ultrasound elastography showing inhomogeneous hardness (blue = significant hardness; green = intermediate hardness).



Fig. 3 Endoscopic ultrasound tattooing: injection of purified carbon particle just proximal to the mass as the needle is withdrawn to the surface of the pancreas.




Fig. 4 Epidermoid cyst in an intrapancreatic accessory spleen. Note the thin epithelial lining (top), the layer of splenic tissue, and the associated pancreatic parenchyma (bottom).

represent a PNET according to outside pathologic diagnosis.

Elastography is a means of measuring tissue stiffness. Malignant tissue is harder than benign tissue and elastography may be able to differentiate between them [2]. The system is set up to use a hue color map (red-green-blue), in which hard tissue areas are shown in dark blue, medium-hard tissue areas in cyan, intermediate hardness tissue areas in green, medium-soft tissue areas in yellow, and soft tissue areas in red [3]. In the current case, the epidermoid cyst demonstrated inhomogeneous hardness (mixture of blue and green). EUS-elastography has been used for the diagnosis of pancreatic cancer and malignant lymphadenopathy with variable sensitivity, specificity, and accuracy in different studies [3–5].

Endoscopy_UCTN_Code_CCL_1AF_2AZ_3AB

Competing interests: None

**M. A. Khashab¹, M. I. Canto¹,
V. K. Singh¹, R. H. Hruban²,
M. A. Makary³, S. Giday¹**

¹ Department of Medicine and Division of Gastroenterology and Hepatology, Sol Goldman Pancreatic Cancer Research Center, Johns Hopkins Medical Institutions, Baltimore, Maryland, USA

² Department of Pathology, Sol Goldman Pancreatic Cancer Research Center, Johns Hopkins Medical Institutions, Baltimore, Maryland, USA

³ Department of Surgery, Sol Goldman Pancreatic Cancer Research Center, Johns Hopkins Medical Institutions, Baltimore, Maryland, USA

References

- 1 Halpert B, Alden ZA. Accessory spleens in or at the tail of the pancreas. A survey of 2 700 additional necropsies. *Arch Pathol* 1964; 77: 652–654
- 2 Gao L, Parker KJ, Lerner RM et al. Imaging of the elastic properties of tissue – a review. *Ultrasound Med Biol* 1996; 22: 959–977
- 3 Iglesias-Garcia J, Larino-Noia J, Abdulkader I et al. EUS elastography for the characterization of solid pancreatic masses. *Gastrointest Endosc* 2009; 70: 1101–1108
- 4 Saftoiu A, Vilman P, Ciurea T et al. Dynamic analysis of EUS used for the differentiation of benign and malignant lymph nodes. *Gastrointest Endosc* 2007; 66: 291–300
- 5 Saftoiu A, Vilman P, Gorunescu F et al. Neural network analysis of dynamic sequences of EUS elastography used for the differential diagnosis of chronic pancreatitis and pancreatic cancer. *Gastrointest Endosc* 2008; 68: 1086–1094

Bibliography

DOI 10.1055/s-0030-1256272

Endoscopy 2011; 43: E193–E194

© Georg Thieme Verlag KG Stuttgart · New York ·

ISSN 0013-726X

Corresponding author

S. Giday

Center for Digestive Health

1817 North Mills Avenue

Orlando

Florida 32803

USA

samgiday@gmail.com