

Cyanoacrylate glue in the management of blue rubber bleb nevus syndrome



Fig. 1 Case 1. Hemangiomas of the foot.



Fig. 2 Case 1. Capsule endoscopy picture of a bluish, nipple-like rubbery vascular lesion of the jejunum, compatible with a blue rubber bleb nevus.





Fig. 3 a Case 1. Hemangioma of the proximal jejunum. b Solidification of the glue inside the lumen. Inadvertent sticking of the needle inside the lesion or adherence of the glue to the needle was avoided by flushing at least 2 ml water for injection after each aliquot of Glubran 2 and Lipiodol. c Typical lesion of the transverse colon.

Case 1. A 16-year-old boy with blue rubber bleb nevus syndrome (BRBNS) was admitted for occult bleeding. He had a history of gastric and colonic hemangiomas treated 2 years earlier by argon plasma coagulation and alcohol injection respectively. Clinical examination showed a few hemangiomas on his back and feet (● Fig. 1).

Capsule endoscopy (Pillcam SB2; Given Imaging, Yoqneam, Israel) disclosed three blebs of the small bowel (● Fig. 2), and the patient underwent double-balloon enteroscopy (Fujinon, Saitama, Japan).

A 2-cm bleb was found at the jejunum, two smaller lesions at the ileum, and one at the transverse colon. N-butyl-2-cyanoacrylate and methacryloxysulfolane (Glubran 2), 1 ml, diluted with Lipiodol, 1 ml, was injected in aliquots of 1 ml per injection and up to 4 ml at each lesion (● Fig. 3). No complications occurred. At 6-month follow-up the patient had no anemia.

Case 2. An 11-year-old girl with BRBNS was addressed for hematochezia due to colonic blebs. Colonoscopy revealed a bleeding lesion at the cecum (● Fig. 4). After initial failure to stop the bleeding with a hemostatic grasper (Coagrasper, Olympus), Glubran 2 was injected successfully. Two more blebs of the right colon were treated similarly. After 2 years the patient remained asymptomatic.




Fig. 4 Case 2. Bleeding hemangioma of the cecum near the ileocecal valve.

BRBNS is a rare condition characterized by multiorgan cavernous hemangiomas. Gastrointestinal bleeding is the major problem, and several endoscopic therapeutic approaches have been proposed: Nd:YAG laser, bipolar or argon plasma coagulation, band ligation, snare resection, and sclerotherapy [1–4]. Our patients were treated successfully by Glubran 2 injection, commonly used in the management of gastric varices. Endoscopic obliteration of cavernous hemangiomas with

n-butyl-2-cyanoacrylate (Histoacryl) has been previously described, in the rectum [5]. However, these appear to be the first cases of cyanoacrylate glue injection to control bleeding in the small bowel and colon in the setting of BRBNS.

Endoscopy_UCTN_Code_TTT_1AO_2AD

Competing interests: None

**G. Mavrogenis¹, D. Coumaros¹,
D. Tzilves¹, E. Rapti¹, G. Stefanidis²,
J. Leroy³, F. Becmeur⁴**

¹ Department of Gastroenterology, University Hospital, Strasbourg, France

² Department of Gastroenterology, Athens Naval Hospital, Athens, Greece

³ Department of Digestive Surgery, University Hospital, Strasbourg, France.

⁴ Department of Pediatric Surgery, University Hospital, Strasbourg, France

References

- 1 *Shahed M, Hagenmüller F, Rösch T et al.* A 19-year-old female with blue rubber bleb nevus syndrome. Endoscopic laser photocoagulation and surgical resection of gastrointestinal angiomata. *Endoscopy* 1990; 22: 54–56

- 2 *Maunoury V, Turck D, Brunetaud JM et al.* Blue rubber bleb nevus syndrome. 3 cases treated with a Nd:YAG laser and bipolar electrocoagulation [in French with English abstract]. *Gastroenterol Clin Biol* 1990; 14: 593–595

- 3 *Bak YT, Oh CH, Kim JH.* Blue rubber bleb nevus syndrome: endoscopic removal of the gastrointestinal hemangiomas. *Gastrointest Endosc* 1997; 45: 90–92

- 4 *Arguedas MR, Wilcox CM.* At the focal point. Blue rubber bleb nevus syndrome. *Gastrointest Endosc* 1999; 50: 544

- 5 *Zurakowski J, Swiercz P, Wróblewski T et al.* Diffuse cavernous hemangioma of rectosigmoid colon treated with n-butyl-2-cyanoacrylate injections. *Endoscopy* 2008; 40: E120–121

Bibliography

DOI 10.1055/s-0030-1256425

Endoscopy 2011; 43: E291–E292

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

D. Coumaros, MD

IRCAD/EITS

University Hospital

1 Place de l'Hôpital

67091 Strasbourg

France

Fax: +33-3-88751521

coumarosd@wanadoo.fr