

SYNLETT Spotlight 342

N-Sulfinyl Imines

Compiled by Ping Song

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Ping Song was born in Taizhou, Jiangsu, P. R. of China. She received her B.Sc. degree in Pharmaceutical Engineering from the Nanjing University of Technology, P. R. of China. She is currently working as a postgraduate at the same university under the supervision of Prof. Zhenjiang Li. Her current research is focused on the asymmetric version of the Morita–Baylis–Hillman reaction.

State Key Laboratory of Materials-Oriented Chemical Engineering, College of Biotechnology and Pharmaceutical Engineering, Nanjing University of Technology, Nanjing 210009, P. R. of China
E-mail: zjli@njut.edu.cn




Introduction

N-Sulfinyl imines (sulfonimines) play an important role in asymmetric reactions. They display unique reactivity and stereoselectivity in the synthesis of amino group containing natural products and bioactive compounds. Furthermore, a wide variety of sulfonimines is efficiently prepared for many types of asymmetric reactions, and the chiral sulfinyl in the resultant product is easily removed under comparatively mild conditions.¹

Preparation


Three synthetic routes were developed, for example, asymmetric oxidation,² iminolysis of sulfinate esters,³ and condensation of a sulfonamide with aldehydes or ketones.⁴ The most common and versatile method is the direct condensation of aldehydes or ketones with sulfonamide.




Scheme 1

Abstracts


(A) Sulfonimines were used to prepare functionalized amines with high stereoselectivity. Organometallic reagents, such as Grignard reagents^{5a} and organolithium,⁵ are added to sulfonimines to get the desired products in high diastereoselectivity. Both aliphatic and aromatic sulfonimines proceeded in very high diastereoselective ratios when arylboronic acid was employed.⁶




(B) The sulfonimines-mediated asymmetric Strecker reaction provided efficiently chiral α -amino acids.⁷ Polyhydroxy α -amino acids were derived from polyhydroxy sulfonimines through smooth deprotection of the sulfinyl.⁸ Quaternary α -stereogenic centers of α -amino acids were controlled by tuning the solvents.⁹ The (*S,R*)-product was afforded predominantly in hexane while the contrary (*R,R*)-isomer was the major product in DMF.



(C) β -Amino esters or acids were prepared efficiently via addition of the sodium enolate of methyl acetate to sulfonimines in high diastereoselectivity.^{10a} Lithium enolate effected the better yield.^{10b} The stereoselective Michael–nucleophilic addition domino reaction from sulfonimines was another route towards β -amino esters.¹¹



(D) β -Amino ketones were prepared by addition of prochiral lithium enolates of Weinreb amides to sulfonimines.^{5,12} Reduction of *N*-sulfinyl β -amino ketones led to *syn*- and *anti*-1,3-amino alcohols.¹²




SYNLETT 2011, No. 2, pp 0287–0288

Advanced online publication: 05.01.2011


DOI: 10.1055/s-0030-1259277; Art ID: V34910ST

© Georg Thieme Verlag Stuttgart · New York


(E) The aza-Diels–Alder reactions of sulfinimines as dienophile with Rawal dienes resulted in dihydropyridones with ee values up to 90%.^{13a} When a Lewis acid catalyst was added, both activated and non-activated dienes could be used in this reaction.^{13b}



(F) The pure sulfinimines have been applied in the aza-Baylis–Hillman reaction. The resulting allylic amines reacted with electrophiles led to highly functionalized 3-sulfinyl and 3-sulfonyl 2,5-*cis*-dihydropyrroles.¹⁴



(G) Addition of suitably protected α -amino acid to pure sulfinimines led to *syn*- and *anti*- α,β -diamino esters with high dr and good yields.¹⁵ The water content in THF was an important factor determining the selectivity.^{15b}



References

- (1) (a) Zhou, P.; Chen, B. C.; Davis, F. A. *Tetrahedron* **2004**, *60*, 8003. (b) Morton, D.; Stockman, R. A. *Tetrahedron* **2006**, *62*, 8869.
- (2) Davis, F. A.; Reddy, R. E.; Szweczyk, J. M.; Reddy, G. V.; Portonovo, P. S.; Zhang, H.; Fanelli, D.; Reddy, R. T.; Zhou, P.; Carroll, P. J. *J. Org. Chem.* **1997**, *62*, 2555.
- (3) Davis, F. A.; Reddy, T.; Reddy, R. E. *J. Org. Chem.* **1992**, *57*, 6387.
- (4) (a) Cogan, D. A.; Liu, G.; Kim, K.; Backes, B. J.; Ellman, J. A. *J. Am. Chem. Soc.* **1998**, *120*, 8011. (b) Weix, D. A.; Ellman, J. A. *Org. Lett.* **2003**, *5*, 1317.
- (5) (a) Cogan, D. A.; Liu, G.; Ellman, J. A. *Tetrahedron* **1999**, *55*, 8883. (b) Davis, F. A.; Song, M. *Org. Lett.* **2007**, *9*, 2413.
- (6) Weix, D. J.; Shi, Y.; Ellman, J. A. *J. Am. Chem. Soc.* **2005**, *127*, 1092.
- (7) (a) Davis, F. A.; Zhou, P.; Reddy, G. V. *J. Org. Chem.* **1994**, *59*, 3243. (b) Hua, D. H.; Miao, S. W.; Chen, J. S.; Iguchi, S. *J. Org. Chem.* **1991**, *56*, 4. (c) Hua, D. H.; Lagneau, N.; Wang, H.; Chen, J. *Tetrahedron: Asymmetry* **1995**, *6*, 349.
- (8) Davis, F. A.; Prasad, K. R.; Carroll, P. J. *J. Org. Chem.* **2002**, *67*, 7802.
- (9) Wang, H.; Zhao, X.; Li, Y.; Lu, L. *Org. Lett.* **2006**, *8*, 1379.
- (10) (a) Davis, F. A.; Reddy, R. E.; Szweczyk, J. M. *J. Org. Chem.* **1995**, *60*, 7037. (b) Davis, F. A.; Szweczyk, J. M.; Reddy, R. E. *J. Org. Chem.* **1996**, *61*, 2222.
- (11) Kamimura, A.; Okawa, H.; Morisaki, Y.; Ishikawa, S.; Uno, H. *J. Org. Chem.* **2007**, *72*, 3569.
- (12) (a) Kochi, T.; Tang, T. P.; Ellman, J. A. *J. Am. Chem. Soc.* **2002**, *124*, 6518. (b) Davis, F. A.; Gaspari, P. M.; Nolt, B. M.; Xu, P. *J. Org. Chem.* **2008**, *73*, 9619.
- (13) (a) Kawecky, R. *Tetrahedron: Asymmetry* **2006**, *17*, 1420. (b) Andreassen, T.; Haland, T.; Hansen, L. K.; Gautun, O. R. *Tetrahedron Lett.* **2007**, *48*, 8413.
- (14) Viso, A.; Fernández de la Pradilla, R.; Ureña, M.; Colomer, I. *Org. Lett.* **2008**, *10*, 4775.
- (15) (a) Davis, F. A.; Deng, J. H. *Org. Lett.* **2004**, *6*, 2789. (b) Davis, F. A.; Zhang, Y. F.; Qiu, H. *Org. Lett.* **2007**, *9*, 833.