


C. F. CARTER, H. LANGE, D. SAKAI, I. R. BAXENDALE, S. V. LEY\* (UNIVERSITY OF CAMBRIDGE, UK AND MITSUBISHI TANABE PHARMA CORPORATION, YOKOHAMA, JAPAN)  
Diastereoselective Chain-Elongation Reactions Using Microreactors for Applications in Complex Molecule Assembly  
*Chem. Eur. J.* **2011**, *17*, 3398–3405.

# Crotylation and Homopropargylation by Flow Technology



**Significance:** A new flow reaction technology for diastereoselective Roush crotylation and Marshall homopropargylation as well as in-line purification protocols was described. Thus, the flow Roush reaction of aldehydes **1** with boronate **2** gave the corresponding homoallyl alcohols **3** in 70–94% yield with *syn* selectivity (*syn/anti* = 3.6–20:1). Similarly, the diastereoselective flow Marshall homopropargylation of allene **(P)-4** with aldehyde **(S)-5** was carried out with  $\text{BF}_3\cdot\text{OEt}_2$  to give homopropargyl alcohol **6** in 75% yield.

**Comment:** An automated multi-step flow protocol [(1) the reduction of ester **(S)-7** with DIBAL; (2) the purification with IRA-743/SiO<sub>2</sub>; (3) the treatment with boronate **2a** in a microreactor; (4) the purification with IRA-743] afforded **3a** in 78% yield. The authors indicated that flushing the pumps, valves, and reaction coils with isopropyl alcohol, followed by acetone and then the dry solvent of choice under an inert atmosphere for at least two hours was needed for a suitable process (to eliminate all traces of water from the reactor).

**SYNFACTS Contributors:** Yasuhiro Uozumi, Yoichi M. A. Yamada  
*Synfacts* 2011, 6, 0685–0685 Published online: 19.05.2011  
DOI: 10.1055/s-0030-1260385; Reg-No.: Y05011SF