R. MOSSETTI, D. SAGGIORATO, G. C. TRON* (UNIVERSITÁ DEGLI STUDI DEL PIEMONTE ORIENTALE 'A. AVOGADRO', NOVARO, ITALY)

Exploiting the Acylating Nature of the Imide-Ugi Intermediate: A Straightforward Synthesis of Tetrahydro-1,4benzodiazepin-2-ones

J. Org. Chem. 2011, 76, 10258-10262.

Tetrahydro-1,4-benzodiazepin-2-ones from **Ugi MCR Intermediates**

$$\begin{array}{c} \text{NC} \\ + \text{ AcOH } + \\ \text{R1} \end{array} \begin{array}{c} \text{CHO} \\ + \\ \text{NH} \\ \text{R2} \end{array} \begin{array}{c} \text{CH}_2\text{Cl}_2, 4 \text{ Å MS}, 2 \text{ h} \\ \text{or 40 °C, 3 d (R^1 = Ar)} \end{array}$$

Significance: Tron and co-workers set to test the very interesting concept of setting a nucleophilic intramolecular trap for the intermediate Ugi imide, normally generated in a MCR reaction. The Ugi imides A were synthesized in a straightforward fashion and, after purification, subjected to a Staudinger reduction followed by heating, which one derivatives. Only trimethylphosphine was re-

 $R^2 = H$, *n*-Bu, *c*-Pr, Bn, 4-FC₆H₄CH₂, 4-ClC₆H₄CH₂

R³ = H, 5-Br, 4,5-diOMe, 4,5-benzo

 $R^1 = n$ -Hex, Cy, *i*-Pr, Ph, Bn, 4-ClC₆H₄, 4-MeOC₆H₄, 4-FC₆H₄, thiophen-2-yl

resulted in the formation of 1,4-benzodiazepine-2ported to effect the reduction as the corresponding iminophosphoranes generated from other phosphines were too stable to hydrolysis even at reflux. The yields achieved per step were usually good, around 70%, giving rise to final product yields of 14-46%. The substrate scope was reasonably well studied.

Comment: Benzodiazepines are widely appreciated for various biological activities, for example, antitumor and anti-HIV (S. Forso Mini-Rev. Med. Chem. 2010, 7, 68). Surprisingly, it appears that only three synthetic strategies for tetrahydro-1,4benzodiazepin-2-ones were reported prior to the current procedure. It is procedurally simple and utilizes mostly inexpensive starting materials. The only drawback of this procedure is the need for trimethylphosphine, which is somewhat expensive and has pyrophoric tendencies. Sadly, the scale of reactions performed was not evident either in the article or in the supporting information.

SYNFACTS Contributors: Victor Snieckus, Toni Rantanen Synfacts 2012, 8(3), 0259 Published online: 20.02.2012 DOI: 10.1055/s-0031-1290287; Reg-No.: V00512SF

Category

Synthesis of **Heterocycles**

Key words

tetrahvdro-1.4benzodiazepin-2ones

Ugi reaction

trimethylphosphine

