# **SYNLETT Spotlight 426**

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

## **Bis(pinacolato)diboron**

## Compiled by Rémy Hemelaere

Rémy Hemelaere was born in 1987 in Menin, Belgium. He graduated from National Polytechnic School of Chemical Engineering and Technology of Caen (Ensicaen, France) in 2010 and received his M.Sc. degree in organic chemistry in the same year. He is currently pursuing his Ph.D. under the supervision of Dr. François Carreaux at the University of Rennes 1, France. His Ph.D. research project focusses on allylboronate chemistry and the utilisation of ruthenium metathesis catalysts. He is developing methodologies for the total synthesis of natural molecules.

Université de Rennes 1 - Institut des Sciences Chimiques de Rennes - UMR CNRS 6226, 263 Avenue du Général Leclerc, Campus de Beaulieu - Bâtiment 10B, 35042 Rennes, France E-mail: remy.hemelaere@univ-rennes1.fr

## Introduction

Bis(pinacolato)diboron (B2pin2 Figure 1, CAS: 73183-34-3) is an air-stable, odourless, colorless powder that is commercially available. Being a major tool for the introduction of boron atoms into organic compounds, it is widely known as a good reagent to prepare substrates for the Suzuki-Miyaura reaction.

An extensive scope of reactions, including C-H activation of  $C(sp^2)$  –H and  $C(sp^3)$  –H bonds,<sup>1</sup> borylation of  $\alpha,\beta$ -unsaturated derivatives and substitutions of allylic carbonates, have been recently described in the literature. Over



Borylated products, obtained from these new strategies, could successfully be used in oxidation, allylation or coupling reactions.

[Rh(OMe)(cod)]<sub>2</sub>

(0.5 mol%, Rh)

silica-TRIP (0.5 mol%)

Cu(Ot-Bu) (5.0 mol%)

(*R*,*R*)-QuinoxP\* (5.0 mol%)

hexane, 40-60 °C, 1



B<sub>2</sub>pin<sub>2</sub>

Figure 1

 $R^1 = Alk, NR^2_2$ 

+

## Abstracts

#### (A) C(sp<sup>3</sup>)-H Activation

Very recently, Sawamura et al. reported a rhodium-catalyzed  $C(sp^3)$ -H borylation of amides and urea derivatives at the position adjacent to nitrogen with a silica-supported triarylphosphine ligand (silica-TRIP).<sup>4</sup> The reaction was carried out under mild conditions with good to excellent yield. Compounds obtained with this method can undergo Suzuki-Miyaura coupling.

#### (B) Allylic Substitution

Under mild conditions, allylic carbonates can be borylated by treatment with bis(pinacolato)diboron in the presence of a copper catalyst to give the corresponding allyl boronates.<sup>5</sup> Using a chiral ligand, excellent enantioselectivities were obtained.6

## (C) Borylation of Aldehydes

Treatment of aldehydes with B<sub>2</sub>pin<sub>2</sub>, in the presence of a copper catalyst, led to the formation of diboration products as stable compounds in the solid state. A mechanistic study was performed confirming the insertion of the carbonyl group into the copperboron bond. A selective hydrolysis of the B-O bond during chromatographic purification can provide the corresponding a-hydroxyboronates.7

SYNLETT 2013, 24, 0659-0660 Advanced online publication: 12.02.2013 DOI: 10.1055/s-0032-1318171; Art ID: ST-2012-V0433-V © Georg Thieme Verlag Stuttgart · New York





Bpir

Bpin

Мe

70-90% vield

### (D) Borylation of Imines

Ellman et al. reported that the (ICy)CuOt-Bu–B<sub>2</sub>pin<sub>2</sub> system can promote asymmetric borylation of chiral *N-tert*-butanesulfinyl aldimines.<sup>8</sup> More recently, Fernandez and co-workers described the enantioselective synthesis of  $\alpha$ -amino boronate esters via an addition of B<sub>2</sub>pin<sub>2</sub> to *N*-tosyl aldimines without transition-metal complexes.<sup>9</sup>

#### (E) Borylation of Allylic Alcohols

Szabó and co-workers described the borylation of allylic alcohols using a SCS palladium pincer complex catalyst. Depending on the solvent system, a regioselective formation of linear or branched allyl boronates is possible. Using an aldehyde, allylic alcohols were obtained via a one-pot borylation–allylation process.<sup>10</sup>







#### (F) *Borylation of Alkynes* Catalytic systems used for the introduction of boron atoms into al-

kynes have been improved over the past ten years in order to obtain better regioselectivity. Hoveyda et al. developed a highly selective method to synthesize internal vinylboronates from terminal alkynes by using N-heterocyclic carbene (NHC) complexes of copper(I).<sup>11</sup> Excellent regioselectivities were observed with a high level of functional group tolerance.

#### (G) Borylation of Primary Alkyl Halides

Starting from primary alkyl bromides, Biscoe et al. reported the synthesis of alkylboronates, catalyzed by complexes of palladium. Diverse functional groups were tolerated on the starting material, such as nitrile, alcohol, ester and amide. Moderate to good yields were obtained with complete selectivity over secondary bromides.<sup>12</sup>

(H) Addition to  $\alpha,\beta$ -Unsaturated Carbonyl Compounds

Over the past eight years, research has focused on  $\beta$ -borylation of  $\alpha$ , $\beta$ -unsaturated esters, <sup>13</sup> amides, <sup>14</sup> and ketones.<sup>15</sup> These processes are efficiently catalyzed by N-heterocyclic carbene (NHC) complexes of copper(I). Hoveyda's group applied this strategy to trisubstituted alkenes of acyclic  $\alpha$ , $\beta$ -unsaturated esters. With a chiral NHC– copper complex, they developed an enantioselective pathway to boron-substituted quaternary carbons.

## References

- (a) Maleczka, R. E.; Shi, F.; Holmes, D.; Smith, M. R. J. Am. Chem. Soc. 2003, 125, 7792. (b) Murphy, J. M.; Tzschucke, C. C.; Hartwig, J. F. Org. Lett. 2007, 9, 757.
- (2) (a) Pelz, N. F.; Woodward, A. R.; Burks, H. E.; Sieber, J. D.; Morken, J. P. J. Am. Chem. Soc. 2004, 126, 16328. (b) Yuan, W.; Ma, S. Adv. Synth. Catal. 2012, 354, 1867.
- (3) Cho, H. Y.; Yu, Z.; Morken, J. P. Org. Lett. 2011, 13, 5267.
- (4) Kawamorita, S.; Miyazaki, T.; Iwai, T.; Ohmiya, H.;
- Sawamura, M. J. Am. Chem. Soc. 2012, 134, 12924.
  (5) Ito, H.; Kawakami, C.; Sawamura, M. J. Am. Chem. Soc.
- 2005, 127, 16034.
  (6) Ito, H.; Ito, S.; Sasaki, Y.; Matsuura, K.; Sawamura, M. J. Am. Chem. Soc. 2007, 129, 14856.
- (7) Laitar, D. S.; Tsui, E. Y.; Sadighi, J. P. J. Am. Chem. Soc. 2006, 128, 11036.
- (8) Beenen, M. A.; An, C.; Ellman, J. A. J. Am. Chem. Soc. 2008, 130, 6910.

- (9) Solé, C.; Gulyás, H.; Fernandez, E. Chem. Commun. 2012, 48, 3769.
- (10) Selander, N.; Szabó, K. J. Org. Chem. 2009, 74, 5695.
- (11) Jang, H.; Zhugralin, A. R.; Lee, Y.; Hoveyda, A. H. J. Am. Chem. Soc. 2011, 133, 7859.
- Joshi-Pangu, A.; Ma, X.; Diane, M.; Iqbal, D.; Kribs, R. J.; Huang, R.; Wang, C-Y.; Biscoe, M. R. J. Org. Chem. 2012, 77, 6629.
- (13) (a) Lee, J.; Yun, J. Angew. Chem. Int. Ed. 2008, 47, 145.
  (b) O'Brien, J. M.; Lee, K.; Hoveyda, A. H. J. Am. Chem. Soc. 2010, 132, 10630. (c) Lee, J. Ch. H.; McDonald, R.; Hall, D. G. Nat. Chem. 2011, 3, 894.
- (14) Chea, H.; Sim, H.; Yun, J. Adv. Synth. Catal. 2009, 351, 855.
- (15) (a) Chen, I.; Yin, L.; Itano, W.; Kanai, M.; Shibasaki, M. J. Am. Chem. Soc. 2009, 131, 11664. (b) Wu, H.; Radomkit, S.; O'Brien, J. M.; Hoveyda, A. H. J. Am. Chem. Soc. 2012, 134, 8277.

Bpin

90-98% yield

90-98% ee



B<sub>2</sub>pin<sub>2</sub>

R = Alk. Ar

NHC (5.0 mol%)

CuCl (5.0 mol%)

t-BuONa (13.0 mol%)

MeOH (1.2 equiv)

THF, -78 °C, 18 h

EtO