

An unexpected finding on gastroscopy: gastro-gastric fistula with *Helicobacter pylori* and *Giardia lamblia*




Fig. 1 View of the incisura during the index gastroscopy showing a deep cratered ulcer in a 66-year-old woman with a 6-week history of vomiting, dysphagia, black stool, with weight loss, and melena on per rectum examination. The angle of view in **Fig. 2** is indicated by the arrow.




Fig. 2 J maneuver in the antrum and withdrawal, allowing a closeup view of the ulcer base, showing two fistulae in the gastric body. The proximal stomach and the proximal gastroscope are clearly seen.




Fig. 3 Histologic section from an ulcer biopsy sample.

A 66-year-old woman presented with a 6-week history of vomiting, dysphagia, black stool, and weight loss, with melena on per rectum examination. Gastroscopy revealed a large chronic ulcer at the incisura with two gastro-gastric fistulae between the antrum and the body (**Fig. 1**). **Fig. 2** shows the proximal endoscope markings visible through a fistula. Biopsies and computed tomography (CT) of the abdomen did not show any evidence of neoplasia, but histologic examination revealed numerous *Giardia lamblia* parasites (**Fig. 3**). *Helicobacter pylori* was not seen but the rapid urease test was positive. The patient was treated with intravenous

pantoprazole, oral metronidazole, and eradication therapy, and no more bleeding was observed. *H. pylori* was successfully eradicated and following discharge the patient gained weight with no further signs of gastrointestinal bleeding. Repeat endoscopy showed healing of the ulcer but persistent incisura deformity and gastro-gastric fistulation.

Although peptic ulcer disease, Crohn's, and cancer have been postulated to cause fistulation, we could find no published data to confirm this. Reports of gastro-gastric fistulae in the literature are almost exclusively related to obesity surgery. This complication occurs in up to 1.2% of Roux-

en-Y procedures, [1] which seems to be the most common cause of gastro-gastric fistulae. The presence of *Giardia* seems incidental although it has previously been linked to *H. pylori* infection [2,3]. In a large retrospective case series of patients with giardiasis, 8.7% had gastric colonization but this was not associated with any specific gastric histology [4]. In addition, since *Giardia* is not usually associated with ulceration in its more usual habitat of the small bowel [4], it can be assumed to be an unrelated finding. We believe our patient most likely had chronic peptic ulcer disease related to *H. pylori* with, perhaps, perforation and subsequent fistula formation.

Endoscopy_UCTN_Code_CCL_1AB_2AD_3AC

Competing interests: None

J. Weeks, P. Mooney, G. Lipscomb, J. M. Pearson, A. Ong, S. Singh

Department of Gastroenterology,
Royal Bolton Hospital NHS Foundation
Trust, Bolton, United Kingdom

References

- 1 Carrodeguas L, Szomstein S, Soto F et al. Management of gastrogastric fistulas after divided Roux-en-Y gastric bypass surgery for morbid obesity: analysis of 1,292 consecutive patients and review of literature. *Surg Obes Relat Dis* 2005; 1: 467–474
- 2 Misra V, Misra SP, Dwivedi M et al. *Giardia lamblia* trophozoites in gastric biopsies. *Indian J Pathol Microbiol* 2006; 49: 519–523
- 3 Molina Infante J, Ferrando Lamana L. Novel endoscopic findings in gastric giardiasis. *Endoscopy* 2009; 41: E275
- 4 Oberhuber G, Kastner N, Stolte M. Giardiasis: a histologic analysis of 567 cases. *Scand J Gastroenterol* 1997; 32: 48–51

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1326259>
Endoscopy 2013; 45: E118
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Dr S. Singh

Department of Gastroenterology
Royal Bolton Hospital NHS Foundation Trust
Minerva Road, Farnworth
Bolton, BL4 0JR
United Kingdom
Fax: +44 1204 390 141
Salil.Singh@rbh.nhs.uk