

Nasopharyngeal carcinoma with esophageal metastasis presenting with progressive dysphagia



Fig. 1 Contrast-enhanced thoracic computed tomography (CT) scan showing metastatic spread to axillary, prevascular, paratracheal, precarinal, subcarinal, paraesophageal, perigastric, para-aortic, and intercaro-aortic lymph nodes.




Fig. 2 Esophagoscopy showing lumen narrowing and a protruding mass between 30 and 35 cm from the upper incisors.




Fig. 3 Histology of the esophageal biopsy specimen. **a** In situ hybridization for Epstein-Barr virus (EBV)-encoded RNA-1 (EBER-1) with DAB (3,3'-diaminobenzidine) chromogen showing brown positive nuclear staining (original magnification $\times 200$). **b** Immunostaining for high molecular weight cytokeratin with AEC (3-amino-9-ethylcarbazole) chromogen showing red positive cytoplasmic staining (original magnification $\times 200$). The arrow indicates cells that are showing dual staining.

A 49-year-old man attended our department complaining of progressive difficulty in swallowing over a period of 2 weeks. He reported no other discomfort. His medical history revealed that he had been diagnosed with nasopharyngeal carcinoma (cT3N3bM0; WHO Type III) in September 2008 and had undergone chemotherapy and radiotherapy concurrently. A complete response was achieved initially, but regional recurrence in the right side of his neck required him to undergo neck dissection in June 2010. Progressive dysphagia brought him to our department 4 months later. Imaging studies of the head and neck region produced negative findings, but an esophagogram revealed

narrowing in the middle third of his esophagus.

Contrast-enhanced thoracic computed tomography (CT) demonstrated multiple nodal metastases in the thoracic cavity and enlargement of the retrocardiac segment of the esophagus (Fig. 1). Esophagoscopy demonstrated a protruding mass located 30–35 cm from the upper incisors (Fig. 2). Biopsy showed a poorly differentiated carcinoma; metastasis from the nasopharyngeal carcinoma was suspected. In addition, in situ hybridization showed that the neoplastic epithelial cells were positive for both Epstein-Barr virus (EBV)-encoded RNA-1 (EBER-1; Fig. 3a) and high molecular weight cytokeratin

(Fig. 3b). These findings were consistent with metastatic nasopharyngeal carcinoma, as in situ hybridization results from a previous nasopharyngeal specimen were also positive for EBER-1 in neoplastic epithelial cells. The patient underwent palliative chemotherapy but died of his disease 5 months later.

Distant spread is more common with nasopharyngeal carcinoma than with other head and neck squamous cell carcinomas. The most frequent sites of distant metastases are bone, lung, liver, and distant lymph nodes [1,2]. A clinical diagnosis of metastatic cancer in the esophagus arising from nasopharyngeal carcinoma is extremely rare. To the best of our knowledge, this type of metastasis has not been previously described in the English literature.

In conclusion, secondary esophageal carcinoma from nasopharyngeal carcinoma is rare, presents as a distant recurrence, and has a very poor prognosis.

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AB

Competing interests: None

S.-Y. Yang, C.-Y. Chien

Department of Otolaryngology, Kaohsiung Chang Gung Memorial Hospital and Chang Gung University College of Medicine, Kaohsiung, Taiwan

References

- 1 Altun M, Fandi A, Dupuis O et al. Undifferentiated nasopharyngeal cancer (UCNT): current diagnostic and therapeutic aspects. *Int J Radiat Oncol Biol Phys* 1995; 32: 859–877
- 2 Hsu MM, Tu SM. Nasopharyngeal carcinoma in Taiwan. Clinical manifestations and results of therapy. *Cancer* 1983; 52: 362–368

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1326274>
Endoscopy 2013; 45: E120–E121
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

C.-Y. Chien, MD
Department of Otolaryngology
Kaohsiung Chang Gung Memorial Hospital
123 Ta-Pei Road
Niao-Song District
Kaohsiung 833
Taiwan
Fax: +886-7-7313855
cychien3965@adm.cgmh.org.tw