

Small-intestinal hemorrhage caused by treatment with sorafenib for hepatocellular carcinoma and diagnosed by capsule endoscopy

Sorafenib, a multikinase inhibitor, is the first-line treatment for unresectable progressive hepatocellular carcinoma (HCC) and has been approved by both the United States Food and Drug Administration (FDA) and the European Medicines Agency (EMA) [1,2]. Although there are studies reporting its effectiveness, there are also reports about adverse events related to treatment with this drug. Among the noted side effects is gastrointestinal hemorrhage, although only a few studies have reported the frequency of this complication [3,4]. This is the first case report of small-intestinal hemorrhage following sorafenib treatment, and includes a description of the finding of hemorrhagic lesions as determined by capsule endoscopy. An 80-year-old woman with hepatitis C liver cirrhosis and being treated at a local hospital underwent ultrasound, which showed a mass lesion in the S8 region (40 mm in diameter). She was referred to our hospital for further investigation and treatment of the mass lesion. Based on the findings of dynamic computed tomography (CT) and ethoxybenzyl magnetic resonance imaging (EOB-MRI), the patient was diagnosed as having HCC. We recommended surgical treatment and transcatheter arterial chemoembolization (TACE) but the patient declined this treatment option. Therefore, sorafenib 400 mg/day was started on day 20 in the hospital. On day 28, tarry stools were observed, and serum chemistry investigations revealed red blood count $369 \times 10^4/\mu\text{L}$, hemoglobin 11.8 g/dL, and hematocrit 34.1%, which suggested normocytic hypochromic anemia. On day 30, esophagogastroduodenoscopy revealed no hemorrhagic esophageal varices with LiF1CbRC(-) (Fig. 1). On day 35, colonoscopy revealed dark blood throughout the colon but no evident hemorrhagic lesions (Fig. 2). On day 36, the small intestine was examined with a capsule endoscope. Oozing of blood from several hemorrhagic lesions was observed. This was diagnosed as small-



Fig. 1 Esophagogastroduodenoscopy in an 80-year-old woman with hepatocellular carcinoma and tarry stools showing no hemorrhagic varices (LiF1CbRC(-)) in the esophagus.



Fig. 2 Colonoscopy showed no active hemorrhagic lesions.

intestinal hemorrhage (Fig. 3) as a side effect of sorafenib therapy. The symptoms quickly improved on cessation of the treatment. Capsule endoscopy on day 90 showed scarlike lesions in the distal small intestine, indicating significant improvement (Fig. 4).

To our knowledge, this is the first report of hemorrhage in the small intestine caused by sorafenib treatment and diagnosed by capsule endoscopy.

Endoscopy_UCTN_Code_CCL_1AB_2AZ_3AD

Competing interests: None




Fig. 3 First report of hemorrhagic lesions in the small intestine detected by capsule endoscopy (arrows). a Proximal small intestine. b Distal small intestine.



Fig. 4 Second capsule endoscopy of the small intestine showing improvement (arrows).

Y. Takahashi^{1,2}, S. Fukunishi², T. Nishikawa^{1,2}, S. Nouda², Y. Sasaki¹, M. Sanomura¹, E. Umegaki², K. Higuchi²

¹ Department of Gastroenterology, Hokusetsu General Hospital, Osaka, Japan

² 2nd Department of Internal Medicine, Osaka Medical College, Osaka, Japan

References

1 *Wilhelm SM, Carter C, Tang I et al.* BAY 43-9006 exhibits broad spectrum oral anti-tumor activity and targets the RAM/MEK/ERK pathway and receptor tyrosine kinase

involved in tumor progression and angiogenesis. *Cancer Res* 2004; 64: 7099–7109

2 *Chang YS, Adnane JZ, Trail PA et al.* Sorafenib (BAY 43-9006) inhibits tumor growth and vascularization and induces tumor apoptosis and hypoxia in RCC xenograft models. *Cancer Chemother Pharmacol* 2007; 59: 561–574

3 *Llovet JM, Ricci S, Mazzeferro V et al.* Sorafenib in advanced hepatocellular carcinoma. *N Engl J Med* 2008; 359: 378–390

4 *Cheng AL, Kang YK, Chen Z et al.* Efficacy and safety of sorafenib in patients in the Asia-Pacific region with advanced hepatocellular carcinoma: a phase randomized, double-blind, placebo-controlled trial. *Lancet Oncol* 2009; 10: 25–34

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1326641>
Endoscopy 2013; 45: E179–E180
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

S. Fukunishi

2-7 Daigaku-machi
Takatsuki
Osaka 569-8686
Japan
Fax: +81-726-846532
in2104@poh.osaka-med.ac.jp