

SYNLETT Spotlight 455

5-Alkyl and 5,5-Dialkyl Meldrum's Acids

Compiled by Inese Mierina



This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Inese Mierina was born in Riga, Latvia, in 1985. She received her B.Sc. (2007) and Mg.Sc. (2009) in chemical engineering from Riga Technical University (RTU). She is currently working toward her Ph.D. under supervision of Professor Mara Jure at RTU. Her research is focused on natural antioxidant analogues and plant material extracts as potential antioxidants for vegetable oils.


Riga Technical University, Faculty of Material Science and Applied Chemistry, Azenes Str. 14/24, 1048 Riga, Latvia
E-mail: inesem@ktf.rtu.lv

Introduction

5-Alkyl and 5,5-dialkyl Meldrum's acids (**1** and **2**, respectively) are used in total synthesis.¹ Monoalkylated derivatives **1** are synthesized from Meldrum's acid **3** by condensation with aldehyde² (or by acylation³) followed by hydrogenation.

5,5-Dialkyl Meldrum's acids **2** can be obtained by alkylation of Meldrum's acid **3** or its monoalkylated derivatives **1** (Scheme 1).⁴ Herein, reactions of Meldrum's acids **1** and


2 proceeding with destruction of 1,3-dioxane cycle are reviewed.




Scheme 1

Abstracts


(A) Hydrolysis of Meldrum's acid **2** leads to malonic or acetic acid derivatives.⁵ Radical reduction of compound **2** with SmI_2 and H_2O forms 3-hydroxypropanoic acids **4** selectively. The first step is activation of Meldrum's acid via coordination of SmI_2 to the carbonyl group, followed by electron transfer.⁶



(B) Detz et al.⁷ have reported that dimethyl malonate **5** can be obtained from propargylic derivative **2**. The authors propose that the first step of the reaction cascade is copper-mediated addition of Meldrum's acid **2** to the triple bond and sequential methanolysis of the dioxane cycle forming lactone **6**, which is further cleaved with methoxide leading to compound **5**.




(C) β -Substituted aldehydes **7** can be synthesized by Lewis base promoted hydrosilylation of Meldrum's acids **1** with phenylsilane, followed by hydrolysis. In situ treatment of aldehyde **7** with an amine and sequential hydrogenation of the formed imine with H_2 in the presence of Pd/C or with $\text{NaBH}(\text{OAc})_3$ gives γ -substituted amines **8**.⁸



i) 1) Et_3N (2 equiv), PhSiH_3 (3 equiv), THF, r.t.; 2) H_2O
ii) 1) Et_3N (2 equiv), PhSiH_3 (3 equiv), THF, r.t.; 2) $\text{R}^3\text{R}^4\text{NH}$ (2 equiv), MeOH; 3) H_2 , Pd/C

(D) 5-Alkyl Meldrum's acids **1** are used for the rapid synthesis of 2-alkyl acrylates **9** via Mannich-type reactions. The advantage of the method is the clean conversion into products due to the formation of volatile by-products – acetone, carbon dioxide, and dimethylamine.⁹




SYNLETT 2014, 25, 0155–0156

Advanced online publication: 21.11.2013


DOI: 10.1055/s-0033-1340156; Art ID: ST-2013-V0462-V

© Georg Thieme Verlag Stuttgart · New York


(E) Derivatives of 5-(but-3-enyl) Meldrum's acid **10** are suitable for the synthesis of cyclopentanols **11** via radical cyclization upon treatment with SmI_2 in H_2O .⁶ *Exo-trig/exo-trig* radical cyclization cascade occurs, when substituent R^2 is an alkene or alkyne; such a transformation gives fused bicyclic system **12**.¹⁰




(F) Meldrum's acid can act as a carbon-based leaving group. Catalytic hydrogenolysis of Meldrum's acids **2** ($\text{R}^1 = \text{H}$) is an excellent route for the synthesis of compounds **13** both with secondary and tertiary benzylic stereocenters in 65–96% yield under mild reaction conditions. As the reaction proceeds with inversion at the stereocenter, an $\text{S}_{\text{N}}2$ mechanism is proposed.¹¹ Treatment of Meldrum's acid derivatives **2** ($\text{R}^1 = \text{H}, \text{Me}$) with nucleophiles in the presence of Lewis acids furnishes compounds **14** and **15**. The yields vary from 51% to quantitative.¹²




(G) Johnson and co-workers reported the hydroperoxidation of 5-alkyl Meldrum's acids **1** with O_2 in the presence of $\text{Cu}(\text{NO}_3)_2$.¹³ These reaction conditions are compatible with unsaturated bonds in substituent R^1 . The peroxides **16** are suitable for intramolecular oxidation of unsaturated bonds via electrophilic activation; such an approach was used for the synthesis of lactones **17** and **18**.




(H) Addition of 5-substituted Meldrum's acid **1** to prop-2-ynal **19** forms 3-(1,3-dioxan-5-yl)-4,4-dimethoxy-but-2-enal. The thermolysis of 2,3-unsaturated aldehyde intermediate provides a synthetic procedure to 2*H*-pyran-2-one **20**. Hydrolysis of acetal moiety results in 4-formyl pyran-2-one **21**.¹⁴



(I) Copper- and iron-¹⁵ or silver-catalyzed¹⁶ tandem cyclization–hydrolysis–decarboxylation of 5-propargyl Meldrum's acid **2** is an efficient approach for the synthesis of *Z*- γ -alkylidene lactones **22**. The compatibility of copper(I) and iron(III) is not established yet. The authors¹⁵ suggest that copper(I) activates the alkyne moiety, but iron(III) interacts with the oxygen atom.



(J) Intermolecular cleavage of Meldrum's acid derivatives with nucleophiles is well known. Sapi et al.¹⁷ reported the tandem deprotection–intramolecular cyclization of Meldrum's acids **23** applied for the synthesis of lactones **24a** and lactame **24b**.



References

- Ivanov, A. S. *Chem. Soc. Rev.* **2008**, *37*, 789.
- Tóth, G.; Kövér, K. E. *Synth. Commun.* **1995**, 3067.
- Nutaitis, C. F.; Schultz, R. A.; Obaza, J.; Smith, F. X. *J. Org. Chem.* **1980**, *45*, 4606.
- Chan, C.-C.; Huang, X. *Synthesis* **1982**, 452.
- (a) Scheuer, P. J.; Cohen, S. G. *J. Am. Chem. Soc.* **1958**, *80*, 4933. (b) Jacobs, R. T.; Wright, A. D.; Smith, F. X. *J. Org. Chem.* **1982**, *47*, 3769.
- Collins, K. D.; Oliveira, J. M.; Guazzelli, G.; Sautier, B.; De Grazia, S.; Matsubara, H.; Helliwell, M.; Procter, D. J. *Chem.–Eur. J.* **2010**, *16*, 10240.
- Detz, R. J.; Abiri, Z.; le Griel, R.; Hiemstra, H.; van Maarseveen, J. H. *Chem.–Eur. J.* **2011**, *17*, 5921.
- Frost, C. G.; Hartley, B. C. *J. Org. Chem.* **2009**, *74*, 3599.
- Frost, C. G.; Penrose, S. D.; Gleave, R. *Synthesis* **2009**, 627.
- Sautier, B.; Lyons, S. E.; Webb, M. R.; Procter, D. J. *Org. Lett.* **2012**, *14*, 146.
- Wilsily, A.; Nguyen, Y.; Fillion, E. *J. Am. Chem. Soc.* **2009**, *131*, 15606.
- Mahoney, S. J.; Lou, T.; Bondarenko, G.; Fillion, E. *Org. Lett.* **2012**, *14*, 3474.
- Krabbe, S. W.; Do, D. T.; Johnson, J. S. *Org. Lett.* **2012**, *14*, 5932.
- Akué-Gédu, R.; Hénichart, J.-P.; Couturier, D.; Rigo, B. *Tetrahedron Lett.* **2004**, *45*, 9197.
- Li, S.; Jia, W.; Jiao, N. *Adv. Synth. Catal.* **2009**, 351, 569.
- Jia, W.; Li, S.; Yu, M.; Chen, W.; Jiao, N. *Tetrahedron Lett.* **2009**, *50*, 5406.
- (a) Dardennes, E.; Kovács-Kulyassa, Á.; Renzetti, A.; Sapi, J.; Laronze, J.-Y. *Tetrahedron Lett.* **2003**, *44*, 221. (b) Dardennes, E.; Kovács-Kulyassa, Á.; Boisbrun, M.; Petermann, C.; Laronze, J.-Y.; Sapi, J. *Tetrahedron: Asymmetry* **2005**, *16*, 1329. (c) Dardennes, E.; Gérard, S.; Petermann, C.; Sapi, J. *Tetrahedron: Asymmetry* **2010**, *21*, 208.