



S. REGATI, Y. HE, M. THIMMAIAH, P. LI, S. XIANG, B. CHEN,\* J. C.-G. ZHAO\*  
(UNIVERSITY OF TEXAS AT SAN ANTONIO, USA AND FUJIAN NORMAL UNIVERSITY,  
FUZHOU, P. R. OF CHINA)  
Enantioselective Ring-Opening of *meso*-Epoxides by Aromatic Amines Catalyzed by a Homochiral Metal-Organic Framework  
*Chem. Commun.* **2013**, *49*, 9836–9838.

# Asymmetric Ring-Opening of *meso*-Epoxides with a Chiral Metal-Organic Framework



## Typical results:



**Significance:** A chiral MOF (**UTSA-32**) was prepared from (*S*)-4,4'-dibromo-6,6'-dichloro-2,2'-diethoxy-1,1'-binaphthyl in three steps (eq. 1). **UTSA-32** catalyzed the enantioselective ring opening of *meso*-epoxides with anilines to give the corresponding  $\alpha$ -hydroxyamines in 69–95% yield with 12–89% ee (11 examples, eq. 2).

**Comment:** **UTSA-32** was characterized by single-crystal X-ray diffraction, PXRD, TGA, and BET analyses. Single-crystal X-ray diffraction analysis of **UTSA-32** revealed that the binuclear zinc clusters are bridged by the carboxylic groups of organic linker **L** to form a three-dimensional framework.

**SYNFACTS Contributors:** Yasuhiro Uozumi, Takao Osako  
*Synfacts* 2013, 9(12), 1355 Published online: 18.11.2013  
DOI: 10.1055/s-0033-1340237; Reg-No.: Y11913SF