



H. MIYAMURA, G. C. Y. CHOO, T. YASUKAWA, W.-J. YOO, S. KOBAYASHI* (THE UNIVERSITY OF TOKYO, JAPAN)

A Heterogeneous Layered Bifunctional Catalyst for the Integration of Aerobic Oxidation and Asymmetric C–C Bond Formation

Chem. Commun. **2013**, *49*, 9917–9919.

A Heterogeneous Bifunctional Au/Pd Nanoparticle Organocatalyst



Significance: The preparation of a polymer-incarcerated Au/Pd-coated organocatalyst [PI(Au/Pd)-CO] for the sequential aerobic oxidation–Michael reaction between γ -substituted allylic alcohols and dibenzylmalonate was described. The reaction of 3-phenyl-2-propen-1-ol (**1**) and dibenzylmalonate (**2**) proceeded in the presence of PI(Au/Pd)-CO to give dibenzyl 2-(3-oxo-1-phenylpropyl)propane-1,3-dioate (**3**) in 75% yield with 90% ee (other 9 examples: 34–83% yield, 74–91% ee).

Comment: The Au/Pd organocatalyst was prepared by the copolymerization of monomers **5**, **6** and **7** in the presence of V-601 [dimethyl 2,2'-azo-bis(2-methylpropionate)] as a radical initiator followed by the formation of the cross-linked shell layer of a polymer **9** in which bimetallic Au/Pd nanoparticles were incorporated. Thus, PI(Au/Pd)-CO was composed of an organocatalytic resin core and a polymeric Au/Pd nanoparticle shell.

SYNFACTS Contributors: Yasuhiro Uozumi, Yoichi M. A. Yamada, Takuma Sato

Synfacts 2014, 10(1), 0102 Published online: 13.12.2013

DOI: 10.1055/s-0033-1340401; Reg-No.: Y14213SF

2014 © THIEME STUTTGART • NEW YORK