




W. TANG, S. JOHNSTON, C. LI, J. A. IGGO, J. BACSA, J. XIAO* (UNIVERSITY OF LIVERPOOL, UK)


Cooperative Catalysis: Combining an Achiral Metal Catalyst with a Chiral Brønsted Acid Enables Highly Enantioselective Hydrogenation of Imines

Chem. Eur. J. **2013**, *19*, 14187–14193.

Enantioselective Hydrogenation of Imines Using Cooperative Catalysis



Selected examples:



Significance: Optically active amines are common in many fine chemicals, agrochemicals, and pharmaceuticals. The authors report a cooperative metal-organocatalytic system utilizing a chiral Brønsted acid and an achiral iridium catalyst (see below for a Review on transfer hydrogenation).

Review: C. Zheng, S.-L. You *Chem. Soc. Rev.* **2012**, *41*, 2498–2518.

Comment: The authors have reported the cooperative use of a chiral iridium catalyst with a chiral phosphoric acid in the asymmetric hydrogenation of acyclic imines with H₂ (*J. Am. Chem. Soc.* **2008**, *130*, 14450). Here, they report an achiral iridium catalyst with a chiral phosphoric acid in a similar reaction. Alkyl imines, which are known to be difficult substrates for asymmetric hydrogenation, were shown to be excellent substrates in this system, giving enantioselectivities up to 97%.

SYNFACTS Contributors: Mark Lautens, Zafar Qureshi
Synfacts 2014, 10(1), 0067 Published online: 13.12.2013
DOI: 10.1055/s-0033-1340443; Reg-No.: L15313SF