


## Endoscopic ultrasound-guided retrograde pancreatic stent placement for the treatment of stenotic jejunopancreatic anastomosis after a Whipple procedure



**Fig. 1** a–e Images of the pancreatitis and the initial procedure: **a** computed tomography (CT) showing a dilated main pancreatic duct (MPD), fluid collection at the anastomosis (arrowhead), and a thickened jejunal wall (arrows); **b** scope placed at the anastomosis; **c** color Doppler image showing the needle striking into the duct (arrow); **d** plaques visible in the MPD (arrows); **e** a pancreatic stent placed through the jejunum.


Endoscopic ultrasound (EUS)-guided [1, 2] or percutaneous [3] rendezvous methods have been used for the treatment of stenosis at the jejunopancreatic anastomosis following pancreatoduodenectomy. However, the use of EUS-guided retrograde pancreatic duct stenting has not been reported, even though it may be pre-

ferable from the point of view of complications.

A 50-year-old man, who had been suffering from repetitive pancreatitis for 1 year due to stenosis at the jejunopancreatic anastomosis after pancreatoduodenectomy (Fig. 1a), underwent EUS-guided retrograde pancreatic duct stenting. Prior to the procedure, despite careful searching with a forward-viewing scope, the orifice of the pancreatic duct could not be detected due to severe luminal inflammation. A convex-type EUS scope (GF-UCT240; Olympus, Tokyo, Japan) was advanced to the anastomotic site (Fig. 1b) and, using color Doppler, the puncture line was adjusted to avoid blood vessels. A 19-gauge needle (SonoTip Pro Control; Medi-globe, Achenmühle, Germany) was inserted into the main pancreatic duct (Fig. 1c), and advancement of a guide

wire (VisiGlide, 0.025-inch; Olympus) fully into the duct was confirmed using contrast medium (Fig. 1d). Dilation was unsuccessful with a bougie catheter (Soehendra, 4–7 Fr; Cook Medical, Winston-Salem, North Carolina, USA), but was easily completed using a diathermic sheath (Cysto-Gastro Set, 6 Fr; Endo-Flex, Voerde, Germany) [4]. A plastic stent (Geenen, 5 Fr; Cook Medical) was placed (Fig. 1e) and the patient's symptoms disappeared immediately.

Two months later, as scheduled, the stent was upsized to a 7 Fr with balloon dilation (Quantum TTC, 6 mm; Cook Medical) (Fig. 2a). No complications occurred during these procedures and the patient was subsequently healthy (Fig. 2b). Compared with the rendezvous method, the retrograde procedure presented here is a one-step, one-scope method that is



**Fig. 2** a, b The second procedure and following CT: **a** balloon dilation of the stenotic jejunopancreatic anastomosis for upsizing of the stent; **b** CT view showing healing of the jejunal and peripancreatic inflammation and the pancreatic stent in place in the nondilated MPD.

not performed through the abdominal cavity; hence, it carries a lower risk of pancreatic juice leakage and other complications [5]. This method is worthwhile when attempting to rescue a stenotic pancreatojejunostomy after a Whipple resection.

Endoscopy\_UCTN\_Code\_TTT\_1AS\_2AD

**Competing interests:** None

**Hiroyuki Matsubayashi<sup>1</sup>, Yoshihiro Kishida<sup>1</sup>, Kunihiro Shinjo<sup>1</sup>, Kenichiro Imai<sup>1</sup>, Kinichi Hotta<sup>1</sup>, Masaki Tanaka<sup>1</sup>, Naomi Kakushima<sup>1</sup>, Kohei Takizawa<sup>1</sup>, Takashi Mizuno<sup>2</sup>, Yukiyasu Okamura<sup>2</sup>, Hiroyuki Ono<sup>1</sup>**

<sup>1</sup> Division of Endoscopy, Shizuoka Cancer Center, Suntogun, Shizuoka, Japan

<sup>2</sup> Department of Hepatopancreaticobiliary Surgery, Shizuoka Cancer Center, Suntogun, Shizuoka, Japan

## References

- 1 Kikuyama M, Itoi T, Ota Y et al. Therapeutic endoscopy for stenotic pancreatodigestive tract anastomosis after pancreatoduodenectomy (with videos). *Gastrointest Endosc* 2011; 73: 376–382
- 2 Mallery S, Matlock J, Freeman ML. EUS-guided rendezvous drainage of obstructed biliary and pancreatic ducts: report of 6 cases. *Gastrointest Endosc* 2004; 59: 100–107
- 3 Ota Y, Kikuyama M, Suzuki S et al. Percutaneous pancreatic-duct puncture with rendezvous technique can treat stenotic pancreatojejunostomy. *Digest Endosc* 2010; 22: 228–231
- 4 Kawakami H, Kuwatani M, Eto K et al. Resolution of a refractory severe biliary stricture using a diathermic sheath. *Endoscopy* 2012; 44 (Suppl. 02): E119–120
- 5 Will U, Fuedner F, Thieme AK et al. Transgastric pancreatography and EUS-guided drainage of the pancreatic duct. *J Hepatobiliary Pancreat Surg* 2007; 14: 377–382

## Bibliography

**DOI** <http://dx.doi.org/10.1055/s-0033-1358923>  
*Endoscopy* 2013; 45: E435–E436  
 © Georg Thieme Verlag KG  
 Stuttgart · New York  
 ISSN 0013-726X

## Corresponding author

**Hiroyuki Matsubayashi, MD, PhD**

Division of Endoscopy  
 1007 Shimonagakubo  
 Nagaizumi, Suntogun  
 Shizuoka  
 411-8777  
 Japan  
 Fax: +81-55-9895692  
 h.matsubayashi@scchr.jp