

A fish bone perforation of the esophagus

A 73-year-old man presented with odynophagia and retrosternal pain of 3 days' duration following a fish meal. Physical examination disclosed normal vital signs and a temperature of 37.2°C. Laboratory studies showed elevated white blood cells of 13 900/ μ L and C-reactive protein of 14.8 mg/dL. A computed tomography (CT) of the chest revealed a suspicious fish bone that measured 3 cm in length and had perforated through the esophageal wall (● Fig. 1 and ● Fig. 2). Three-dimensional CT showed the bone (blue matter) penetrating close to the left common carotid artery (● Fig. 3). Subsequent upper endoscopy revealed only a small submucosal nodule, which was located at 19 cm from the incisors, not an impacted fish bone in the upper esophagus (● Fig. 4). A tiny white linear scar (arrow) was observed on its top, suggesting the site of perforation (● Fig. 5). Surgical exploration was performed via a lateral neck incision, and the fish bone was successfully retrieved. The postoperative course was uneventful.

Most ingested foreign bodies can pass through the gastrointestinal tract spontaneously. However, 10%–20% of such bodies require nonoperative intervention and 1% need surgery [1]. Based on a large-scale retrospective study including 316 cases of foreign bodies in the esophagus [2], the most common foreign bodies in the pharynx and the upper esophagus were fish bones. The risk of complications was increased with a longer duration of impaction (>24 hours), bone type, and longer bone length (>3 cm). The current case had all of these risk factors. As for endoscopic features of fish bones, most visible bodies can be retrieved by biopsy forceps [3]. Extremely rare cases with imbedded or perforating fish bones may present submucosal tumor-like nodules [4, 5], as in this case.

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AH

Competing interests: None




Fig. 1 An axial computed tomography of the chest disclosed a suspicious fish bone that measured 3 cm in length (black arrow) and had perforated the esophagus (white arrow).




Fig. 2 A coronal computed tomography of the chest showed the bone (black arrow) and the esophagus (white arrow).



Fig. 3 Three-dimensional computed tomography showed the bone (blue matter) penetrating close to the left common carotid artery.




Fig. 4 Upper endoscopy showed only a small submucosal nodule in the upper esophagus not an impacted fish bone.




Fig. 5 A tiny white linear scar (arrow) was observed on the nodule, suggesting the site of perforation.

Akira Hokama¹, Kayoko Uechi¹, Eriko Takeshima², Chiharu Kobashigawa¹, Atsushi Iraha¹, Tetsu Kinjo², Kazuto Kishimoto¹, Fukunori Kinjo², Jiro Fujita¹

¹ Department of Infectious, Respiratory, and Digestive Medicine, University of the Ryukyus, Okinawa, Japan

² Department of Endoscopy, University of the Ryukyus, Okinawa, Japan

References

- 1 Eisen GM, Baron TH, Dominitz JA et al. Guideline for the management of ingested foreign bodies. *Gastrointest Endosc* 2002; 55: 802–806
- 2 Sung SH, Jeon SW, Son HS et al. Factors predictive of risk for complications in patients with oesophageal foreign bodies. *Dig Liver Dis* 2011; 43: 632–635
- 3 Ihama Y, Hokama A, Iraha A et al. Esophageal perforation by fish bone ingestion. *Gastrointest Endosc* 2011; 74: 921
- 4 Chu YC, Chiu HH. A completely imbedded fish bone presenting as an esophageal tumor-like lesion: an unusual presentation. *Gastrointest Endosc* 2008; 68: 1190–1191; discussion 1191
- 5 Tomimori K, Nakasone H, Hokama A et al. Liver abscess. *Gastrointest Endosc* 2004; 59: 397–398

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1364952>
Endoscopy 2014; 46: E216–E217
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Akira Hokama, MD

Department of Infectious, Respiratory,
 and Digestive Medicine
 University of the Ryukyus
 207 Uehara, Nishihara
 Okinawa 903-0215
 Japan
 Fax: +81-98-8951414
hokama-a@med.u-ryukyu.ac.jp