



## To twist or not to twist: a case of ERCP in situs inversus totalis



**Fig. 1** Computed tomography (CT) scan confirming situs inversus with the liver seen on the patient's left side and the spleen on the right.



**Fig. 2** Fluoroscopy image during endoscopic retrograde cholangiopancreatography (ERCP) without use of the mirror-image technique showing the position of the endoscope.



**Fig. 3** Endoscopic views during endoscopic retrograde cholangiopancreatography (ERCP) showing: **a** performance of a normal sphincterotomy; **b** balloon sweeps during the procedure; **c** pigment-type stones being removed during the ERCP.

A 57-year-old woman with history of hypercholesterolemia and situs inversus totalis presented with a chief complaint of epigastric pain and poor appetite for 2 days. The epigastric pain was 7/10 in intensity with no radiation. On physical examination, she had no abdominal scars but there was evidence of hepatosplenomegaly and epigastric tenderness on palpation, although Murphy's sign was negative; bowel sounds were normal on auscultation. The results of laboratory testing revealed normal aspartate transaminase (AST) and alanine transaminase (ALT) levels, but an elevated total bilirubin of 1.3 mg/dL. The alkaline phosphatase (ALP) level was 112 IU/L (normal 45–115 IU/L) and the  $\gamma$ -glutamyltransferase (GGT) was 195 IU/L (normal 0–42 IU/L). Biliary ultrasound revealed a moderately dilated common bile duct and multiple gall stones. A computed tomography (CT) scan of the abdomen and pelvis confirmed the diagnosis of situs inversus totalis with hepatosplenomegaly (► **Fig. 1**).

The patient underwent endoscopic retrograde cholangiopancreatography (ERCP) for her proven choledocholithiasis. Because of the patient having situs inversus totalis, she was placed in a prone position with the endoscopist on the right side of the table (► **Fig. 2**). During the ERCP, the endoscope was rotated through 180° in the second portion of duodenum to allow for the anatomical anomaly. The ampulla was identified with difficulty; however, wire-guided cannulation was then successfully performed. The first cholangiogram demonstrated filling defects and a sphincterotomy was performed (► **Fig. 3**). After this, four pigment-type stones were removed and a subsequent cholangiogram showed that no filling defects remained. During conventional ERCP in a patient without anatomical anomalies, the patient is placed in the left lateral decubitus position with the endoscopist on the left side of the table [1]. There have been a few reports of successful cases where modifications of the conventional ERCP technique have been used [2,3]. These have included alterations in the position of the patient prior to the procedure, during the procedure, and/or alteration in the position of the endoscopist [1,4]. Our case demonstrates that a skilled endoscopist can successfully carry out ERCP while maintaining a patient with situs inversus in the prone position without using a mirror-image technique or resorting to laparotomy [5].

Endoscopy\_UCTN\_Code\_TTT\_1AR\_2AB

Competing interests: None

**Kumkum Sarkar Patel<sup>1</sup>, Jay Nitin Patel<sup>1</sup>,  
Siddharth Mathur<sup>2</sup>, Yitzchak Moshe-  
nyat<sup>3</sup>**

<sup>1</sup> Department of Internal Medicine,  
Winthrop-University Hospital, Mineola,  
New York, USA

<sup>2</sup> Department of Gastroenterology,  
Brooklyn Hospital Center, Brooklyn,  
New York, USA

<sup>3</sup> Department of Gastroenterology,  
Lutheran Medical Center, Brooklyn,  
New York, USA

## References

- 1 Chowdhury A, Chatterjee BK, Das U et al. ERCP in situs inversus: do we need to turn the other way? *Indian J Gastroenterol* 1997; 16: 155–156
- 2 Venu RP, Geenen JE, Hogan WJ et al. ERCP and endoscopic sphincterotomy in patients with situs inversus. *Gastrointest Endosc* 1985; 31: 338–340
- 3 McDermott JP, Caushaj PF. ERCP and laparoscopic cholecystectomy for cholangitis in a 66-year-old male with situs inversus. *Surg Endosc* 1994; 8: 1227–1229
- 4 Nordback I, Airo I. ERCP and endoscopic papillotomy in complete abdominal situs inversus. *Gastrointest Endosc* 1988; 34: 150
- 5 García-Fernández FJ, Infantes JM, Torres Y et al. ERCP in complete situs inversus viscerum using a “mirror image” technique. *Endoscopy* 2010; 42 (Suppl. 02): E316–E317

## Bibliography

**DOI** <http://dx.doi.org/10.1055/s-0034-1377213>  
*Endoscopy* 2014; 46: E304–E305  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

## Corresponding author

**Kumkum Sarkar Patel, MD, MPH**  
260 First Street, Apt. B13  
Mineola  
NY 11501  
USA  
[kumkum.sarkar@gmail.com](mailto:kumkum.sarkar@gmail.com)  
[kspatel@winthrop.org](mailto:kspatel@winthrop.org)