


## Extensive squamous metaplasia in the rectum stained with a Lugol's solution in patient with ulcerative colitis and primary sclerosing cholangitis



**Fig. 1** A whitish epithelium in the distal rectum adjacent to the inflamed rectal mucosa.



**Fig. 2** The rectal mucosa and the squamous metaplasia stained with Lugol's solution.

A case of extensive squamous metaplasia in a patient with longstanding ulcerative colitis and primary sclerosing cholangitis is presented. The patient was a 52-year-old man with a 20-year history of pancolitis who was referred for surveillance colonoscopy. No suspicious lesions had been found on previous examinations. Inspection of the rectum revealed a whitish epithelium lining the distal portion of the bowel, which was almost complete in circumferential extent and approximately 2 cm long at its maximum extent (► Fig. 1). Lugol's solution, known for its interaction with glycogen in a squamous epithelium, was used as a stain. An irregular border between the rectal mucosa and the metaplastic epithelium was clearly demarcated, and details of the surface of the latter could be easily observed (► Fig. 2). As expected, histopathology confirmed the presence of squamous epithelium in the rectum. Although extremely rare, squamous metaplasia and squamous carcinoma of the rectum have been observed in longstanding ulcerative colitis [1–3]. In addition to chronic inflammation, human papilloma virus infection may play a role in the metaplasia–dysplasia–carcinoma sequence [1]. To the best of our knowledge, the combination of ulcerative colitis, squamous metaplasia, and primary sclerosing cholangitis has not been previously described.

Endoscopy\_UCTN\_Code\_CCL\_1AD\_2AD

**Competing interests:** None

**Edyta Zagórowicz<sup>1,2</sup>,  
Andrzej Paweł<sup>1,2</sup>,  
Anna Rawa-Golebiewska<sup>2</sup>,  
Janina Orłowska<sup>3</sup>**

<sup>1</sup> Department of Gastroenterological Oncology, The Maria Skłodowska-Curie Memorial Cancer Centre and Institute of Oncology, Warsaw, Poland

<sup>2</sup> Department of Gastroenterology, Hepatology and Clinical Oncology, Medical Centre for Postgraduate Education, Warsaw, Poland

<sup>3</sup> Department of Pathology, The Maria Skłodowska-Curie Memorial Cancer Centre and Institute of Oncology, Warsaw, Poland

### References

- 1 Cheng H, Sitrin MD, Satchidanand SK et al. Colonic squamous cell carcinoma in ulcerative colitis: report of a case and review of the literature. *Can J Gastroenterol* 2007; 21: 47–50
- 2 Fu K, Tsujinaka Y, Hamahata Y et al. Squamous metaplasia of the rectum associated with ulcerative colitis diagnosed using narrow-band imaging. *Endoscopy* 2008; 40 (Suppl. 02): E45–46
- 3 Lord JD, Upton MP, Hwang JH. Confocal endomicroscopic evaluation of colorectal squamous metaplasia and dysplasia in ulcerative colitis. *Gastrointest Endosc* 2011; 73: 1064–1066

### Bibliography

**DOI** <http://dx.doi.org/10.1055/s-0034-1377374>  
Endoscopy 2014; 46: E377  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

### Corresponding author

**Edyta Zagórowicz, MD**  
Department of Gastroenterology  
The Maria Skłodowska-Curie Memorial Cancer  
Centre and Institute of Oncology  
5 Roentgen Street  
02-781 Warsaw  
Poland  
Fax: +48-22-5463035  
ezagorowicz@wp.pl