


F. XU,* M. J. ZACUTO,* Y. KOHMURA, J. ROSEN, A. GIBB, M. ALAM, J. SCOTT,
D. TSCHAEN (MERCK RESEARCH LABORATORIES, RAHWAY, USA AND MERCK SHARP &
DOHME RESEARCH LABORATORIES HODDESDON, UK)
Asymmetric Synthesis of Highly Functionalized Tetrahydropyran DPP-4 Inhibitor
Org. Lett. **2014**, *16*, 5422–5425.

Asymmetric Synthesis of a DPP-4 Inhibitor



Significance: The target tetrahydropyran DPP-4 inhibitor was of interest for the treatment of type 2 diabetes. The synthesis depicted features three tandem ruthenium-catalyzed reactions: (1) an asymmetric transfer hydrogenation of ketone **A** with dynamic kinetic resolution (2) a cycloisomerization to form a dihydropyran ring and (3) an oxidation. The overall yield of the synthesis is 25%.

Comment: Extensive optimization of the asymmetric transfer hydrogenation established that significant contributors to the yield, dr and er included the use of the pentafluoro-substituted DAIPEN catalyst **B**, DABCO as the base and THF as the solvent. The reductive amination of ketone **I** with NaBH(OAc)₃ dramatically improved (dr = 19:1) using DMAc as solvent when the bis(tosylate) salt **J** was neutralized with Et₃N followed by pH buffering with HOAc.

SYNFACTS Contributors: Philip Kocienski
Synfacts 2015, 11(1), 0003 Published online: 15.12.2014
DOI: 10.1055/s-0034-1379643; Reg-No.: K05714SF

2015 © THIEME STUTTGART • NEW YORK

Category

Synthesis of Natural Products and Potential Drugs

Key words

DPP-4 inhibitors

asymmetric transfer hydrogenation

dynamic kinetic resolution

cycloisomerization

reductive amination

SYNFACTS
of the month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.