

Z. ZHANG,* W. ZHANG, J. LI, Q. LIU, T. LIU, G. ZHANG* (HENAN NORMAL UNIVERSITY, XINXIANG, P. R. OF CHINA)

Synthesis of Multisubstituted Pyrroles from Doubly Activated Cyclopropanes Using an Iron-Mediated Oxidation Domino Reaction

J. Org. Chem. **2014**, *79*, 11226–11233.

Iron-Mediated Synthesis of Pyrroles from Cyclopropanes

Category


Synthesis of Heterocycles

Key words

pyrroles

cyclopropanes

iron catalysis



$\text{R}^1 = \text{Me, Ph, 4-MeOC}_6\text{H}_4$

$\text{R}^2 = \text{OEt, 4-MeC}_6\text{H}_4\text{NH, 4-ClC}_6\text{H}_4\text{NH, 4-MeOC}_6\text{H}_4\text{NH}$

$\text{R}^3 = \text{Cy, } n\text{-Bu, Ph, 4-MeC}_6\text{H}_4, 2\text{-MeC}_6\text{H}_4, 3\text{-MeC}_6\text{H}_4, 4\text{-ClC}_6\text{H}_4, 2\text{-ClC}_6\text{H}_4, 3\text{-ClC}_6\text{H}_4, 4\text{-MeOC}_6\text{H}_4, \text{C}_6\text{H}_4\text{CH}_2, 4\text{-MeC}_6\text{H}_4\text{CH}_2$
 $\text{R}^4 = \text{H, Me}$

Significance: The synthesis of highly substituted pyrroles **3** from cyclopropanes **1** and amines **2** via an iron-mediated sequential ring opening–cyclization–dehydrogenation reaction is reported. The conditions were optimized using different solvents, reaction times, and iron catalysts. The scope was studied and cyclopropanes **1** bearing methyl and aryl substituents at R^1 were tested with the latter giving better yields. EWG- and EDG-substituted aryl amines as well as $\text{R}^2 = \text{OEt}$ were well tolerated. Also, a methyl group in the cyclopropane ($\text{R}^4 = \text{Me}$) was suitable, furnishing 1,2,3,5-substituted pyrrole **3** in good yield. A series of aromatic and aliphatic amines were screened as well: aliphatic amines furnished pyrroles **3** in low yields, while aromatic amines gave better results, although the yields decrease according to the position of the phenyl substituent. A reaction mechanism involving a radical process was suggested based on radical trapping experiments.

Comment: Cyclopropane derivatives can be used as precursors to synthesize a variety of useful heterocyclic motifs (C. A. Carson, M. A. Kerr *Chem. Soc. Rev.* **2009**, *38*, 3051; J. R. Green, V. Snieckus *Synlett* **2014**, *25*, 2258). The present work reports an efficient synthesis of tri- and tetra-substituted pyrroles in moderate to good yields from readily available cyclopropanes and amines. The reaction shows a broad substrate scope; both EWG and EDG in **1** and **2** affording pyrroles **3** in comparable yields. While anilines with different patterns of substitution furnished **3** in good yields, a large excess of amine was required when benzylamines were used. The readily available cyclopropanes can be easily prepared in high yields by the reaction of a dicarbonyl compound with 1,2-dibromoethane (Z. Zhang et al. *Angew. Chem. Int. Ed.* **2007**, *46*, 1726).

SYNFACTS Contributors: Victor Snieckus, Livia Cristina Frota
Synfacts 2015, 11(1), 0023 Published online: 15.12.2014
DOI: 10.1055/s-0034-1379704; **Reg-No.:** V15114SF

2015 © THIEME STUTTGART • NEW YORK