


A Mukaiyama–Claisen Approach to 3,5-Diketo Esters

Qinggang Wang
Benjamin List*

Max-Planck-Institut für Kohlenforschung, Kaiser-Wilhelm-Platz 1,
45470 Mülheim an der Ruhr, Germany
list@mpi-muelheim.mpg.de

Dedicated to ‘uncle Peter’, in deep admiration for his marvelous
contributions to chemical synthesis, including *Synlett*.



Received: 02.12.2014
Accepted after revision: 14.01.2015
Published online: 10.02.2015
DOI: 10.1055/s-0034-1380145; Art ID: st-2014-d0995-l

Abstract A thermally promoted synthesis of 3,5-diketo esters via a Mukaiyama–Claisen reaction of 4H-1,3-dioxin-4-one derivatives with silyl enolates has been developed. The desired oligocarbonyl compounds were obtained with moderate to good yields.

Key words 3,5-diketo esters, Mukaiyama–Claisen reaction, thermal reaction, silylated nucleophiles

Oligocarbonyl compounds, especially 3,5-diketo ester derivatives, are important building blocks for the synthesis of polyketides and polyols, such as erythromycin A, rosuvastatin, tetracycline and wailupemycin F (Figure 1).^{1–3}





Figure 1 Representative examples of polyketides and polyols

Therefore, approaches towards their synthesis are of interest. Several methodologies have previously been developed to synthesize oligocarbonyl compounds.⁴ Among them, Claisen condensation approaches have proven efficient in building up the carbon backbone. The Langer group recently reported the reaction of silylated nucleophiles with acyl chloride to give 3,5-diketo esters.^{4a–d} Tanabe and co-workers utilized a similar strategy to access analogous 1,3-dicarbonyl compounds.^{4e,f} However, to our knowledge, esters themselves have not been used as electrophiles in Mukaiyama-type Claisen condensations. Notably, 4H-1,3-dioxin-4-one derivatives are useful synthetic equivalents for 1,3-dicarbonyls. In the context of a different study, we have recently described a single example in which a 3,5-diketo ester was accessed by reacting an enol silane with a 4H-1,3-dioxin-4-one derivative.^{5,6} Herein, we explore the generality of this unique transformation and report an improved thermal approach for the synthesis of 3,5-diketo esters via a Mukaiyama–Claisen reaction of 4H-1,3-dioxin-4-one derivatives with silyl ketene acetals.

Under optimized conditions both aliphatic and α,β -unsaturated 4H-1,3-dioxin-4-one derivatives could be employed in the thermally promoted Mukaiyama–Claisen reaction. Simply treating compounds **1** with an excess of silyl ketene acetals **2a** or **2b**, gave the corresponding 3,5-diketo esters in good yields after desilylation (Table 1). Using an *N*-Boc carbamate containing substrate, the corresponding tricarbonyl products **3a** and **3b** were obtained in 85% and 78% yield, respectively. With an *O*-TBS protected substrate, the reactions proceeded smoothly, giving **3c** and **3d** in 85% and 84% yield, respectively. Reactions involving a conjugated alkenyl substrate required higher temperature (110 °C), and the corresponding products **3e** and **3f** were obtained in 61% and 73% yield, respectively. Using the simple, methyl-substituted substrate, the desired tricarbonyl compounds **3g** and **3h** were obtained in 73% and 75% yield, respectively.

Table 1 Substrate Scope of the Mukaiyama–Claisen Reaction


Entry ^a	Product	Yield (%)
1	3a 	85
2	3b 	78
3	3c 	85
4	3d 	84
5 ^b	3e 	61
6 ^b	3f 	73
7	3g 	73
8	3h 	75

^a Reactions were carried out with **1** (0.1 mmol) and **2** (0.4 mmol) in toluene (1 mL) for 4–5 h at 90 °C.

^b Reactions were carried out at 110 °C.

Interestingly, when a silyl enol ether (**2c**) was explored as the nucleophile, an intramolecular cyclization was observed and 2-methyl-6-phenyl-4H-pyran-4-one (**4**) was isolated in 59% yield (Scheme 1). Presumably, under these conditions, a triketone is initially formed and subsequently undergoes cyclization to heterocycle **4**.⁷

In summary, a thermally promoted synthesis of 3,5-diketone esters via a Mukaiyama–Claisen reaction of 4H-1,3-dioxin-4-one derivatives with silylated enolate nucleophiles has been developed.⁸ The desired oligocarbonyl compounds were obtained in good yields. This methodology may find applications in the synthesis of bioactive polyketides or polyols.

**Scheme 1** Intramolecular cyclization observed using silyl enol ether **2c** as nucleophile

Acknowledgment

Generous support by the Max-Planck-Society and the European Research Council (Advanced Grant ‘High Performance Lewis Acid Organocatalysis, HIPOCAT’) is gratefully acknowledged.


Supporting Information

Supporting information for this article is available online at <http://dx.doi.org/10.1055/s-0034-1380145>.

References and Notes

- (a) Koskinen, A. M. P.; Karisalmi, K. *Chem. Soc. Rev.* **2005**, *34*, 677. (b) Aginagalde, M.; Bello, T.; Masdeu, C.; Vara, Y.; Arrieta, A.; Cossío, F. P. *J. Org. Chem.* **2010**, *75*, 7435. (c) Shimada, N.; Stewart, C.; Bow, W. F.; Jolit, A.; Wong, K.; Zhou, Z.; Tius, M. A. *Angew. Chem. Int. Ed.* **2012**, *51*, 5727. (d) Pandit, R. P.; Lee, Y. R. *Org. Biomol. Chem.* **2014**, *12*, 4407.
- (a) Himmelsbach, M.; Lintvedt, R. L.; Zehetmair, J. K.; Nanny, M.; Heeg, M. J. *J. Am. Chem. Soc.* **1987**, *109*, 8003. (b) Chizhov, D. L.; Sevenard, D. V.; Lork, E.; Pashkevich, K. I.; Roschenthaler, G.-V. *J. Fluorine Chem.* **2004**, *125*, 1137. (c) Miyatake-Onozabal, H.; Barrett, A. G. M. *Org. Lett.* **2010**, *12*, 5573. (d) Jiang, J.-A.; Huang, W.-B.; Zhai, J.-J.; Liu, H.-W.; Cai, Q.; Xu, L.-X.; Wang, W.; Ji, Y.-F. *Tetrahedron* **2013**, *69*, 627.
- (a) Work, S. T.; Hauser, C. R. *J. Org. Chem.* **1963**, *28*, 725. (b) Bartoli, G.; Bosco, M.; Cimarelli, C.; Dalpozzo, R.; Guercio, G.; Palmieri, G. *J. Chem. Soc., Perkin. Trans. 1* **1993**, 2081. (c) Sánchez-Larios, E.; Thai, K.; Bilodeau, F.; Gravel, M. *Org. Lett.* **2011**, *13*, 4942. (d) Kalaitzis, J. A.; Cheng, Q.; Thomas, P. M.; Kelleher, N. L.; Moore, B. S. *J. Nat. Prod.* **2009**, *72*, 469.
- (a) Rahn, T.; Nguyen, V. T. N.; Dang, T. H. T.; Ahmed, Z.; Methling, K.; Lalk, M.; Fischer, C.; Spannenberg, A.; Langer, P. *J. Org. Chem.* **2007**, *72*, 1957. (b) Rahn, T.; Dang, T. H. T.; Spannenberg, A.; Fischer, C.; Langer, P. *Org. Biomol. Chem.* **2008**, *6*, 3366. (c) Rahn, T.; Appel, B.; Baumann, W.; Jiao, H.; Börner, A.; Fischer, C.; Langer, P. *Org. Biomol. Chem.* **2009**, *7*, 1931. (d) Büttner, S.; Desens, W.; Michalik, D.; Langer, T. *Eur. J. Org. Chem.* **2011**, 6663. (e) Iliada, A.; Takai, K.; Okabayashi, T.; Misaki, T.; Tanabe, Y. *Chem. Commun.* **2005**, 3171. (f) Iliada, A.; Osada, J.; Nagase, R.; Misaki, T.; Tanabe, Y. *Org. Lett.* **2007**, *9*, 1859. (g) Reynolds, P. W.; Deloach, J. A. *J. Am. Chem. Soc.* **1984**, *106*, 4566.

- (5) (a) Wang, Q.; van Gemmeren, M.; List, B. *Angew. Chem. Int. Ed.* **2014**, *53*, 13592. (b) Sridharan, V.; Ruiz, M.; Menndez, J. C. *Synthesis* **2010**, 1053. (c) Roh, E. J.; Keller, J. M.; Olah, Z.; Iadarola, M. J.; Jacobson, K. A. *Bioorg. Med. Chem.* **2008**, *16*, 9349. (d) Toueg, J.; Prunet, J. *Synlett* **2006**, 2807.
- (6) An acyl ketene intermediate has recently been proposed in thermal reactions of 4*H*-1,3-dioxin-4-ones (Scheme 2):



- (7) (a) Arimoto, H.; Nishiyama, S.; Yamamura, S. *Tetrahedron Lett.* **1990**, *31*, 5619. (b) Lovell, S.; Subramony, P.; Kahr, B. *J. Am. Chem. Soc.* **1999**, *121*, 7020. (c) Paterson, I.; Chen, D. Y.-K.; Aceña, J. L.; Franklin, A. S. *Org. Lett.* **2000**, *2*, 1513. (d) Sakakura, A.; Watanabe, H.; Nakagawa, S.; Ishihara, K. *Chem. Asian J.* **2007**, *2*, 477. (e) Sakakura, A.; Watanabe, H.; Ishihara, K. *Org. Lett.* **2008**, *10*, 2569.
- (8) A cooling-finger-capped Schleck tube with a stirring bar was charged with the corresponding 4*H*-1,3-dioxin-4-one derivative **1** (0.1 mmol), silylated nucleophile **2** (0.4 mmol), and anhydrous toluene (1 mL). The resulting mixture was heated to 90 °C or 110 °C for 4–5 h until full consumption of the starting material was observed (monitored by TLC). The solvent was removed under reduced pressure, MeOH (1 mL) and KHF₂ (10 equiv) were added, and the reaction mixture was stirred at r.t. for 10 min. After filtering through a thin layer of Celite, the residue was purified by column chromatography (isohexanes–EtOAc, 5:1 to 1:1).