

Cytokine-Suppressive Activity of a Hydroxylated Alkylamide from *Echinacea purpurea*

Martha Leyte-Lugo¹, Daniel A. Todd¹, Travis V. Gulledge², Monika Juzumaite², Frederick S. Carter¹, Scott M. Laster², Nadja B. Cech¹

¹ Department of Chemistry and Biochemistry, The University of North Carolina at Greensboro, Greensboro, NC, United States

² Department of Biological Sciences, North Carolina State University, Raleigh, NC, United States

Abstract

Echinacea purpurea has been widely used to treat upper respiratory tract infections. Bioassay-guided fractionation of hexane and ethanol extracts of this plant yielded 4[(2-methylbutyl)amino]-4-oxo-2-butenoic acid (**1**) and 8,11-dihydroxy-dodeca-2*E*,4*E*,9*E*-trienoic acid isobutylamide (**2**), both of which are new to *Echinacea*. Compound **2** suppressed the production of TNF- α from RAW 264.7 cells with an IC₅₀ value of 6.8 μ M, while **1** was inactive. Neither compound showed cytotoxicity at concentrations up to 100 μ M. These findings support a growing body of literature demonstrating that *E. purpurea* contains an array of compounds that suppress cytokine secretion by macrophage-type cells *in vitro*.

Key words

Echinacea purpurea · Asteraceae · alkylamides · alkamides · anti-inflammatory · tumor necrosis factor- α

Supporting information available online at <http://www.thieme-connect.de/products>

Preparations from *Echinacea purpurea* (L.) Moench (Asteraceae) are used in the treatment and prevention of upper respiratory infections [1]. The mechanisms that underlie the purported efficacy of *E. purpurea* have been subject to debate. However, recent studies have indicated that some *E. purpurea* extracts and constituents possess *in vitro* activity consistent with an anti-inflammatory effect [2, 3]. Along these lines, our group has shown that *E. purpurea* root extracts suppress the secretion of inflammatory cytokines by cultured immune cells. The activity of the extracts could not entirely be explained by the presence of known compounds [4, 5]. Thus, we sought to isolate and identify additional compounds with cytokine-suppressive activity from *E. purpurea*. Bioassay-guided fractionation of *E. purpurea* root extracts yielded two alkylamides new to *E. purpurea* (Fig. 1). From a hexane extract, 4[(2-methylbutyl)amino]-4-oxo-2-butenoic acid (**1**) was isolated, while a mixture of diastereoisomers (as indicated by LC-MS, Fig. 1 S, Supporting Information), 8,11-dihydroxy-dodeca-2*E*,4*E*,9*E*-trienoic acid isobutylamide (**2**), was obtained from an aqueous ethanol extract. Compound **1** is commercially available as a synthetic compound (Aldlab Chemicals, Woburn, MA; Aurora Building Blocks, San Diego, CA), but its characterization has not been described and it has not, to our knowledge, been reported in the literature. Compound **2** is a mixture of diastereoisomers, for which the NMR spectroscopic data matches




Fig. 1 Alkylamides from the roots of *E. purpurea*. Two compounds were isolated and identified, 4[(2-methylbutyl)amino]-4-oxo-2-butenoic acid (**1**) and 8,11-dihydroxy-dodeca-2*E*,4*E*,9*E*-trienoic acid isobutylamide (**2**). The structure of an additional known alkylamide from *E. purpurea*, dodeca-2*E*,4*E*-dienoic acid isobutylamide (**3**), is shown for comparison.

that of a known constituent of *Spilanthes callimorpha* A.H.Moore (Asteraceae), 8,11-dihydroxy-dodeca-2*E*,4*E*,9*E*-trienoic acid isobutylamide [6]. Interestingly, although at least 17 different alkylamides have been reported from *E. purpurea* [4, 7–12], **1** and **2** are the first carboxylated and hydroxylated alkylamides, respectively, to be identified from this species. Importantly, although **1** and **2** were originally isolated and purified from extracts of different polarity (hexane and aqueous ethanol, respectively), both of these compounds were detected in both the hexane and aqueous ethanol crude extracts (Fig. 2 S, Supporting Information). The presence of **1** and **2** in the aqueous ethanol extract is particularly interesting given that it is common to use aqueous ethanol extracts of *E. purpurea* as dietary supplements. However, it should also be noted that **1** and **2** appear, based on LC-MS data (Fig. 2 S, Supporting Information), to be present only at relatively low levels compared to other alkylamides.

Compound **1** was obtained as a yellow powder with a molecular formula of C₉H₁₅NO₃ at *m/z* 186.1126 (calcd. for [M + H]⁺, C₉H₁₆NO₃, 186.11302), indicating three degrees of unsaturation. Spectral data (¹H and ¹³C NMR, Table 1) revealed that the compound is an α -unsaturated N-methylbutylamide. Characteristic ¹H-NMR signals at δ 3.10 (dd, H-1'), 1.56 (m, H-2'), and ~0.88 (H-4' and H-5') further supported a methylbutylamide moiety, as did the HMBC correlations observed from H-1' to C-1, C-3', and C-5' (Fig. 2). Signals of olefinic protons at δ 6.92 (d, H-2) and 6.64 (d, H-3) confirmed the presence of one double bond conjugated with the amide carbonyl. The *E* configuration was assigned based on the coupling constants (*J* = 15.6). The terminal carboxylic acid was confirmed with the HMBC correlations from H-2 to C-4, and from H-3 to C-1 and C-4 (Fig. 2).

Table 1 NMR spectroscopic data^a (400 MHz for ¹H and 100 MHz for ¹³C) for 4[(2-methylbutyl)amino]-4-oxo-2-butenic acid (**1**).

Position	δ_{H} , m (J in Hz)	δ_{C}
1		166.6
2	6.92, d (15.6)	137.1
3	6.64, d (15.2)	132.3 ^b
4		168.1
1'	3.10, dd (7.2, 6.0)	46.4
2'	1.56, m	36.1
3'	1.39, m	28.1
1.13, m		
4'	0.87, t (7.6)	11.6
5'	0.88, d (6.4)	17.5

^a ¹H and ¹³C chemical shifts with reference to methanol-*d*₄ (δ_{H} = 3.31 ppm) and methanol-*d*₄ (δ_{C} = 49.01 ppm), respectively; ^b Obtained from HSQC




Fig. 2 HMBC correlations of **1**.

The inflammatory response is mediated, in part, by the secretion of proinflammatory cytokines by macrophages. Thus, to explore potential anti-inflammatory activity of **1** and **2**, we evaluated their influence on the secretion of the cytokine tumor necrosis factor- α (TNF- α) by RAW 264.7 macrophage-type cells *in vitro*. To mimic an inflammatory state, these cells were concomitantly treated with lipopolysaccharide (LPS), which induces the production of TNF- α . No significant suppression of LPS-induced TNF- α secretion was observed for **1**, while a dose-dependent suppressive effect was observed for **2** (IC_{50} = 6.8 μM) (● Fig. 3). The potency of **2** was higher than that of the positive control, dodeca-2E,4E-dienoic acid isobutylamide (**3**). Notably, the activity observed in ● Fig. 3 is not attributable to cytotoxic effects. The three

compounds tested were shown to have very little toxicity up to a concentration of 100 μM in the lactate dehydrogenase (LDH) assay (Table 1 S, Supporting Information).

Materials and Methods

Optical rotations of pure compounds were recorded in MeOH on a Rudolph Research Autopol (II) Polarimeter. NMR was conducted using a JEOL ECS-400 NMR spectrometer operating at 400 MHz for ¹H and 100 MHz for ¹³C. High-resolution electrospray ionization mass spectrometry (HRESIMS) was performed on a Thermo LTQ Orbitrap XL mass spectrometer with an electrospray source. A Varian ProStar HPLC with a photodiode array detector was used with a Phenomenex Gemini-NX preparative C₁₈ column (5 μm ; 250 \times 21.2 mm). Flash chromatography was performed on a Teledyne ISCO CombiFlash[®]R_f using 120 g RediSep RF Gold[®] silica columns (20–40 μm particle size). All reagents and solvents were obtained from Fisher Scientific.

The roots of *E. purpurea* were purchased from Pacific Botanicals in Grants Pass, Oregon and species was verified by Nadja B. Cech. In support of the correct assignment of the genus and species, the alkylamide profile for the samples under investigation, as reported in [5] and indicated in Fig. 2 S, Supporting Information, was consistent with that observed for *E. purpurea*. The alkylamide profile differs among *Echinacea* species [7], and is, therefore, a useful indicator for species identification. A voucher specimen (NCU633811) is deposited in the University of North Carolina Herbarium. Two extracts were prepared from these roots, one in hexane and the other in ethanol:water (75:25), as described previously [5].

The hexane extract (4.9 g) was fractionated via flash chromatography with a hexane/chloroform/methanol gradient (flow rate 20 mL/min) to yield 11 primary fractions (FI-FXI). Fraction VII (42 mg) was subjected to purification with reverse-phase preparative HPLC with acetonitrile/water following a gradient of 30–90% acetonitrile over 20 min. Compound **1** (4[(2-methylbutyl)amino]-4-oxo-2-butenic acid) eluted at 7 min (1.0 mg, 96% purity, Fig. 3 S, Supporting Information).

Residue from the ethanolic extract was fractionated into 13 fractions according to previously described methods [5]. Fractions 9




Fig. 3 Influence of **1** and **2** on TNF- α secretion by murine RAW 264.7 macrophage-type cells. Included as a positive control is the *Echinacea* isobutylamide dodeca-2E,4E-dienoic acid isobutylamide (**3**), which has been previously shown to suppress TNF- α secretion in this assay [4]. The data presented are the means of two independent experiments, with error bars representing the standard deviation. Percent of control was determined according to the formula: compound treatment (ng/mL)/LPS treatment (ng/mL) \times 100. Compounds **1–3** were shown not to be cytotoxic up to a concentration of 100 μM (see Table 1 S, Supporting Information).

and 10 were combined based on a similar bioactivity and LC-MS profile for a combined yield of 747 mg. This combined sample was then separated with a hexane/chloroform/methanol gradient (flow rate 40 mL/min) and pooled into five fractions (A–E). Fraction C (330 mg) was subjected to the third stage of separation with a hexane/ethyl acetate/methanol gradient (flow rate 30 mL/min) and pooled into three fractions (I–III). Fraction II (50 mg) was then subjected to further purification with an acetonitrile/water gradient using reversed-phase preparative HPLC (40–50% acetonitrile over 20 min). Compound **2**, 8,11-dihydroxy-dodeca-2E,4E,9E-trienoic acid isobutylamide (1.9 mg, 97% purity as mixture of diastereomers, **Fig. 1 S**, Supporting Information) eluted at 6.4 min.

4[(2-methylbutyl)amino]-4-oxo-2-butenic acid (**1**): pale yellow amorphous powder; $[\alpha]_D^{25} = +2.4$ (c 0.0025, MeOH); ^1H (400 MHz) and ^{13}C NMR (100 MHz) (**Table 1**; **Figs. 4 S** and **5 S**, Supporting Information); HRESIMS m/z 186.1126 $[\text{M} + \text{H}]^+$ (calcd. for $\text{C}_9\text{H}_{16}\text{NO}_3$ 186.1125).

8,11-dihydroxy-dodeca-2E,4E,9E-trienoic acid isobutylamide (**2**): light yellow oil; ^1H (400 MHz) and ^{13}C NMR (100 MHz) data were consistent with the literature (**Table 2 S**, **Figs. 9 S** and **10 S**, Supporting Information) [8]; HRESIMS m/z 282.2056 $[\text{M} + \text{H}]^+$ (calcd. for $\text{C}_{16}\text{H}_{28}\text{NO}_3$ 282.2064).

Cytokine assays were performed using RAW 264.7 cells in 96-well plates. TNF- α secretion was stimulated with LPS from *Salmonella minnesota* R595 (List Biological Laboratories) at 10 ng/mL. TNF- α levels were measured in supernatants after 18 h using an ELISA kit (R&D Systems). Dodeca-2E,4E-dienoic acid isobutylamide (**3**) (Chromadex) served as a positive control. To evaluate cytotoxicity, RAW 264.7 cells were seeded into 96-well plates, treated for 18 h, and the supernatants were analyzed for LDH activity using a commercially available kit (Thermo Fisher Scientific).

Supporting information

NMR spectra, chromatograms for **1** and **2** and the crude *E. purpurea* extracts, and cytotoxicity results are available as Supporting Information.

Acknowledgements

Support for this research was provided by grant #1R15AT007259 from the National Center for Complementary and Alternative Medicine, a component of the National Institutes of Health. We thank Tyler N. Graf and Dr. Nicholas H. Oberlies for technical assistance.

Conflict of Interest

The authors report no conflict of interest.

References

- 1 Gruenewald, J, Brendler T, Jaenicke C. PDR for herbal medicines. Montvale: Medical Economics Co.; 1998: 261–266
- 2 Zhai Z, Solcoc A, Wud L, Wurtele ES, Kohute ML, Murphy PA, Cunnick JE. *Echinacea* increases arginase activity and has anti-inflammatory properties in RAW 264.7 macrophage cells indicative of alternative macrophage activation. *J Ethnopharmacol* 2009; 122: 76–85
- 3 Sharma SM, Anderson M, Schoop SR, Hudson JB. Bactericidal and anti-inflammatory properties of a standardized *Echinacea* extract (Echinaforce): dual actions against respiratory bacteria. *Phytomedicine* 2010; 17: 563–568
- 4 Cech NB, Kandhi V, Davis JM, Hamilton A, Eads D, Laster SM. *Echinacea* and its alkylamides: effects on the influenza A-induced secretion of cytokines, chemokines, and PGE2 from RAW 264.7 macrophage-like cells. *Int Immunopharmacol* 2010; 10: 1268–1278
- 5 Todd DA, Gullidge TV, Britton ER, Oberhofer M, Leyte-Lugo M, Moody A, Shymanovich T, Grubbs LF, Juzumaita M, Graf TN, Oberlies NH, Faeth SH, Laster SM, Cech NB. Ethanolic *Echinacea purpurea* extracts contain a mixture of cytokine-suppressive and cytokine-inducing compounds, including some that originate from endophytic bacteria. *PLoS One* 2015; 10: e0124276
- 6 Li GP, Shen BC, Zhao JF, Yang XD, Li L. Two new alkamides from *Spiranthes callimorpha*. *J Integr Plant Biol* 2007; 49: 1608–1610
- 7 Bauer R, Remiger P. TLC and HPLC analysis of alkamides in *Echinacea* drugs. *Planta Med* 1989; 55: 367–371
- 8 Binns SE, Livesey JF, Arnason JT, Baum BR. Phytochemical variation in *Echinacea* from roots and flowerheads of wild and cultivated populations. *J Agric Food Chem* 2002; 50: 3673–3687
- 9 Clifford LJ, Nair MG, Rana J, Dewitt DL. Bioactivity of alkamides isolated from *Echinacea purpurea* (L.) Moench. *Phytomedicine* 2002; 9: 249–253
- 10 Chen Y, Fu T, Tao T, Yang J, Chang Y, Wang M, Kim L, Qu L, Cassidy J, Scalzo R, Wang X. Macrophage activating effects of new alkamides from the roots of *Echinacea* species. *J Nat Prod* 2005; 68: 773–776
- 11 Cech NB, Eleazer MS, Shoffner LT, Crosswhite MR, Davis AC, Mortenson AM. High performance liquid chromatography/electrospray ionization mass spectrometry for simultaneous analysis of alkamides and caffeic acid derivatives from *Echinacea purpurea* extracts. *J Chromatogr A* 2006; 1103: 219–228
- 12 Spelman K, Wetschler MH, Cech NB. Comparison of alkylamide yield in ethanolic extracts prepared from fresh versus dry *Echinacea purpurea* utilizing HPLC-ESI-MS. *J Pharm Biomed Anal* 2009; 49: 1141–1149

received June 10, 2015

revised

accepted June 23, 2015

Bibliography

DOI <http://dx.doi.org/10.1055/s-0035-1557764>
Planta Med Lett 2015; 2: e25–e27

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 2199-157X

Correspondence

Nadja B. Cech

Department of Chemistry and Biochemistry
The University of North Carolina at Greensboro
301 Mclver St.
Greensboro, NC 24702
United States
Phone: +1 336 334 3017
Fax: +1 336 334 5402
nadja_cech@uncg.edu

License terms

