


Y. GAO, P. TANG, H. ZHOU, W. ZHANG, H. YANG, N. YAN, G. HU, D. MEI, J. WANG,* D. MA* (PEKING UNIVERSITY, BEIJING, ZHEJIANG UNIVERSITY OF TECHNOLOGY, HANGZHOU, ISRAEL CHEMICALS LTD., SHANGHAI, P. R. OF CHINA; NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE; AND PACIFIC NORTHWEST NATIONAL LABORATORY, RICHLAND, USA)

Graphene Oxide Catalyzed C–H Bond Activation: The Importance of Oxygen Functional Groups for Biaryl Construction

Angew. Chem. Int. Ed. **2016**, *55*, 3124–3128.

Graphene Oxide Promoted C–H Arylation of Benzene with Aryl Halides



Significance: Graphene oxide promoted the C–H arylation of benzene with aryl halides in the presence of *t*-BuOK to give the corresponding biaryls in ≤92% GC yield (15 examples).

Comment: The graphene oxide was characterized by TEM, AFM, Raman spectroscopy, XPS, SEM, and BET analyses. In the reaction of benzene with 4-iodoanisole, the catalytic activity of graphene oxide was superior to that of the other carbon materials (carbon nanotubes: 19% GC yield, active carbon: 11%, carbon black: 19%, natural graphite: 8%).